

EVALUATION OF MOBILE SOURCE EMISSIONS AND TRENDS USING DETAILED CHEMICAL AND PHYSICAL MEASUREMENTS

Robert Harley (harley@ce.berkeley.edu)

Department of Civil and Environmental Engineering
University of California, Berkeley

Mar 4, 2014

EPA STAR Project Meeting in Ann Arbor (OTAQ)

Acknowledgments




- UC Berkeley: Tim Dallmann, Drew Gentner, Arthur Chan, Allen Goldstein, Gabriel Isaacman, Steven DeMartini, Brian McDonald, and Dave Worton.
- Aerodyne: Ezra Wood, Tim Onasch, Scott Herndon, John Franklin, Ed Fortner, Doug Worsnop
- LBNL: Tom Kirchstetter, Kevin Wilson
- Research funding:
 - ▣ US Environmental Protection Agency (Grant # RD834553)

A Highway Tunnel Laboratory

Vehicle emissions measured at Caldecott tunnel in SF Bay area:


⊙ Light-Duty Gasoline: 1994-97, 1999, 2001, 2004, 2006, 2010

⊙ Heavy-Duty Diesel Trucks: 1996-97, 2006, 2010




Pollutant	Tunnel Measurement Method
CO ₂	Infrared absorption
Nitric Oxide (NO)	Chemiluminescence
NO ₂ , CO HCHO, C ₂ H ₄	Tunable infrared laser spectroscopy
PM mass & composition	Aerosol mass spectrometer
Black Carbon (BC)	Aethalometer
Light absorption & scattering (532 nm)	Photoacoustic spectrometer and reciprocal nephelometer
Light absorption (630 nm)	Multi-angle absorption photometer
Light extinction (630 nm)	Cavity attenuation phase-shift

On-Road NO_x Emission Factor Trends




McDonald et al. (JGR 2012)

Fuel Sales Trends, 1990-2010




McDonald et al. (JGR 2012)

National On-Road NO_x Emission Trends



McDonald et al. (JGR 2012)


Aerosol Mass Spectrometer (SP-AMS)




- Heated tungsten vaporizer combined with laser to vaporize organic *and* refractory aerosol (e.g., soot)
- Both vaporizers on at all times
- Operate in fast MS mode to capture individual truck plumes

Onasch et al.
(AS&T 2012)

Sample AMS Data – Diesel Truck Plume



Capturing Individual Truck Exhaust Plumes





Chemical
speciation of
exhaust particles,
including trace
elements

Independent
measurements of
NO and NO₂


Peak in CO₂
denotes capture of
exhaust plume

HDDT Emission Factor Distributions




Dallmann et al.
(ACPD 2014)

Cumulative Contributions to Total Emissions from Heavy-Duty Diesel Trucks



Dallmann et al.
(ES&T 2012)

OA mass spectra similar for Gasoline and Diesel




Dallmann et al.
(ACPD 2014)

GC-MS Analysis of Organic Aerosol

Previous GC-MS analyses of vehicular OA emissions typically identify only a small fraction (~5%) of total mass.



We analyzed tunnel OA by photoionization mass spectrometry using vacuum ultraviolet (VUV) photons instead of electron ionization (EI).




Contacts: Allen Goldstein (UCB) & Kevin Wilson (Lawrence Berkeley National Lab)



Electron Ionization (EI) versus Vacuum Ultraviolet (VUV) Ionization




Sample GC-MS Results for Tunnel OA



Worton et al. (ES&T, in review)


Chemical Composition of Tunnel OA



Worton et al. (ES&T, in review)



Diesel Fuel Speciation

(Gentner et al. PNAS 2012)




Gasoline and Diesel and SOA Yields

(Gentner et al. PNAS 2012)



Diesel Contribution to On-Road Emissions

Stabilized Running Emissions – as of 2010



Dallmann et al.
(ES&T 2013)

Summary

- On-road engines are important air pollution source
 - ▣ In 2010, diesel was dominant on-road source of BC, POA, and NO_x
 - ▣ Emission factor distributions are becoming **increasingly** skewed
 - High-emitting tail of distribution responsible for majority of running emissions

- Novel approaches used to characterize emissions
 - ▣ Aerosol Mass Spectrometer (SP-AMS)
 - BC, OA, zinc and phosphorus (lube oil additives) measured in individual truck plumes
 - POA mass spectra **very similar** for gasoline & diesel engine emissions & lube oil
 - ▣ GC-MS analysis using Vacuum Ultraviolet (VUV) photons
 - EI analysis (70 eV) of diesel and lube oil leads to near-total fragmentation of parent molecular ions, and leaves most of the emitted HC mass unidentified (“UCM”)
 - use of softer (9-10.5 eV) photo-ionization preserves molecular ions; **greatly enhances** ability to identify and quantify organics present in diesel fuel and vehicle emissions
 - SOA yield per unit mass of diesel fuel emitted is ~6X higher than gasoline yield

Publications

- Dallmann et al. (2012). On-Road Measurements of Gas and Particle Phase Pollutant Emission Factors for Individual Heavy-Duty Diesel Trucks. *Environ. Sci. Technol.* **46**, 8511–8518.
- Dallmann et al. (2013). Quantifying On-Road Emissions from Gasoline-Powered Motor Vehicles: Accounting for the Presence of Medium- and Heavy-Duty Diesel Trucks. *Environ. Sci. Technol.* **47**, 13873-13881.
- Dallmann et al. (2014). Characterization of particulate matter emissions from on-road gasoline and diesel vehicles using a soot particle aerosol mass spectrometer. *Atmos. Chem. Phys. Disc.* **14**, 4007-4049.
- Gentner et al. (2012). Elucidating Secondary Organic Aerosol from Diesel and Gasoline Vehicles Through Detailed Characterization of Organic Carbon Emissions. *PNAS* **109**, 18318-18323.


Publications



- Gentner et al. (2013). Chemical Composition of Gas-Phase Organic Carbon Emissions from Motor Vehicles and Implications for Ozone Production. *Environ. Sci. Technol.* **47**, 11837-11848.
- McDonald et al. (2012). Long-Term Trends in Nitrogen Oxide Emissions from Motor Vehicles at National, State, and Air Basin Scales. *Journal of Geophysical Research* **117**, D00V18, doi: 10.1029/2012JD018304.
- Worton et al. (2014). Lubricating Oil Dominates Primary Organic Aerosol Emissions from Motor Vehicles. *Environ. Sci. Technol.*, in review.

Measured Tunnel PM Concentrations

(Teflon & Quartz Filters, SP-AMS Data)



Dallmann et al.
(ACPD 2014)