

**APPENDIX A-1b**

**CHEMICAL AND PHYSICAL PROPERTIES AND  
TOXICITY REFERENCE LEVELS**

**TABLE A-1b-1**

**CHEMICAL-SPECIFIC INPUTS FOR  
ACENAPHTHYLENE (208-96-8)**

| <b>Parameter</b>                       | <b>Reference and Explanation</b>                                  | <b>Value</b> |
|--|---|--------------|
| <b>Chemical/Physical Properties</b>    |   |              |
| $MW$ (g/mole)                          | CRC Handbook (1995)   | 152.20       |
| $T_m$ (K)                              | CRC Handbook (1995)   | 365.65       |
| $V_p$ (atm)                            | --  | ND           |
| $S$ (mg/L)                             | --  | ND           |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value was cited from CRC Handbook (1995).                     | 8.29E-05     |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 4.39E-02     |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 7.53E-06     |
| $K_{ow}$ (unitless)                    | --  | ND           |
| $K_{oc}$ (mL/g)                        | --  | ND           |
| $Kd_s$ (cm <sup>3</sup> /g)            | --  | ND           |
| $Kd_{sw}$ (L/Kg)                       | --  | ND           |
| <b>Dermal Exposure Factors</b>         |   |              |
| $K_p$ (cm/hr)                          | $K_p$ value was obtained from U.S. EPA (1992b).                   | 1.55E-01     |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).                     | 7.42E-01     |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).                   | 6.23E+00     |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).                     | 1.00E+00     |
| <b>Biotransfer Factors for Animals</b> |   |              |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --  | ND           |
| $BAF_{fish}$ (L/kg FW)                 | --  | ND           |

**TABLE A-1b-1****CHEMICAL-SPECIFIC INPUTS FOR  
ACENAPHTHYLENE (208-96-8)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997b)  | 6.00E-02     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | U.S. EPA (1997b)  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 2.1E-01      |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | U.S. EPA (1997c)  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

**TABLE A-1b-2**

**CHEMICAL-SPECIFIC INPUTS FOR  
ACRYLAMIDE (79-06-1)**

| <b>Parameter</b>                           | <b>Reference and Explanation</b>  | <b>Value</b> |
|--|---|--------------|
| <b>Chemical/Physical Properties</b>        |   |              |
| <i>MW</i> (g/mole)                         | Budavari, O'Neil, Smith, and Heckelman (1989)   | 71.08        |
| <i>T<sub>m</sub></i> (K)                   | Budavari, O'Neil, Smith, and Heckelman (1989)   | 357.65       |
| <i>V<sub>p</sub></i> (atm)                 | <i>V<sub>p</sub></i> value cited in U.S. EPA (1995g)  | 9.20E-06     |
| <i>S</i> (g/100ml H <sub>2</sub> O)        | Geometric mean value cited in U.S. EPA (1994c).   | 2.15E+02     |
| <i>H</i> (atm·m <sup>3</sup> /mol)         | <i>H</i> value cited in U.S. EPA (1995g)  | 3.00E-10     |
| <i>D<sub>a</sub></i> (cm <sup>2</sup> /s)  | <i>D<sub>a</sub></i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 9.70E-02     |
| <i>D<sub>w</sub></i> (cm <sup>2</sup> /s)  | <i>D<sub>w</sub></i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 1.06E-05     |
| <i>K<sub>ow</sub></i> (unitless)           | --  | 1.10E-01     |
| <i>K<sub>oc</sub></i> (mL/g)               | --  | 1.10E-01     |
| <i>Kd<sub>s</sub></i> (cm <sup>3</sup> /g) | <i>Kd<sub>s</sub></i> value was calculated by using the correlation equation with <i>K<sub>oc</sub></i> that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate <i>Kd<sub>s</sub></i> , because the value varies, depending on the fraction of organic carbon in soil. Recommended <i>Kd<sub>s</sub></i> value was calculated by using the <i>K<sub>oc</sub></i> value that is provided in this table.   | 1.10E-03     |
| <i>Kd<sub>sw</sub></i> (L/Kg)              | <i>Kd<sub>sw</sub></i> value was calculated by using the correlation equation with <i>K<sub>oc</sub></i> that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate <i>Kd<sub>sw</sub></i> , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended <i>Kd<sub>sw</sub></i> value was calculated by using the <i>K<sub>oc</sub></i> value that is provided in this table. | 9.00E-03     |
| <b>Dermal Exposure Factors</b>             |   |              |
| <i>Kp</i> (cm/hr)                          | <i>Kp</i> value was obtained from U.S. EPA (1992b).   | 1.50E-04     |
| <i>t</i>                                   | <i>t</i> value was obtained from U.S. EPA (1992b).  | 2.40E-01     |
| <i>t*</i>                                  | <i>t*</i> value was obtained from U.S. EPA (1992b).   | 5.70E-01     |
| <i>B</i>                                   | <i>B</i> value was obtained from U.S. EPA (1992b).  | 1.10E-05     |
| <b>Biotransfer Factors for Animals</b>     |   |              |
| <i>BCF<sub>fish</sub></i> (L/kg FW tissue) | <i>BCF<sub>s</sub></i> were used for compounds with a log <i>K<sub>ow</sub></i> value below 4.0, as cited in U.S. EPA (1995g). <i>BCF<sub>fish</sub></i> value calculated using the correlation equation with <i>K<sub>ow</sub></i> obtained from Veith, Macek, Petrocelli, and Caroll (1980)   | 7.70E-02     |
| <i>BAF<sub>fish</sub></i> (L/kg FW)        | --  | NA           |

**TABLE A-1b-2**

**CHEMICAL-SPECIFIC INPUTS FOR  
ACRYLAMIDE (79-06-1)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S.EPA (1997b)   | 2.00E-04     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | U.S.EPA (1997b)   | 4.50E+00     |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 7.00E-04     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | U.S.EPA (1997b)   | 4.55E+00     |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

**TABLE A-1b-3**

**CHEMICAL-SPECIFIC INPUTS FOR  
ALLYL CHLORIDE (107-05-1)**

| <b>Parameter</b>                           | <b>Reference and Explanation</b>  | <b>Value</b> |
|--|---|--------------|
| <b>Chemical/Physical Properties</b>        |   |              |
| <i>MW</i> (g/mole)                         | Montgomery and Welkom (1991)  | 76.53        |
| <i>T<sub>m</sub></i> (K)                   | Montgomery and Welkom (1991)  | 138.65       |
| <i>V<sub>p</sub></i> (atm)                 | <i>V<sub>p</sub></i> value cited in U.S. EPA (1995g).   | 4.80E-01     |
| <i>S</i> (mg/L)                            | <i>S</i> value cited in U.S. EPA (1995g).   | 3.40E+03     |
| <i>H</i> (atm·m <sup>3</sup> /mol)         | <i>H</i> value cited in U.S. EPA (1995g)  | 1.10E-02     |
| <i>D<sub>a</sub></i> (cm <sup>2</sup> /s)  | <i>D<sub>a</sub></i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 1.17E-01     |
| <i>D<sub>w</sub></i> (cm <sup>2</sup> /s)  | <i>D<sub>w</sub></i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 1.08E-05     |
| <i>K<sub>ow</sub></i> (unitless)           | <i>K<sub>ow</sub></i> value cited in U.S. EPA (1995g).  | 2.80E+01     |
| <i>K<sub>oc</sub></i> (mL/g)               | <i>K<sub>oc</sub></i> value cited in U.S. EPA (1995g).  | 2.70E+01     |
| <i>Kd<sub>s</sub></i> (cm <sup>3</sup> /g) | <i>Kd<sub>s</sub></i> value was calculated by using the correlation equation with <i>K<sub>oc</sub></i> that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate <i>Kd<sub>s</sub></i> , because the value varies, depending on the fraction of organic carbon in soil. Recommended <i>Kd<sub>s</sub></i> value was calculated by using the <i>K<sub>oc</sub></i> value that is provided in this table.   | 2.70E-01     |
| <i>Kd<sub>sw</sub></i> (L/Kg)              | <i>Kd<sub>sw</sub></i> value was calculated by using the correlation equation with <i>K<sub>oc</sub></i> that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate <i>Kd<sub>sw</sub></i> , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended <i>Kd<sub>sw</sub></i> value was calculated by using the <i>K<sub>oc</sub></i> value that is provided in this table. | 2.02E+00     |
| <b>Dermal Exposure Factors</b>             |   |              |
| <i>Kp</i> (cm/hr)                          | <i>Kp</i> value was obtained from U.S. EPA (1992b).   | 7.00E-03     |
| <i>t</i>                                   | <i>t</i> value was obtained from U.S. EPA (1992b).  | 2.60E-01     |
| <i>t*</i>                                  | <i>t*</i> value was obtained from U.S. EPA (1992b).   | 6.20E-01     |
| <i>B</i>                                   | <i>B</i> value was obtained from U.S. EPA (1992b).  | 2.80E-03     |
| <b>Biotransfer Factors for Animals</b>     |   |              |
| <i>BCF<sub>fish</sub></i> (L/kg FW tissue) | <i>BCF<sub>s</sub></i> were used for compounds with a log <i>K<sub>ow</sub></i> value below 4.0, as cited in U.S. EPA (1995g). <i>BCF<sub>fish</sub></i> value calculated using the correlation equation with <i>K<sub>ow</sub></i> obtained from Veith, Macek, Petrocelli, and Caroll (1980)   | 3.70E+00     |
| <i>BAF<sub>fish</sub></i> (L/kg FW)        | --  | NA           |

**TABLE A-1b-3**

**CHEMICAL-SPECIFIC INPUTS FOR  
ALLYL CHLORIDE (107-05-1)**

| <b>Parameter</b>  | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>  |   |              |
| <i>RfD</i> (mg/kg/day)  | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 2.86E-04     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>                           | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                                     | --  | 1.0E-03      |
| <i>Inhalation</i> <sup>1</sup> <i>CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>  | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)  | --  | ND           |

Note: NA = Not applicable, ND = No data available

**TABLE A-1b-4**

**CHEMICAL-SPECIFIC INPUTS FOR  
BENZIDINE (92-87-5)**

| Parameter                                  | Reference and Explanation   | Value    |
|--|---|----------|
| <b>Chemical/Physical Properties</b>        |   |          |
| <i>MW</i> (g/mole)                         | Budavari, O'Neil, Smith, and Heckelman (1989)   | 184.23   |
| <i>T<sub>m</sub></i> (K)                   | Budavari, O'Neil, Smith, and Heckelman (1989)   | 390.65   |
| <i>V<sub>p</sub></i> (atm)                 | --  | ND       |
| <i>S</i> (g/2500ml)                        | Geometric mean value cited in U.S. EPA (1994c).   | 1.0      |
| <i>H</i> (atm·m <sup>3</sup> /mol)         | <i>H</i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 3.88E-11 |
| <i>D<sub>a</sub></i> (cm <sup>2</sup> /s)  | <i>D<sub>a</sub></i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-02 |
| <i>D<sub>w</sub></i> (cm <sup>2</sup> /s)  | <i>D<sub>w</sub></i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 1.50E-05 |
| <i>K<sub>ow</sub></i> (unitless)           | --  | 4.60E+01 |
| <i>K<sub>oc</sub></i> (mL/g)               | --  | 4.30E+01 |
| <i>Kd<sub>s</sub></i> (cm <sup>3</sup> /g) | <i>Kd<sub>s</sub></i> value was calculated by using the correlation equation with <i>K<sub>oc</sub></i> that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate <i>Kd<sub>s</sub></i> , because the value varies, depending on the fraction of organic carbon in soil. Recommended <i>Kd<sub>s</sub></i> value was calculated by using the <i>K<sub>oc</sub></i> value that is provided in this table.   | 4.30E-01 |
| <i>Kd<sub>sw</sub></i> (L/Kg)              | <i>Kd<sub>sw</sub></i> value was calculated by using the correlation equation with <i>K<sub>oc</sub></i> that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate <i>Kd<sub>sw</sub></i> , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended <i>Kd<sub>sw</sub></i> value was calculated by using the <i>K<sub>oc</sub></i> value that is provided in this table. | 3.22E+00 |
| <b>Dermal Exposure Factors</b>             |   |          |
| <i>Kp</i> (cm/hr)                          | <i>Kp</i> value was obtained from U.S. EPA (1992b).   | 2.20E-03 |
| <i>t</i>                                   | <i>t</i> value was obtained from U.S. EPA (1992b).  | 1.20E+00 |
| <i>t*</i>                                  | <i>t*</i> value was obtained from U.S. EPA (1992b).   | 2.80E+00 |
| <i>B</i>                                   | <i>B</i> value was obtained from U.S. EPA (1992b).  | 4.60E-03 |
| <b>Biotransfer Factors for Animals</b>     |   |          |
| <i>BCF<sub>fish</sub></i> (L/kg FW tissue) | <i>BCFs</i> were used for compounds with a log <i>K<sub>ow</sub></i> value below 4.0, as cited in U.S. EPA (1995g). <i>BCF<sub>fish</sub></i> value calculated using the correlation equation with <i>K<sub>ow</sub></i> obtained from Veith, Macek, Petrocelli, and Carroll (1980)   | 3.00E+00 |
| <i>BAF<sub>fish</sub></i> (L/kg FW)        | --  | NA       |



**TABLE A-1b-4**

**CHEMICAL-SPECIFIC INPUTS FOR  
BENZIDINE (92-87-5)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S.EPA (1997b)   | 3.00E-03     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | 2.30E+02     |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 1.10E-02     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | 2.35E+02     |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | U.S. EPA (1995f)  | 2.5E+01      |

Note: NA = Not applicable, ND = No data available

**TABLE A-1b-5**

**CHEMICAL-SPECIFIC INPUTS FOR  
BENZO(GHI)PERYLENE (191-24-2)**

| <b>Parameter</b>                       | <b>Reference and Explanation</b>   | <b>Value</b> |
|--|--|--------------|
| <b>Chemical/Physical Properties</b>    |  |              |
| $MW$ (g/mole)                          | CRC Handbook (1995)  | 276.34       |
| $T_m$ (°K)                             | --   | ND           |
| $V_p$ (atm)                            | --   | ND           |
| $S$ (mg/L)                             | --   | ND           |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value is assumed to be zero, because the $V_p$ and $S$ values are zero for all metals, except mercury. | 7.40229E-07  |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database in U.S. EPA (1994f).                                       | 2.01E-02     |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database in U.S. EPA (1994f).                                       | 5.26E-06     |
| $K_{ow}$ (unitless)                    | --   | ND           |
| $K_{oc}$ (mL/g)                        | --   | ND           |
| $Kd_s$ (mL/g)                          | --   | ND           |
| <b>Dermal Exposure Factors</b>         |  |              |
| $K_p$ (cm/hr)                          | $K_p$ value was obtained from U.S. EPA (1992b).  | 1.62E+00     |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 4.24E+00     |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 2.00E+01     |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 3.16E+02     |
| <b>Biotransfer Factors for Animals</b> |  |              |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --   | ND           |
| $BAF_{fish}$ (L/kg FW)                 | --   | ND           |

**TABLE A-1b-5**

**CHEMICAL-SPECIFIC INPUTS FOR  
BENZO(GHI)PERYLENE (191-24-2)**

| <b>Parameter</b>                                   | <b>Reference and Explanation</b>             | <b>Value</b> |
|--|--|--------------|
| <b>Health Benchmarks</b>                           |  |              |
| <i>RfD</i> (mg/kg/day)                             | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>          | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                    | --   | ND           |
| <i>Inhalation CSF</i><br>(mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>   | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                                 | --   | ND           |

Note: NA = Not applicable, ND = No data available

**TABLE A-1b-6**

**CHEMICAL-SPECIFIC INPUTS FOR  
BIS-(2-CHLOROETHOXY) METHANE (111-91-1)**

| <b>Parameter</b>                              | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Chemical/Physical Properties</b>           |   |              |
| <i>MW</i> (g/mole)                            | Montgomery and Welkom (1990)  | 173.04       |
| <i>T<sub>m</sub></i> (°K)                     | Montgomery and Welkom (1990)  | 240.35       |
| <i>V<sub>p</sub></i> (1mm @53 °C)             | Montgomery and Welkom (1990)  | 1.0          |
| <i>S</i> (mg/L@ 25 °C)                        | All metals, except mercury, are assumed to be insoluble in water.   | 81,000.00    |
| <i>H</i> (atm·m <sup>3</sup> /mol)            | U.S. EPA (1998c)  | 3.78E-07     |
| <i>D<sub>a</sub></i> (cm <sup>2</sup> /s)     | <i>D<sub>a</sub></i> value was calculated using the equation cited in U.S. EPA (1996a).   | 3.20E-02     |
| <i>D<sub>w</sub></i> (cm <sup>2</sup> /s)     | <i>D<sub>w</sub></i> value was calculated using the equation cited in U.S. EPA (1996a).   | 8.46E-06     |
| <i>K<sub>ow</sub></i> (unitless)              | Montgomery and Welkom (1990)  | 1.82E+01     |
| <i>K<sub>oc</sub></i> (mL/g)                  | Montgomery and Welkom (1990)  | 1.14E+02     |
| <i>Kd<sub>s</sub></i> (mL/g)                  | <i>Kd<sub>s</sub></i> value was calculated by using the correlation equation with <i>K<sub>oc</sub></i> that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate <i>Kd<sub>s</sub></i> , because the value varies, depending on the fraction of organic carbon in soil. Recommended <i>Kd<sub>s</sub></i> value was calculated by using the <i>K<sub>oc</sub></i> value that is provided in this table.   | 1.14E+00     |
| <i>Kd<sub>sw</sub></i> (L/Kg)                 | <i>Kd<sub>sw</sub></i> value was calculated by using the correlation equation with <i>K<sub>oc</sub></i> that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate <i>Kd<sub>sw</sub></i> , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended <i>Kd<sub>sw</sub></i> value was calculated by using the <i>K<sub>oc</sub></i> value that is provided in this table. | 8.55E+00     |
| <b>Dermal Exposure Factors</b>                |   |              |
| <i>K<sub>p</sub></i> (cm/hr)                  | <i>K<sub>p</sub></i> value was obtained from U.S. EPA (1992b).  | 131E-03      |
| <i>t</i>                                      | <i>t</i> value was obtained from U.S. EPA (1992b).  | 9.95E-01     |
| <i>t*</i>                                     | <i>t*</i> value was obtained from U.S. EPA (1992b).   | 2.39E+00     |
| <i>B</i>                                      | <i>B</i> value was obtained from U.S. EPA (1992b).  | 1.82E-03     |
| <b>Biotransfer Factors for Animals</b>        |   |              |
| <i>BCF<sub>fish</sub></i><br>(L/kg FW tissue) | --  | ND           |
| <i>BAF<sub>fish</sub></i> (L/kg FW)           | --  | ND           |

**TABLE A-1b-6****CHEMICAL-SPECIFIC INPUTS FOR  
BIS-(2-CHLOROETHOXY) METHANE (111-91-1)**

| <b>Parameter</b>                                   | <b>Reference and Explanation</b>             | <b>Value</b> |
|--|--|--------------|
| <b>Health Benchmarks</b>                           |  |              |
| <i>RfD</i> (mg/kg/day)                             | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>          | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                    | --   | ND           |
| <i>Inhalation CSF</i><br>(mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>   | National Primary Drinking Water Regulations. |              |
| Aquatic TRV (µg/l)                                 | --   | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-7

**CHEMICAL-SPECIFIC INPUTS FOR  
BUTANOL (71-36-3)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | U.S. EPA (1995g)   | 74.12    |
| $T_m$ (K)                              | --   |          |
| $Vp$ (atm)                             | U.S. EPA (1995g)   | 8.60E-03 |
| $S$ (mg/L)                             | U.S. EPA (1995g)   | 7.50E+04 |
| $H$ (atm·m <sup>3</sup> /mol)          | U.S. EPA (1995g)   | 8.81E-06 |
| $D_a$ (cm <sup>2</sup> /s)             | U.S. EPA (1995g)   | 8.00E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | U.S. EPA (1995g)   | 9.30E-06 |
| $K_{ow}$ (unitless)                    | U.S. EPA (1995g)   | 6.30E+00 |
| $K_{oc}$ (mL/g)                        | U.S. EPA (1995g)   | 6.10E+00 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 6.10E-02 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 4.58E-01 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 2.50E-03 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 2.50E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 5.90E-01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 6.30E-04 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$<br>(L/kg FW tissue)       | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 1.20E+00 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-7****CHEMICAL-SPECIFIC INPUTS FOR  
BUTANOL (71-36-3)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997c)  | 1.00E-01     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 3.50E-01     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

**TABLE A-1b-8**

**CHEMICAL-SPECIFIC INPUTS FOR  
BUTYL-4,6-DINITROPHENOL, 2-SEC (DINOSEB) (88-85-7)**

| <b>Parameter</b>                       | <b>Reference and Explanation</b>   | <b>Value</b>     |
|--|--|------------------|
| <b>Chemical/Physical Properties</b>    |  |                  |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 240.22           |
| $T_m$ (K)                              | U.S. EPA (1995b)   | 311.15 to 414.15 |
| $V_p$ (atm)                            | U.S. EPA (1995g)   | 9.90E-05         |
| $S$ (mg/L)                             | U.S. EPA (1995g)   | 5.20E+01         |
| $H$ (atm·m <sup>3</sup> /mol)          | U.S. EPA (1995g)   | 4.56E-07         |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-02         |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-06         |
| $K_{ow}$ (unitless)                    | U.S. EPA (1995g)   | 1.40E+03         |
| $K_{oc}$ (mL/g)                        | U.S. EPA (1995g)   | 1.20E+02         |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 1.20E+00         |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 9.00E+00         |
| <b>Dermal Exposure Factors</b>         |  |                  |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 1.10E-02         |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 2.60E+00         |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 8.30E+00         |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 1.40E-01         |
| <b>Biotransfer Factors for Animals</b> |  |                  |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 8.00E+01         |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA               |



**TABLE A-1b-8**

**CHEMICAL-SPECIFIC INPUTS FOR  
BUTYL-4,6-DINITROPHENOL, 2-SEC (DINOSEB) (88-85-7)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997c)  | 1.00E-03     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 3.50E-03     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-9

**CHEMICAL-SPECIFIC INPUTS FOR  
2-CHLORO-1,3-BUTADIENE (CHLOROPRENE) (126-99-8)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | U.S. EPA (1995g)   | 88.54    |
| $T_m$ (K)                              | --   |          |
| $Vp$ (atm)                             | U.S. EPA (1995g)   | 2.80E-01 |
| $S$ (mg/L)                             | U.S. EPA (1995g)   | 6.30E+02 |
| $H$ (atm·m <sup>3</sup> /mol)          | --   | ND       |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d)   | 1.04E-01 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d)   | 1.00E-05 |
| $K_{ow}$ (unitless)                    | U.S. EPA (1995g)   | 1.20E+02 |
| $K_{oc}$ (mL/g)                        | U.S. EPA (1995g)   | 1.10E+02 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table. 5g)   | 1.10E+00 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 8.25E+00 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 1.60E-02 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 3.00E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 7.30E-01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 1.20E-02 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 1.10E+01 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-9****CHEMICAL-SPECIFIC INPUTS FOR  
2-CHLORO-1,3-BUTADIENE (CHLOROPRENE) (126-99-8)**

| <b>Parameter</b>                                   | <b>Reference and Explanation</b>             | <b>Value</b> |
|--|--|--------------|
| <b>Health Benchmarks</b>                           |  |              |
| <i>RfD</i> (mg/kg/day)                             | U.S. EPA (1997b)                             | 2.00E-02     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>          | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                    | U.S. EPA (1997b)                             | 2.00E-03     |
| <i>Inhalation CSF</i><br>(mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>   | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                                 | --   | ND           |

Note: NA = Not applicable, ND = No data available

**TABLE A-1b-10**

**CHEMICAL-SPECIFIC INPUTS FOR  
CHROMIUM (+3) (16065-38-1)**

| <b>Parameter</b>                       | <b>Reference and Explanation</b>  | <b>Value</b>   |
|--|---|--|
| <b>Chemical/Physical Properties</b>    |   |  |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)   | 51.996   |
| $T_m$ (K)                              | Budavari, O'Neil, Smith, and Heckelman (1989)   | 2173.15  |
| $V_p$ (atm)                            | --  | 0  |
| $S$ (mg/L)                             | --  | 0  |
| $H$ (atm·m <sup>3</sup> /mol)          | --  | 0  |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d)  | 1.01E-01   |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d)  | 4.63E-05   |
| $K_{ow}$ (unitless)                    | --  | NA   |
| $K_{oc}$ (mL/g)                        | --  | NA   |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was obtained from U.S. EPA (1996a), which provides pH-based values that were estimated by using the MINTEQ2 geochemical speciation model.                    | 1.2E+03 at<br>pH=4.9;<br>1.8E+06 at<br>pH=6.8;<br>4.3E+06 at<br>pH=8.0 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value is assumed to be the same as the $Kd_s$ value, because organic carbon does not play a major role in sorption for the metals, as cited in US. EPA (1994e). | 1.2E+03 at<br>pH=4.9;<br>1.8E+06 at<br>pH=6.8;<br>4.3E+06 at<br>pH=8.0 |
| <b>Dermal Exposure Factors</b>         |   |  |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).  | 1.00E-03   |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).   | ND   |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).   | ND   |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).   | ND   |
| <b>Biotransfer Factors for Animals</b> |   |  |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --  | ND   |
| $BAF_{fish}$ (L/kg FW)                 | --  | ND   |

**TABLE A-1b-10****CHEMICAL-SPECIFIC INPUTS FOR  
CHROMIUM (+3) (16065-38-1)**

| <b>Parameter</b>                                   | <b>Reference and Explanation</b>             | <b>Value</b> |
|--|--|--------------|
| <b>Health Benchmarks</b>                           |  |              |
| <i>RfD</i> (mg/kg/day)                             | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>          | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                    | --   | ND           |
| <i>Inhalation CSF</i><br>(mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>   | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                                 | --   | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-11

**CHEMICAL-SPECIFIC INPUTS FOR  
CIS-1,3-DICHLOROPROPENE (10061-01-5)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 110.97   |
| $T_m$ (°K)                             | --   |          |
| $Vp$ (atm)                             | U.S. EPA (1995g)   | 4.99E-02 |
| $S$ (mg/L)                             | U.S. EPA (1995g)   | 2.70E+03 |
| $H$ (atm·m <sup>3</sup> /mol)          | U.S. EPA (1995g)   | 1.76E-03 |
| $D_a$ (cm <sup>2</sup> /s)             | U.S. EPA (1995g)   | 5.85E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | U.S. EPA (1995g)   | 1.10E-05 |
| $K_{ow}$ (unitless)                    | U.S. EPA (1995g)   | 1.00E+02 |
| $K_{oc}$ (mL/g)                        | U.S. EPA (1995g)   | 9.30E+01 |
| $Kd_s$ (mL/g)                          | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 9.30E-01 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 6.97E+00 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 1.10E-02 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 4.20E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 1.00E+00 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 1.00E-02 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW)                 | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 5.30E+00 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-11**

**CHEMICAL-SPECIFIC INPUTS FOR  
CIS-1,3-DICHLOROPROPENE (10061-01-5)**

| <b>Parameter</b>                                   | <b>Reference and Explanation</b>             | <b>Value</b> |
|--|--|--------------|
| <b>Health Benchmarks</b>                           |  |              |
| <i>RfD</i> (mg/kg/day)                             | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>          | U.S. EPA (1997c)                             | 1.75E-01     |
| <i>RfC</i> (mg/m <sup>3</sup> )                    | --   | ND           |
| <i>Inhalation CSF</i><br>(mg/kg/day) <sup>-1</sup> | --   | 1.75E-01     |
| <i>MCL</i>   | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                                 | U.S. EPA (1995f)                             | 2.40E+01     |

Note: Not applicable, ND = No data available

**TABLE A-1b-12**

**CHEMICAL-SPECIFIC INPUTS FOR  
COBALT (7440-48-4)**

| <b>Parameter</b>                               | <b>Reference and Explanation</b>                               | <b>Value</b> |
|--|--|--------------|
| <b>Chemical/Physical Properties</b>            |  |              |
| <i>MW</i> (g/mole)                             | Montgomery and Welkom (1991)                                   | 58.93        |
| <i>T<sub>m</sub></i> (K)                       | Montgomery and Welkom (1991)                                   | 1766.15      |
| <i>V<sub>p</sub></i> (atm)                     | --   | NA           |
| <i>S</i> (mg/L)                                | --   | NA           |
| <i>H</i> (atm·m <sup>3</sup> /mol)             | --   | NA           |
| <i>D<sub>a</sub></i> (cm <sup>2</sup> /s)      | --   | NA           |
| <i>D<sub>w</sub></i> (cm <sup>2</sup> /s)      | --   | NA           |
| <i>K<sub>ow</sub></i> (unitless)               | --   | NA           |
| <i>K<sub>oc</sub></i> (mL/g)                   | --   | NA           |
| <i>Kd<sub>s</sub></i> (cm <sup>3</sup> /g)     | --   | NA           |
| <i>Kd<sub>sw</sub></i> (L/Kg)                  | --   | NA           |
| <b>Dermal Exposure Factors</b>                 |  |              |
| <i>K<sub>p</sub></i> (cm/hr)                   | <i>K<sub>p</sub></i> value was obtained from U.S. EPA (1992b). | 1.00E-03     |
| <i>t</i>                                       | <i>t</i> value was obtained from U.S. EPA (1992b).             | ND           |
| <i>t*</i>                                      | <i>t*</i> value was obtained from U.S. EPA (1992b).            | ND           |
| <i>B</i>                                       | <i>B</i> value was obtained from U.S. EPA (1992b).             | ND           |
| <b>Biotransfer Factors for Animals</b>         |  |              |
| <i>BCF<sub>fish</sub></i><br>(L/kg, FW tissue) | --   | ND           |
| <i>BAF<sub>fish</sub></i> (L/kg FW)            | --   | ND           |



**TABLE A-1b-12**

**CHEMICAL-SPECIFIC INPUTS FOR  
COBALT (7440-48-4)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S.EPA (1997b)   | 6.00E-02     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 2.10E-01     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | .   | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-13

**CHEMICAL-SPECIFIC INPUTS FOR  
COPPER (744-050-8)**

| Parameter                              | Reference and Explanation   | Value    |
|--|---|----------|
| <b>Chemical/Physical Properties</b>    |   |          |
| $MW$ (g/mole)                          | Montgomery and Welkom (1991)  | 63.55    |
| $T_m$ (K)                              | Montgomery and Welkom (1991)  | 1356.15  |
| $V_p$ (atm)                            | --  | NA       |
| $S$ (mg/L)                             | --  | NA       |
| $H$ (atm·m <sup>3</sup> /mol)          | --  | NA       |
| $D_a$ (cm <sup>2</sup> /s)             | --  | NA       |
| $D_w$ (cm <sup>2</sup> /s)             | --  | NA       |
| $K_{ow}$ (unitless)                    | --  | NA       |
| $K_{oc}$ (mL/g)                        | --  | NA       |
| $Kd_s$ (cm <sup>3</sup> /g)            | --  | 2.20E+01 |
| $Kd_{sw}$ (L/Kg)                       | --  | 2.20E+01 |
| <b>Dermal Exposure Factors</b>         |   |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).  | 1.00E-03 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).   | ND       |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).   | ND       |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).   | ND       |
| <b>Biotransfer Factors for Animals</b> |   |          |
| $BCF_{fish}$<br>(L/kg, FW tissue)      | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980) | 0.0E+00  |
| $BAF_{fish}$ (L/kg FW)                 | --  | NA       |

**TABLE A-1b-13**

**CHEMICAL-SPECIFIC INPUTS FOR  
COPPER (744-050-8)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S.EPA (1997b)   | 4.00E-02     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 1.40E-01     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | 1.3          |
| Aquatic TRV (µg/l)                              | Ambient Water Quality Criteria, U.S. EPA, Office of Water.  | 9.00E+00     |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-14

CHEMICAL-SPECIFIC INPUTS FOR  
CYCLOTETRAMETHYLENETETRANITRAMINE (HMX) (2691-41-0)

| Parameter                              | Reference and Explanation   | Value    |
|--|---|----------|
| <b>Chemical/Physical Properties</b>    |   |          |
| $MW$ (g/mole)                          | CRC Handbook (1995)   | 296.16   |
| $T_m$ (K)                              | CRC Handbook (1995)   | 559.15   |
| $Vp$ (atm)                             | --  | ND       |
| $S$ (mg/L)                             | --  | ND       |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value was obtained from WATER8 model database (U.S. EPA 1995d).   | 2.60E-15 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from WATER8 model database (U.S. EPA 1995d). | 8.00E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from WATER8 model database (U.S. EPA 1995d). | 8.00E-06 |
| $K_{ow}$ (unitless)                    | --  | ND       |
| $K_{oc}$ (mL/g)                        | --  | ND       |
| $Kd_s$ (cm <sup>3</sup> /g)            | --  | ND       |
| $Kd_{sw}$ (L/Kg)                       | --  | ND       |
| <b>Dermal Exposure Factors</b>         |   |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).                        | 3.28E-05 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).                         | 6.90E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).                       | ND       |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).                         | ND       |
| <b>Biotransfer Factors for Animals</b> |   |          |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --  | ND       |
| $BAF_{fish}$ (L/kg FW)                 | --  | ND       |

**TABLE A-1b-14**

**CHEMICAL-SPECIFIC INPUTS FOR  
CYCLOTETRAMETHYLENETETRANITRAMINE (HMX) (2691-41-0)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997b)  | 5.00E-02     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 1.75E-01     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-15

**CHEMICAL-SPECIFIC INPUTS FOR  
DIALATE (2303-16-4)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 270.24   |
| $T_m$ (K)                              | --   | ND       |
| $V_p$ (atm)                            | $V_p$ value cited in U.S. EPA (1995g).   | 2.00E-07 |
| $S$ (ppm)                              | $S$ value cited in U.S. EPA (1995b).   | 40       |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 4.83E-05 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-06 |
| $K_{ow}$ (unitless)                    | Value cited in U.S. EPA (1995g).   | 3.10E+04 |
| $K_{oc}$ (mL/g)                        | Value cited in U.S. EPA (1995g).   | 2.60E+04 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 2.60E+02 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 1.82E+03 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 6.60E-02 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 3.90E+00 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 2.00E+01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 3.10E+00 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --   | NA       |
| $BAF_{fish}$ (L/kg FW)                 | $BAFs$ were used for compounds with a log $K_{ow}$ value above 4.0, as cited in U.S. EPA (1995g). $BAF$ values were predicted values calculated by multiplying a food chain multiplier ( $FCM$ ) with an estimated $BCF$ . $BCFs$ were estimated using the correlation equation obtained from Veith, Macek, Petrocelli, and Caroll (1980). $FCMs$ were obtained from U.S. EPA (1995g)  | 1.80E+04 |

**TABLE A-1b-15**

**CHEMICAL-SPECIFIC INPUTS FOR  
DIALATE (2303-16-4)**

| <b>Parameter</b>  | <b>Reference and Explanation</b>                                      | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                                    |   |              |
| <i>RfD</i> (mg/kg/day)                                      | --  | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>                   | U.S. EPA (1997c)  | 6.10E-02     |
| <i>RfC</i> (mg/m <sup>3</sup> )                             | --  | ND           |
| <i>Inhalation URF</i><br>(μg/m <sup>3</sup> ) <sup>-1</sup> | --  | ND           |
| <i>Inhalation CSF</i><br>(mg/kg/day) <sup>-1</sup>          | Value based on <i>Oral CSF</i> assuming route-to-route extrapolation. | 6.10E-01     |
| <i>MCL</i>  | National Primary Drinking Water Regulations.                          | ND           |
| Aquatic TRV (μg/l)  | --  | ND           |

Note: NA= Not applicable, ND= No data available

**TABLE A-1b-16**

**CHEMICAL-SPECIFIC INPUTS FOR  
DIBENZOFURAN (132-64-9)**

| <b>Parameter</b>                       | <b>Reference and Explanation</b>                                  | <b>Value</b> |
|--|---|--------------|
| <b>Chemical/Physical Properties</b>    |   |              |
| $MW$ (g/mole)                          | CRC Handbook (1995)   | 168.19       |
| $T_m$ (K)                              | CRC Handbook (1995)   | 359.65       |
| $V_p$ (atm)                            | --  | ND           |
| $S$ (mg/L)                             | --  | ND           |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).   | 1.056E-04    |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 8.00E-02     |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 8.00E-06     |
| $K_{ow}$ (unitless)                    | --  | ND           |
| $K_{oc}$ (mL/g)                        | --  | ND           |
| $Kd_s$ (cm <sup>3</sup> /g)            | --  | ND           |
| $Kd_{sw}$ (L/Kg)                       | --  | ND           |
| <b>Dermal Exposure Factors</b>         |   |              |
| $K_p$ (cm/hr)                          | $K_p$ value was obtained from U.S. EPA (1992b).                   | 2.06E-01     |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).                     | 6.90E-01     |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).                   | ND           |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).                     | 2.04E+00     |
| <b>Biotransfer Factors for Animals</b> |   |              |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --  | ND           |
| $BAF_{fish}$ (L/kg FW)                 | --  | ND           |



**TABLE A-1b-16**

**CHEMICAL-SPECIFIC INPUTS FOR  
DIBENZOFURAN (132-64-9)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S.EPA (1997a)   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | U.S. EPA (1996c)  | 2.00E+1      |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-17

**CHEMICAL-SPECIFIC INPUTS FOR  
2,6-DICHLOROPHENOL (87-65-0)**

| Parameter                              | Reference and Explanation   | Value               |
|--|---|---------------------|
| <b>Chemical/Physical Properties</b>    |   |                     |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)                     | 163.01              |
| $T_m$ (K)                              | Howard (1989-1993)  | 337.65 to<br>338.65 |
| $V_p$ (atm)                            | --  | ND                  |
| $S$ (mg/L)                             | --  | ND                  |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).   | 2.96E-06            |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 3.47E-02            |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 8.77E-06            |
| $K_{ow}$ (unitless)                    | --  | ND                  |
| $K_{oc}$ (mL/g)                        | --  | ND                  |
| $Kd_s$ (cm <sup>3</sup> /g)            | --  | ND                  |
| $Kd_{sw}$ (L/Kg)                       | --  | ND                  |
| <b>Dermal Exposure Factors</b>         |   |                     |
| $K_p$ (cm/hr)                          | $K_p$ value was obtained from U.S. EPA (1992b).                   | 2.07E-02            |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).                     | 8.63E-01            |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).                   | 2.07E+00            |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).                     | 7.24E-02            |
| <b>Biotransfer Factors for Animals</b> |   |                     |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --  | ND                  |
| $BAF_{fish}$ (L/kg FW)                 | --  | ND                  |

**TABLE A-1b-17**

**CHEMICAL-SPECIFIC INPUTS FOR  
2,6-DICHLOROPHENOL (87-65-0)**

| <b>Parameter</b>                                   | <b>Reference and Explanation</b>             | <b>Value</b> |
|--|--|--------------|
| <b>Health Benchmarks</b>                           |  |              |
| <i>RfD</i> (mg/kg/day)                             | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>          | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                    | --   | ND           |
| <i>Inhalation CSF</i><br>(mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>   | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                                 | --   | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-18

**CHEMICAL-SPECIFIC INPUTS FOR  
2,4-DICHLOROPHENOXYACETIC ACID (94-75-7)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | U.S. EPA (1995g)   | 221.04   |
| $T_m$ (K)                              | --   | ND       |
| $Vp$ (atm)                             | U.S. EPA (1995g)   | 1.40E-05 |
| $S$ (mg/L)                             | U.S. EPA (1995g)   | 6.80E+02 |
| $H$ (atm·m <sup>3</sup> /mol)          | U.S. EPA (1995g)   | 4.50E-06 |
| $D_a$ (cm <sup>2</sup> /s)             | U.S. EPA (1995g)   | 5.88E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | U.S. EPA (1995g)   | 6.49E-06 |
| $K_{ow}$ (unitless)                    | U.S. EPA (1995g)   | 5.05E+02 |
| $K_{oc}$ (mL/g)                        | U.S. EPA (1995g)   | 4.50E+02 |
| $Kd_s$ (mL/g)                          | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.--   | 4.50E+00 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 3.37E+01 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 7.10E-03 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 2.00E+00 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 4.70E+00 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 5.00E-02 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$<br>(L/kg FW tissue)       | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 6.10E+01 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-18**

**CHEMICAL-SPECIFIC INPUTS FOR  
2,4-DICHLOROPHENOXYACETIC ACID (94-75-7)**

| <b>Parameter</b>                                   | <b>Reference and Explanation</b>   | <b>Value</b> |
|--|--|--------------|
| <b>Health Benchmarks</b>                           |  |              |
| <i>RfD</i> (mg/kg/day)                             | U.S. EPA (1997c)   | 1.00E-02     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>          | Calculated by multiplying the Oral CSF for Benzo(a)pyrene by the relative potency factor for Dibenz(a,h)anthracene of 1.0 (U.S.EPA 1993e). | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                    | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg.                            | 3.50E-02     |
| <i>Inhalation CSF</i><br>(mg/kg/day) <sup>-1</sup> |  | ND           |
| <i>MCL</i>   | National Primary Drinking Water Regulations.   | ND           |
| Aquatic TRV (µg/l)                                 | --   | ND           |

Note: NA = Not applicable, ND = No data available

**TABLE A-1b-19**

**CHEMICAL-SPECIFIC INPUTS FOR  
1,3-DICHLOROPROPENE-TRANS (10061-02-6)**

| <b>Parameter</b>                           | <b>Reference and Explanation</b>  | <b>Value</b> |
|--|---|--------------|
| <b>Chemical/Physical Properties</b>        |   |              |
| <i>MW</i> (g/mole)                         | Budavari, O'Neil, Smith, and Heckelman (1989)   | 110.98       |
| <i>T<sub>m</sub></i> (K)                   | --  |              |
| <i>V<sub>p</sub></i> (atm)                 | U.S. EPA (1995g)  | 4.00E-02     |
| <i>S</i> (mg/L)                            | U.S. EPA (1995g)  | 2.80E+03     |
| <i>H</i> (atm·m <sup>3</sup> /mol)         | U.S. EPA (1995g)  | 1.25E-03     |
| <i>D<sub>a</sub></i> (cm <sup>2</sup> /s)  | <i>D<sub>a</sub></i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 5.85E-02     |
| <i>D<sub>w</sub></i> (cm <sup>2</sup> /s)  | <i>D<sub>w</sub></i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 1.10E-05     |
| <i>K<sub>ow</sub></i> (unitless)           | U.S. EPA (1995g)  | 1.00E+02     |
| <i>K<sub>oc</sub></i> (mL/g)               | U.S. EPA (1995g)  | 9.30E+01     |
| <i>Kd<sub>s</sub></i> (cm <sup>3</sup> /g) | <i>Kd<sub>s</sub></i> value was calculated by using the correlation equation with <i>K<sub>oc</sub></i> that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate <i>Kd<sub>s</sub></i> , because the value varies, depending on the fraction of organic carbon in soil. Recommended <i>Kd<sub>s</sub></i> value was calculated by using the <i>K<sub>oc</sub></i> value that is provided in this table.   | 9.30E-01     |
| <i>Kd<sub>sw</sub></i> (L/Kg)              | <i>Kd<sub>sw</sub></i> value was calculated by using the correlation equation with <i>K<sub>oc</sub></i> that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate <i>Kd<sub>sw</sub></i> , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended <i>Kd<sub>sw</sub></i> value was calculated by using the <i>K<sub>oc</sub></i> value that is provided in this table. | 6.97E+00     |
| <b>Dermal Exposure Factors</b>             |   |              |
| <i>K<sub>p</sub></i> (cm/hr)               | <i>K<sub>p</sub></i> value was obtained from U.S. EPA (1992b).  | 1.10E-02     |
| <i>t</i>                                   | <i>t</i> value was obtained from U.S. EPA (1992b).  | 4.20E-01     |
| <i>t*</i>                                  | <i>t*</i> value was obtained from U.S. EPA (1992b).   | 1.00E+00     |
| <i>B</i>                                   | <i>B</i> value was obtained from U.S. EPA (1992b).  | 1.00E-02     |
| <b>Biotransfer Factors for Animals</b>     |   |              |
| <i>BCF<sub>fish</sub></i> (L/kg FW tissue) | <i>BCF<sub>s</sub></i> were used for compounds with a log <i>K<sub>ow</sub></i> value below 4.0, as cited in U.S. EPA (1995g). <i>BCF<sub>fish</sub></i> value calculated using the correlation equation with <i>K<sub>ow</sub></i> obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 5.30E+00     |
| <i>BAF<sub>fish</sub></i> (L/kg FW)        | --  | NA           |

**TABLE A-1b-19**

**CHEMICAL-SPECIFIC INPUTS FOR  
1,3-DICHLOROPROPENE-TRANS (10061-02-6)**

| <b>Parameter</b>                                   | <b>Reference and Explanation</b>                                      | <b>Value</b> |
|--|---|--------------|
| <b>Health Benchmarks</b>                           |   |              |
| <i>RfD</i> (mg/kg/day)                             | --  | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>          | U.S. EPA (1997b)  | 1.75E-01     |
| <i>RfC</i> (mg/m <sup>3</sup> )                    | --  | ND           |
| <i>Inhalation CSF</i><br>(mg/kg/day) <sup>-1</sup> | Value based on <i>Oral CSF</i> assuming route-to-route extrapolation. | 1.75E-01     |
| <i>MCL</i>   | National Primary Drinking Water Regulations.                          | ND           |
| Aquatic TRV (µg/l)                                 | U.S. EPA (1995f)  | 2.40E+01     |

Note: NA= Not applicable, ND= No data available

TABLE A-1b-20

**CHEMICAL-SPECIFIC INPUTS FOR  
DIETHYLSTILBESTROL (56-53-1)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 268.34   |
| $T_m$ (K)                              | Budavari, O'Neil, Smith, and Heckelman (1989)  | 443.65   |
| $V_p$ (atm)                            | U.S. EPA (1995g)   | 1.40E-12 |
| $S$ (mg/L)                             | $S$ value cited in U.S. EPA (1995b).   | 1.30E+04 |
| $H$ (atm·m <sup>3</sup> /mol)          | U.S. EPA (1995g)   | 3.00E-14 |
| $D_a$ (cm <sup>2</sup> /s)             | U.S. EPA (1995g)   | 8.00E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | U.S. EPA (1995g)   | 8.00E-06 |
| $K_{ow}$ (unitless)                    | U.S. EPA (1995g)   | 1.20E+05 |
| $K_{oc}$ (mL/g)                        | U.S. EPA (1995g)   | 9.60E+04 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 9.60E+02 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 7.20E+03 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 1.70E-01 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 3.80E+00 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 1.80E+01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 1.20E+01 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$<br>(L/kg, FW tissue)      | --   | NA       |
| $BAF_{fish}$ (L/kg FW)                 | $BAFs$ were used for compounds with a log $K_{ow}$ value above 4.0, as cited in U.S. EPA (1995g). $BAF$ values were predicted values calculated by multiplying a food chain multiplier ( $FCM$ ) with a geometric mean of various laboratory measured $BCFs$ obtained from various experimental studies cited in U.S. EPA (1998). $FCMs$ were obtained from U.S. EPA (1995g)   | 6.80E+03 |



**TABLE A-1b-20**

**CHEMICAL-SPECIFIC INPUTS FOR  
DIETHYLSTILBESTROL (56-53-1)**

| <b>Parameter</b>                                   | <b>Reference and Explanation</b>                               | <b>Value</b> |
|--|--|--------------|
| <b>Health Benchmarks</b>                           |  |              |
| <i>RfD</i> (mg/kg/day)                             | U.S.EPA (1996d)  | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>          | U.S. EPA (1997b)   | 4.70E+03     |
| <i>RfC</i> (mg/m <sup>3</sup> )                    | U.S. EPA (1997b)   | ND           |
| <i>Inhalation CSF</i><br>(mg/kg/day) <sup>-1</sup> | Value based on Oral CSF assuming route-to-route extrapolation. | ND           |
| <i>MCL</i>   | National Primary Drinking Water Regulations.                   | ND           |
| Aquatic TRV (µg/l)                                 | Ambient Water Quality Criteria, U.S. EPA, Office of Water.     | ND           |

Note: NA = Not applicable, ND = No data available

**TABLE A-1b-21**

**CHEMICAL-SPECIFIC INPUTS FOR  
DIMETHOATE (60-51-5)**

| <b>Parameter</b>                           | <b>Reference and Explanation</b>  | <b>Value</b> |
|--|---|--------------|
| <b>Chemical/Physical Properties</b>        |   |              |
| <i>MW</i> (g/mole)                         | Budavari, O'Neil, Smith, and Heckelman (1989)   | 229.28       |
| <i>T<sub>m</sub></i> (K)                   | --  |              |
| <i>V<sub>p</sub></i> (atm)                 | U.S. EPA (1995g)  | 6.70E-09     |
| <i>S</i> (mg/L)                            | U.S. EPA (1995g)  | 2.50E+04     |
| <i>H</i> (atm·m <sup>3</sup> /mol)         | U.S. EPA (1995g)  | 6.15E-11     |
| <i>D<sub>a</sub></i> (cm <sup>2</sup> /s)  | <i>D<sub>a</sub></i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 2.58E-02     |
| <i>D<sub>w</sub></i> (cm <sup>2</sup> /s)  | <i>D<sub>w</sub></i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 5.82E-06     |
| <i>K<sub>ow</sub></i> (unitless)           | Geometric mean value cited in U.S. EPA (1994c).   | 4.90E+00     |
| <i>K<sub>oc</sub></i> (mL/g)               | <i>K<sub>oc</sub></i> value was calculated by using the correlation equation with <i>K<sub>ow</sub></i> for phthalates and PAHs, cited in U.S. EPA (1994c). <i>K<sub>oc</sub></i> value was calculated by using the recommended <i>K<sub>ow</sub></i> value that is provided in this table.   | 4.80E+00     |
| <i>Kd<sub>s</sub></i> (cm <sup>3</sup> /g) | <i>Kd<sub>s</sub></i> value was calculated by using the correlation equation with <i>K<sub>oc</sub></i> that is cited in U.S. EPA (1993) for an assumed fraction organic carbon of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate <i>Kd<sub>s</sub></i> , because the value varies, depending on the fraction of organic carbon in soil. Recommended <i>Kd<sub>s</sub></i> value was calculated by using the <i>K<sub>oc</sub></i> value that is provided in this table.   | 4.80E-02     |
| <i>Kd<sub>sw</sub></i> (L/Kg)              | <i>Kd<sub>sw</sub></i> value was calculated by using the correlation equation with <i>K<sub>oc</sub></i> that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate <i>Kd<sub>sw</sub></i> , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended <i>Kd<sub>sw</sub></i> value was calculated by using the <i>K<sub>oc</sub></i> value that is provided in this table. | 3.60E-01     |
| <b>Dermal Exposure Factors</b>             |   |              |
| <i>K<sub>p</sub></i> (cm/hr)               | <i>K<sub>p</sub></i> value was obtained from U.S. EPA (1992b).  | 2.40E-04     |
| <i>t</i>                                   | <i>t</i> value was obtained from U.S. EPA (1992b).  | 2.20E+00     |
| <i>t*</i>                                  | <i>t*</i> value was obtained from U.S. EPA (1992b).   | 5.30E+00     |
| <i>B</i>                                   | <i>B</i> value was obtained from U.S. EPA (1992b).  | 4.90E-04     |
| <b>Biotransfer Factors for Animals</b>     |   |              |
| <i>BCF<sub>fish</sub></i> (L/kg FW tissue) | <i>BCFs</i> were used for compounds with a log <i>K<sub>ow</sub></i> value below 4.0, as cited in U.S. EPA (1995g). <i>BCF<sub>fish</sub></i> value calculated using the correlation equation with <i>K<sub>ow</sub></i> obtained from Veith, Macek, Petrocelli, and Carroll (1980)   | 8.40E-01     |
| <i>BAF<sub>fish</sub></i> (L/kg FW)        | --  | NA           |

**TABLE A-1b-21**

**CHEMICAL-SPECIFIC INPUTS FOR  
DIMETHOATE (60-51-5)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S.EPA (1997c)   | 2.00E-04     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 7.00E-04     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA= Not applicable, ND= No data available

TABLE A-1b-22

**CHEMICAL-SPECIFIC INPUTS FOR  
7,12-DIMETHYLBENZ[A]ANTHRACENE (57-97-6)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 256.35   |
| $T_m$ (K)                              | --   |          |
| $V_p$ (atm)                            | Geometric mean value cited in U.S. EPA (1994c).  | 3.80E-12 |
| $S$ (mg/L)                             | Geometric mean value cited in U.S. EPA (1994c).  | 5.00E-02 |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value was calculated by using the theoretical equation from Lyman, Reehl, and Rosenblatt (1982), which defines the constant. Recommended value was calculated by using the $MW$ , $S$ , and $V_p$ values that are provided in this table.  | 3.11E-08 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 4.61E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 4.98E-06 |
| $K_{ow}$ (unitless)                    | Geometric mean value cited in U.S. EPA (1994c).  | 4.20E+06 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value was calculated by using the correlation equation with $K_{ow}$ for all nonionizing organics except phthalates, PAHs, dioxins, and furans as cited in U.S. EPA (1994c). $K_{oc}$ value was calculated by using the recommended $K_{ow}$ value that is provided in this table.  | 3.20E+06 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed fraction organic carbon of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 3.20E+04 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 2.40E+05 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 2.60E+00 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 3.20E+00 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 1.50E+01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 4.20E+02 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | --   | NA       |
| $BAF_{fish}$ (L/kg FW)                 | $BAFs$ were used for compounds with a log $K_{ow}$ value above 4.0, as cited in U.S. EPA (1995g). $BAF$ values were predicted values calculated by multiplying a food chain multiplier ( $FCM$ ) with an estimated $BCF$ . $BCFs$ were estimated using the correlation equation obtained from Veith, Macek, Petrocelli, and Caroll (1980). $FCMs$ were obtained from U.S. EPA (1995g)-   | 1.00E+03 |

**TABLE A-1b-22**

**CHEMICAL-SPECIFIC INPUTS FOR  
7,12-DIMETHYLBENZ[A]ANTHRACENE (57-97-6)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>                                      | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | --  | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | U.S. EPA (1997c)  | 2.50E+01     |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --  | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | Value based on <i>Oral CSF</i> assuming route-to-route extrapolation. | 2.50E+01     |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.                          | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA= Not applicable, ND= No data available

TABLE A-1b-23

**CHEMICAL-SPECIFIC INPUTS FOR  
3-3'-DIMETHYLBENZIDINE (119-93-7)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neill, Smith, and Heckelman (1989)   | 212.28   |
| $T_m$ (K)                              | Budavari, O'Neill, Smith, and Heckelman (1989)   |          |
| $Vp$ (atm)                             | Geometric mean value cited in U.S. EPA (1994c).  | 4.90E-10 |
| $S$ (mg/L)                             | Geometric mean value cited in U.S. EPA (1994c).  | 1.20E+03 |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value was calculated by using the theoretical equation from Lyman, Reehl, and Rosenblatt (1982), which defines the constant. Recommended value was calculated by using the $MW$ , $S$ , and $Vp$ values that are provided in this table.   | 6.29E-11 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 2.83E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 6.17E-06 |
| $K_{ow}$ (unitless)                    | Geometric mean value cited in U.S. EPA (1994c).  | 4.80E+02 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value was calculated by using the correlation equation with $K_{ow}$ for all nonionizing organics except phthalates, PAHs, dioxins, and furans as cited in U.S. EPA (1994c). $K_{oc}$ value was calculated by using the recommended $K_{ow}$ value that is provided in this table.  | 4.30E+02 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed fraction organic carbon of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 4.30E+00 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 3.22E+01 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 7.70E-03 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 1.70E+00 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 4.10E+00 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 4.80E-02 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 1.80E+01 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-23**

**CHEMICAL-SPECIFIC INPUTS FOR  
3-3'-DIMETHYLBENZIDINE (119-93-7)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>                                      | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | --  | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | U.S. EPA (1997c)  | 9.20E+00     |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --  | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | Value based on <i>Oral CSF</i> assuming route-to-route extrapolation. | 9.20E+00     |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.                          | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA= Not applicable, ND= No data available

**TABLE A-1b-24**

**CHEMICAL-SPECIFIC INPUTS FOR  
2,4-DINITRO-6-METHYLPHENOL (534-52-1)**

| <b>Parameter</b>                           | <b>Reference and Explanation</b>   | <b>Value</b> |
|--|--|--------------|
| <b>Chemical/Physical Properties</b>        |  |              |
| <i>MW</i> (g/mole)                         | Budavari, O'Neil, Smith, and Heckelman (1989)                                    | 198.13       |
| <i>T<sub>m</sub></i> (K)                   | Budavari, O'Neil, Smith, and Heckelman (1989)                                    | 360.65       |
| <i>V<sub>p</sub></i> (atm)                 | --   | ND           |
| <i>S</i> (mg/L)                            | --   | ND           |
| <i>H</i> (atm·m <sup>3</sup> /mol)         | <i>H</i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d).             | 4.26E-07     |
| <i>D<sub>a</sub></i> (cm <sup>2</sup> /s)  | <i>D<sub>a</sub></i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 2.93E-02     |
| <i>D<sub>w</sub></i> (cm <sup>2</sup> /s)  | <i>D<sub>w</sub></i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 6.91E-06     |
| <i>K<sub>ow</sub></i> (unitless)           | --   | ND           |
| <i>K<sub>oc</sub></i> (mL/g)               | --   | ND           |
| <i>Kd<sub>s</sub></i> (cm <sup>3</sup> /g) | --   | ND           |
| <i>Kd<sub>sw</sub></i> (L/Kg)              | --   | ND           |
| <b>Dermal Exposure Factors</b>             |  |              |
| <i>K<sub>p</sub></i> (cm/hr)               | <i>K<sub>p</sub></i> value was obtained from U.S. EPA (1992b).                   | 3.78E-03     |
| <i>t</i>                                   | <i>t</i> value was obtained from U.S. EPA (1992b).                               | 1.41E+00     |
| <i>t*</i>                                  | <i>t*</i> value was obtained from U.S. EPA (1992b).                              | 3.39E+00     |
| <i>B</i>                                   | <i>B</i> value was obtained from U.S. EPA (1992b).                               | 1.32E-02     |
| <b>Biotransfer Factors for Animals</b>     |  |              |
| <i>BCF<sub>fish</sub></i> (L/kg FW tissue) | --   | ND           |
| <i>BAF<sub>fish</sub></i> (L/kg FW)        | --   | ND           |



**TABLE A-1b-24**

**CHEMICAL-SPECIFIC INPUTS FOR  
2,4-DINITRO-6-METHYLPHENOL (534-52-1)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>             | <b>Value</b> |
|---|--|--------------|
| <b>Health Benchmarks</b>                        |  |              |
| <i>RfD</i> (mg/kg/day)                          | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --   | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                              | --   | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-25

**CHEMICAL-SPECIFIC INPUTS FOR  
DIPHENYLAMINE (122-39-4)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neill, Smith, and Heckelman (1989)   | 169.23   |
| $T_m$ (K)                              | Budavari, O'Neill, Smith, and Heckelman (1989)   |          |
| $V_p$ (atm)                            | Geometric mean value cited in U.S. EPA (1994c).  | 5.60E-06 |
| $S$ (mg/L)                             | Geometric mean value cited in U.S. EPA (1994c).  | 3.00E+02 |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value was calculated by using the theoretical equation from Lyman, Reehl, and Rosenblatt (1982), which defines the constant. Recommended value was calculated by using the $MW$ , $S$ , and $V_p$ values that are provided in this table.  | 4.96E-07 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 6.80E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 6.31E-06 |
| $K_{ow}$ (unitless)                    | Geometric mean value cited in U.S. EPA (1994c).  | 3.00E+03 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value was calculated by using the correlation equation with $K_{ow}$ for all nonionizing organics except phthalates, PAHs, dioxins, and furans as cited in U.S. EPA (1994c). $K_{oc}$ value was calculated by using the recommended $K_{ow}$ value that is provided in this table.  | 2.60E+03 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed fraction organic carbon of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 2.60E+01 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 1.95E+00 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 5.20E-02 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 9.40E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 5.00E+00 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 3.00E-01 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCF_s$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)   | 8.30E+01 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-25**

**CHEMICAL-SPECIFIC INPUTS FOR  
DIPHENYLAMINE (122-39-4)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997c)  | 2.50E-02     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 8.82E-02     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-26

**CHEMICAL-SPECIFIC INPUTS FOR  
2-ETHOXYETHANOL (110-80-5)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neill, Smith, and Heckelman (1989)   | 90.12    |
| $T_m$ (K)                              | Budavari, O'Neill, Smith, and Heckelman (1989)   |          |
| $V_p$ (atm)                            | Geometric mean value cited in U.S. EPA (1994c).  | 7.00E-03 |
| $S$ (mg/L)                             | Geometric mean value cited in U.S. EPA (1994c).  | 1.20E+01 |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value was calculated by using the theoretical equation from Lyman, Reehl, and Rosenblatt (1982), which defines the constant. Recommended value was calculated by using the $MW$ , $S$ , and $V_p$ values that are provided in this table.  | 1.23E-07 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 9.47E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 9.57E-06 |
| $K_{ow}$ (unitless)                    | Geometric mean value cited in U.S. EPA (1994c).  | 7.90E-01 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value was calculated by using the correlation equation with $K_{ow}$ for all nonionizing organics except phthalates, PAHs, dioxins, and furans as cited in U.S. EPA (1994c). $K_{oc}$ value was calculated by using the recommended $K_{ow}$ value that is provided in this table.  | 8.00E-01 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed fraction organic carbon of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 8.00E-03 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 6.00E-02 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 4.60E-04 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 3.10E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 7.40E-01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 7.90E-05 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 2.20E-01 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-26**

**CHEMICAL-SPECIFIC INPUTS FOR  
2-ETHOXYETHANOL (110-80-5)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | Calculated from <i>RfC</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 5.70E-02     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | U.S. EPA (1997c)  | 2.00E-01     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-27

**CHEMICAL-SPECIFIC INPUTS FOR  
ETHYL ACETATE (141-78-6)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neill, Smith, and Heckelman (1989)   | 88.1     |
| $T_m$ (K)                              | Budavari, O'Neill, Smith, and Heckelman (1989)   |          |
| $V_p$ (atm)                            | Geometric mean value cited in U.S. EPA (1994c).  | 1.20E-01 |
| $S$ (mg/L)                             | Geometric mean value cited in U.S. EPA (1994c).  | 6.40E+04 |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value was calculated by using the theoretical equation from Lyman, Reehl, and Rosenblatt (1982), which defines the constant. Recommended value was calculated by using the $MW$ , $S$ , and $V_p$ values that are provided in this table.  | 1.38E-04 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 7.32E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 9.66E-06 |
| $K_{ow}$ (unitless)                    | Geometric mean value cited in U.S. EPA (1994c).  | 4.90E+00 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value was calculated by using the correlation equation with $K_{ow}$ for all nonionizing organics except phthalates, PAHs, dioxins, and furans as cited in U.S. EPA (1994c). $K_{oc}$ value was calculated by using the recommended $K_{ow}$ value that is provided in this table.  | 4.80E+00 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed fraction organic carbon of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 4.80E-02 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 3.60E-01 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 1.70E-03 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 3.00E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 7.20E-01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 4.90E-04 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 9.90E-01 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-27**

**CHEMICAL-SPECIFIC INPUTS FOR  
ETHYL ACETATE (141-78-6)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>   | <b>Value</b> |
|---|--|--------------|
| <b>Health Benchmarks</b>                        |  |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997c)   | 9.00E-01     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. 1997c) | 3.15E+00     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.   | ND           |
| Aquatic TRV (µg/l)                              | Ambient Water Quality Criteria, U.S. EPA, Office of Water.   | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-28

**CHEMICAL-SPECIFIC INPUTS FOR  
ETHYL ETHER (60-29-7)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neill, Smith, and Heckelman (1989)   | 74.12    |
| $T_m$ (K)                              | Budavari, O'Neill, Smith, and Heckelman (1989)   |          |
| $V_p$ (atm)                            | Geometric mean value cited in U.S. EPA (1994c).  | 7.10E-01 |
| $S$ (mg/L)                             | Geometric mean value cited in U.S. EPA (1994c).  | 6.10E+04 |
| $H$ (atm·m <sup>3</sup> /mol)          | U.S. EPA (1995g)   | 8.70E-04 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 7.40E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 9.30E-06 |
| $K_{ow}$ (unitless)                    | Geometric mean value cited in U.S. EPA (1994c).  | 6.80E+00 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value was calculated by using the correlation equation with $K_{ow}$ for all nonionizing organics except phthalates, PAHs, dioxins, and furans as cited in U.S. EPA (1994c). $K_{oc}$ value was calculated by using the recommended $K_{ow}$ value that is provided in this table.  | 6.50E+00 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed fraction organic carbon of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 6.50E-02 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 4.88E-01 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 2.60E-03 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 2.50E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 5.90E-01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 6.80E-04 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 1.30E+00 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |



**TABLE A-1b-28**

**CHEMICAL-SPECIFIC INPUTS FOR  
ETHYL ETHER (60-29-7)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S.EPA (1996b)   | 2.00E-01     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 7.00E-01     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA= Not applicable, ND= No data available

TABLE A-1b-29

**CHEMICAL-SPECIFIC INPUTS FOR  
ETHYLENE THIOUREA (ETU) (96-45-7)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neill, Smith, and Heckelman (1989)   | 102.17   |
| $T_m$ (°K)                             | Budavari, O'Neill, Smith, and Heckelman (1989)   | 476.65   |
| $V_p$                                  | Geometric mean value cited in U.S. EPA (1994c).  | 1.10E-04 |
| $S$ (mg/L)                             | Geometric mean value cited in U.S. EPA (1994c).  | 1.20E+04 |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value was calculated by using the theoretical equation from Lyman, Reehl, and Rosenblatt (1982), which defines the constant. Recommended value was calculated by using the $MW$ , $S$ , and $V_p$ values that are provided in this table.  | 3.08E-10 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 7.15E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 1.02E-05 |
| $K_{ow}$ (unitless)                    | Geometric mean value cited in U.S. EPA (1994c).  | 2.20E-01 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value was calculated by using the correlation equation with $K_{ow}$ for all nonionizing organics except phthalates, PAHs, dioxins, and furans as cited in U.S. EPA (1994c). $K_{oc}$ value was calculated by using the recommended $K_{ow}$ value that is provided in this table.  | 2.20E-01 |
| $Kd_s$ (mL/g)                          | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 2.20E-03 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 1.65E-02 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 1.50E-04 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 3.70E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 8.80E-01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 2.20E-05 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 7.90E-02 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

| Parameter                                       | Reference and Explanation  | Value    |
|---|--|----------|
| <b>Health Benchmarks</b>                        |  |          |
| <i>RfD</i> (mg/kg/day)                          | --   | ND       |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | Value based on Inhalation CSF assuming route-to-route extrapolation. | 1.10E-01 |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --   | ND       |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | U.S. EPA (1997c)   | 1.10E-01 |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.                         | ND       |
| Aquatic TRV (µg/l)                              | --   | ND       |

Note: ND= Not applicable, ND= No data available

**TABLE A-1b-30**

**CHEMICAL-SPECIFIC INPUTS FOR  
FURAN (110-00-9)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neill, Smith, and Heckelman (1989)   | 68.08    |
| $T_m$ (K)                              | Budavari, O'Neill, Smith, and Heckelman (1989)   |          |
| $Vp$ (atm)                             | Geometric mean value cited in U.S. EPA (1994c).  | 7.90E-01 |
| $S$ (mg/L)                             | Geometric mean value cited in U.S. EPA (1994c).  | 1.00E+04 |
| $H$ (atm·m <sup>3</sup> /mol)          | U.S. EPA (1995g)   | 5.40E-03 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 1.04E-01 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 1.22E-05 |
| $K_{ow}$ (unitless)                    | Geometric mean value cited in U.S. EPA (1994c).  | 2.20E+01 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value was calculated by using the correlation equation with $K_{ow}$ for all nonionizing organics except phthalates, PAHs, dioxins, and furans as cited in U.S. EPA (1994c). $K_{oc}$ value was calculated by using the recommended $K_{ow}$ value that is provided in this table.  | 2.10E+01 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 2.10E-01 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 1.57E+00 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 6.50E-03 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 2.30E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 5.50E-01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 2.20E-03 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (unitless, FW tissue)     | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 3.00E+00 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-30**

**CHEMICAL-SPECIFIC INPUTS FOR  
FURAN (110-00-9)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997b)  | 1.00-03      |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 3.50E-03     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-31

**CHEMICAL-SPECIFIC INPUTS FOR  
GAMMA-HEXACHLOROCYCLOHEXANE (LINDANE) (58-89-9)**

| Parameter                              | Reference and Explanation  | Value     |
|--|--|-----------|
| <b>Chemical/Physical Properties</b>    |  |           |
| $MW$ (g/mole)                          | Montgomery and Welkom (1991)   | 290.85    |
| $T_m$ (K)                              | Montgomery and Welkom (1991)   | 385.65    |
| $Vp$ (mmHG)                            | Geometric mean value cited in U.S. EPA (1994c).  | 9.40 E-06 |
| $S$ (in water)                         | U.S. EPA (1995g)   | 4.20E+00  |
| $H$ (atm·m <sup>3</sup> /mol)          | U.S. EPA (1995g).  | 3.40E-06  |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was calculated using the equation cited in U.S. EPA (1996a).   | 1.42E-02  |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was calculated using the equation cited in U.S. EPA (1996a).   | 7.34E-06  |
| $K_{ow}$ (unitless)                    | Geometric mean value cited in U.S. EPA (1994c)   | 5.40E+03  |
| $K_{oc}$ (mL/g)                        | Geometric mean of measured values obtained from U.S. EPA (1996b).  | 4.60E+03  |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 4.60E+01  |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 3.45E+02  |
| <b>Dermal Exposure Factors</b>         |  |           |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 1.40E-02  |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 5.20E+00  |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 3.50E+01  |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 5.40E-01  |
| <b>Biotransfer Factors for Animals</b> |  |           |
| $BCF_{fish}$ (L/kg FW tissue)          | --   | NA        |
| $BAF_{fish}$ (L/kg FW)                 | $BAFs$ were used for compounds with a log $K_{ow}$ value above 4.0, as cited in U.S. EPA (1995g). $BAF$ values were predicted values calculated by multiplying a food chain multiplier ( $FCM$ ) with an estimated $BCF$ . $BCFs$ were estimated using the correlation equation obtained from Veith, Macek, Petrocelli, and Carroll (1980). $FCMs$ were obtained from U.S. EPA (1995g)   | 1.60E+03  |

**TABLE A-1b-31**

**CHEMICAL-SPECIFIC INPUTS FOR  
GAMMA-HEXACHLOROCYCLOHEXANE (LINDANE) (58-89-9)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S.EPA (1997b)   | 3.00E-04     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | U.S.EPA (1997b)   | 1.30E+00     |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 1.05E-03     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | Value based on Oral CSF assuming route-to-route extrapolation.  | 1.30E+00     |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | 2E-04        |
| Aquatic TRV (µg/l)                              | U.S. EPA (1995f)  | 8E-02        |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-32

**CHEMICAL-SPECIFIC INPUTS FOR  
HEXACHLOROPROPENE (1888-71-7)**

| Parameter                              | Reference and Explanation   | Value    |
|--|---|----------|
| <b>Chemical/Physical Properties</b>    |   |          |
| $MW$ (g/mole)                          | CRC Handbook (1995)   | 248.75   |
| $T_m$ (K)                              | CRC Handbook (1995)   | 200.25   |
| $Vp$ (atm)                             | --  | ND       |
| $S$ (mg/L)                             | --  | ND       |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value was obtained from WATER8 model database (U.S. EPA 1995d).   | 4.70E-03 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from WATER8 model database (U.S. EPA 1995d). | 8.00E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from WATER8 model database (U.S. EPA 1995d). | 8.00E-06 |
| $K_{ow}$ (unitless)                    | --  | ND       |
| $K_{oc}$ (mL/g)                        | --  | ND       |
| $Kd_s$ (cm <sup>3</sup> /g)            | --  | ND       |
| $Kd_{sw}$ (L/Kg)                       | --  | ND       |
| <b>Dermal Exposure Factors</b>         |   |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).                        | ND       |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).                         | 2.88E+00 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).                       | ND       |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).                         | ND       |
| <b>Biotransfer Factors for Animals</b> |   |          |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --  | ND       |
| $BAF_{fish}$ (L/kg FW)                 | --  | ND       |



**TABLE A-1b-32**

**CHEMICAL-SPECIFIC INPUTS FOR  
HEXACHLOROPROPENE (1888-71-7)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>             | <b>Value</b> |
|---|--|--------------|
| <b>Health Benchmarks</b>                        |  |              |
| <i>RfD</i> (mg/kg/day)                          | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --   | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                              | --   | ND           |

Note: NA= Not applicable, ND= No data available

**TABLE A-1b-33**

**CHEMICAL-SPECIFIC INPUTS FOR  
IRON (7439-89-6)**

| <b>Parameter</b>                               | <b>Reference and Explanation</b>                               | <b>Value</b> |
|--|--|--------------|
| <b>Chemical/Physical Properties</b>            |  |              |
| <i>MW</i> (g/mole)                             | Budavari, O'Neill, Smith, and Heckelman (1989)                 | 55.84        |
| <i>T<sub>m</sub></i> (K)                       | Budavari, O'Neill, Smith, and Heckelman (1989)                 | 1808.15      |
| <i>V<sub>p</sub></i> (atm)                     | --   | NA           |
| <i>S</i> (mg/L)                                | --   | NA           |
| <i>H</i> (atm·m <sup>3</sup> /mol)             | --   | NA           |
| <i>D<sub>a</sub></i> (cm <sup>2</sup> /s)      | --   | NA           |
| <i>D<sub>w</sub></i> (cm <sup>2</sup> /s)      | --   | NA           |
| <i>K<sub>ow</sub></i> (unitless)               | --   | NA           |
| <i>K<sub>oc</sub></i> (mL/g)                   | --   | NA           |
| <i>Kd<sub>s</sub></i> (cm <sup>3</sup> /g)     | --   | NA           |
| <i>Kd<sub>sw</sub></i> (L/Kg)                  | --   | NA           |
| <b>Dermal Exposure Factors</b>                 |  |              |
| <i>K<sub>p</sub></i> (cm/hr)                   | <i>K<sub>p</sub></i> value was obtained from U.S. EPA (1992b). | ND           |
| <i>t</i>                                       | <i>t</i> value was obtained from U.S. EPA (1992b).             | 1.00E-03     |
| <i>t*</i>                                      | <i>t*</i> value was obtained from U.S. EPA (1992b).            | ND           |
| <i>B</i>                                       | <i>B</i> value was obtained from U.S. EPA (1992b).             | ND           |
| <b>Biotransfer Factors for Animals</b>         |  |              |
| <i>BCF<sub>fish</sub></i><br>(L/kg, FW tissue) | --   | ND           |
| <i>BAF<sub>fish</sub></i> (L/kg FW)            | --   | ND           |

**TABLE A-1b-33**

**CHEMICAL-SPECIFIC INPUTS FOR  
IRON (7439-89-6)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S.EPA (1997b)   | 3.00E-01     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 1.05E+00     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --.   | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | Ambient Water Quality Criteria, U.S. EPA, Office of Water.  | 1.0E+03      |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-34

CHEMICAL-SPECIFIC INPUTS FOR  
ISOBUTYL ALCOHOL (78-83-1)

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neill, Smith, and Heckelman (1989)   | 74.14    |
| $T_m$ (K)                              | Budavari, O'Neill, Smith, and Heckelman (1989)   |          |
| $Vp$ (atm)                             | Geometric mean value cited in U.S. EPA (1994c).  | 1.40E-02 |
| $S$ (mg/L)                             | Geometric mean value cited in U.S. EPA (1994c).  | 7.60E+04 |
| $H$ (atm·m <sup>3</sup> /mol)          | U.S. EPA (1995g)   | 1.30E-05 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.60E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 9.30E-06 |
| $K_{ow}$ (unitless)                    | Geometric mean value cited in U.S. EPA (1994c).  | 5.60E+00 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value was calculated by using the correlation equation with $K_{ow}$ for all nonionizing organics except phthalates, PAHs, dioxins, and furans as cited in U.S. EPA (1994c). $K_{oc}$ value was calculated by using the recommended $K_{ow}$ value that is provided in this table.  | 5.50E+00 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed fraction organic carbon of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 5.50E-02 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 4.13E-01 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 2.30E-03 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 2.50E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 5.90E-01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 5.60E-04 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 1.00E+00 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-34**

**CHEMICAL-SPECIFIC INPUTS FOR  
ISOBUTYL ALCOHOL (78-83-1)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997b)  | 3.00E-01     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 1.05E+00     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-35

**CHEMICAL-SPECIFIC INPUTS FOR  
KEPONE (143-50-0)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neill, Smith, and Heckelman (1989)   | 490.68   |
| $T_m$ (K)                              | Budavari, O'Neill, Smith, and Heckelman (1989)   |          |
| $Vp$ (atm)                             | Geometric mean value cited in U.S. EPA (1994c).  | 3.90E-10 |
| $S$ (mg/L)                             | Geometric mean value cited in U.S. EPA (1994c).  | 7.60E+00 |
| $H$ (atm·m <sup>3</sup> /mol)          | U.S. EPA (1995g)   | 2.55E-08 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-06 |
| $K_{ow}$ (unitless)                    | Geometric mean value cited in U.S. EPA (1994c).  | 2.00E+05 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value was calculated by using the correlation equation with $K_{ow}$ for all nonionizing organics except phthalates, PAHs, dioxins, and furans as cited in U.S. EPA (1994c). $K_{oc}$ value was calculated by using the recommended $K_{ow}$ value that is provided in this table.  | 1.60E+05 |
| $Kd_s$ (mL/g)                          | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed fraction organic carbon of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 1.60E+03 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 1.20E+04 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 1.10E-02 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 8.60E+01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 4.10E+02 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 2.00E+01 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | --   | NA       |
| $BAF_{fish}$ (L/kg FW)                 | $BAFs$ were used for compounds with a log $K_{ow}$ value above 4.0, as cited in U.S. EPA (1995g). $BAF$ values were predicted values calculated by multiplying a food chain multiplier ( $FCM$ ) with an estimated $BCF$ . $BCFs$ were estimated using the correlation equation obtained from Veith, Macek, Petrocelli, and Caroll (1980). $FCMs$ were obtained from U.S. EPA (1995g)  | 1.80E+03 |

**TABLE A-1b-35**

**CHEMICAL-SPECIFIC INPUTS FOR  
KEPONE (143-50-0)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>                                      | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | --  | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | U.S. EPA (1997b)  | 1.80E+01     |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --  | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | Value based on <i>Oral CSF</i> assuming route-to-route extrapolation. | 1.80E+01     |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.                          | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

**TABLE A-1b-36**

**CHEMICAL-SPECIFIC INPUTS FOR  
MAGNESIUM (7439-95-4)**

| <b>Parameter</b>                       | <b>Reference and Explanation</b>               | <b>Value</b> |
|--|--|--------------|
| <b>Chemical/Physical Properties</b>    |  |              |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 24.30        |
| $T_m$ (K)                              | Budavari, O'Neil, Smith, and Heckelman (1989)  | 924.15       |
| $V_p$ (atm)                            | --   | NA           |
| $S$ (mg/L)                             | --   | NA           |
| $H$ (atm·m <sup>3</sup> /mol)          | --   | NA           |
| $D_a$ (cm <sup>2</sup> /s)             | --   | NA           |
| $D_w$ (cm <sup>2</sup> /s)             | --   | NA           |
| $K_{ow}$ (unitless)                    | --   | NA           |
| $K_{oc}$ (mL/g)                        | --   | NA           |
| $Kd_s$ (cm <sup>3</sup> /g)            | --   | NA           |
| $Kd_{sw}$ (L/Kg)                       | --   | NA           |
| <b>Dermal Exposure Factors</b>         |  |              |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b). | 1.00E-03     |
| $t$                                    | --   | ND           |
| $t^*$                                  | --   | ND           |
| $B$                                    | --   | ND           |
| <b>Biotransfer Factors for Animals</b> |  |              |
| $BCF_{fish}$<br>(L/kg, FW tissue)      | --   | ND           |
| $BAF_{fish}$ (L/kg FW)                 | --   | ND           |



**TABLE A-1b-36**

**CHEMICAL-SPECIFIC INPUTS FOR  
MAGNESIUM (7439-95-4)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>             | <b>Value</b> |
|---|--|--------------|
| <b>Health Benchmarks</b>                        |  |              |
| <i>RfD</i> (mg/kg/day)                          | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --   | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                              | --   | ND           |

Note: NA = Not applicable, ND = No data available

**TABLE A-1b-37**

**CHEMICAL-SPECIFIC INPUTS FOR  
MANGANESE (7439-96-5)**

| <b>Parameter</b>                       | <b>Reference and Explanation</b>                                 | <b>Value</b> |
|--|--|--------------|
| <b>Chemical/Physical Properties</b>    |  |              |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)                    | 54.94        |
| $T_m$ (K)                              | Budavari, O'Neil, Smith, and Heckelman (1989)                    | 1517.15      |
| $V_p$ (atm)                            | --   | ND           |
| $S$ (mg/L)                             | --   | ND           |
| $H$ (atm·m <sup>3</sup> /mol)          | --   | ND           |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d) | 8.00E-02     |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d) | 8.00E-06     |
| $K_{ow}$ (unitless)                    | --   | ND           |
| $K_{oc}$ (mL/g)                        | --   | ND           |
| $Kd_s$ (cm <sup>3</sup> /g)            | --   | ND           |
| $Kd_{sw}$ (L/Kg)                       | --   | ND           |
| <b>Dermal Exposure Factors</b>         |  |              |
| $K_p$ (cm/hr)                          | $K_p$ value was obtained from U.S. EPA (1992b).                  | 1.00E-03     |
| $t$                                    | --   | ND           |
| $t^*$                                  | --   | ND           |
| $B$                                    | --   | ND           |
| <b>Biotransfer Factors for Animals</b> |  |              |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --   | ND           |
| $BAF_{fish}$ (L/kg FW)                 | --   | ND           |

**TABLE A-1b-37**

**CHEMICAL-SPECIFIC INPUTS FOR  
MANGANESE (7439-96-5)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>             | <b>Value</b> |
|---|--|--------------|
| <b>Health Benchmarks</b>                        |  |              |
| <i>RfD</i> (mg/kg/day)                          | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --   | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                              | U.S. EPA (1996c)                             | 8.00E+01     |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-38

CHEMICAL-SPECIFIC INPUTS FOR  
3-METHYLCHOLANTHRENE (56-49-5)

| Parameter                              | Reference and Explanation   | Value    |
|--|---|----------|
| <b>Chemical/Physical Properties</b>    |   |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)   | 268.34   |
| $T_m$ (K)                              | Budavari, O'Neil, Smith, and Heckelman (1989)   | 452.65   |
| $V_p$ (atm)                            | --  | ND       |
| $S$ (mg/L)                             | --  | ND       |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).   | 9.4E-07  |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).   | 2.09E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).   | 5.36E-06 |
| $K_{ow}$ (unitless)                    | --  | ND       |
| $K_{oc}$ (mL/g)                        | --  | ND       |
| $Kd_s$ (cm <sup>3</sup> /g)            | --  | ND       |
| $Kd_{sw}$ (L/Kg)                       | --  | ND       |
| <b>Dermal Exposure Factors</b>         |   |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).  | 1.60E+00 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).   | 3.80E+00 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).   | 1.80E+01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).   | 2.60E+02 |
| <b>Biotransfer Factors for Animals</b> |   |          |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --  | NA       |
| $BAF_{fish}$ (L/kg FW)                 | $BAFs$ were used for compounds with a log $K_{ow}$ value above 4.0, as cited in U.S. EPA (1995g). $BAF$ values were predicted values calculated by multiplying a food chain multiplier ( $FCM$ ) with an estimated $BCF$ . $BCFs$ were estimated using the correlation equation obtained from Veith, Macek, Petrocelli, and Caroll (1980). $FCMs$ were obtained from U.S. EPA (1995g) | 1.96E+06 |

**TABLE A-1b-38**

**CHEMICAL-SPECIFIC INPUTS FOR  
3-METHYLCHOLANTHRENE (56-49-5)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>                                      | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | --  | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | U.S. EPA (1997c)  | 2.60E+01     |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --  | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | Value based on <i>Oral CSF</i> assuming route-to-route extrapolation. | 2.60E+01     |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.                          | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

**TABLE A-1b-39**

**CHEMICAL-SPECIFIC INPUTS FOR  
METHYL METHACRYLATE (80-62-6)**

| <b>Parameter</b>                           | <b>Reference and Explanation</b>  | <b>Value</b> |
|--|---|--------------|
| <b>Chemical/Physical Properties</b>        |   |              |
| <i>MW</i> (g/mole)                         | Budavari, O'Neill, Smith, and Heckelman (1989)  | 100.13       |
| <i>T<sub>m</sub></i> (K)                   | Budavari, O'Neill, Smith, and Heckelman (1989)  |              |
| <i>V<sub>p</sub></i> (atm)                 | Geometric mean value cited in U.S. EPA (1994c).   | 5.10E-02     |
| <i>S</i> (mg/L)                            | Geometric mean value cited in U.S. EPA (1994c).   | 1.60E+04     |
| <i>H</i> (atm·m <sup>3</sup> /mol)         | U.S. EPA (1995g)  | 3.20E-04     |
| <i>D<sub>a</sub></i> (cm <sup>2</sup> /s)  | <i>D<sub>a</sub></i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 7.70E-02     |
| <i>D<sub>w</sub></i> (cm <sup>2</sup> /s)  | <i>D<sub>w</sub></i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.60E-06     |
| <i>K<sub>ow</sub></i> (unitless)           | Geometric mean value cited in U.S. EPA (1994c).   | 2.40E+01     |
| <i>K<sub>oc</sub></i> (mL/g)               | <i>K<sub>oc</sub></i> value was calculated by using the correlation equation with <i>K<sub>ow</sub></i> for all nonionizing organics except phthalates, PAHs, dioxins, and furans as cited in U.S. EPA (1994c). <i>K<sub>oc</sub></i> value was calculated by using the recommended <i>K<sub>ow</sub></i> value that is provided in this table.   | 2.30E+01     |
| <i>Kd<sub>s</sub></i> (cm <sup>3</sup> /g) | <i>Kd<sub>s</sub></i> value was calculated by using the correlation equation with <i>K<sub>oc</sub></i> that is cited in U.S. EPA (1993) for an assumed fraction organic carbon of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate <i>Kd<sub>s</sub></i> , because the value varies, depending on the fraction of organic carbon in soil. Recommended <i>Kd<sub>s</sub></i> value was calculated by using the <i>K<sub>oc</sub></i> value that is provided in this table.   | 2.30E-01     |
| <i>Kd<sub>sw</sub></i> (L/Kg)              | <i>Kd<sub>sw</sub></i> value was calculated by using the correlation equation with <i>K<sub>oc</sub></i> that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate <i>Kd<sub>sw</sub></i> , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended <i>Kd<sub>sw</sub></i> value was calculated by using the <i>K<sub>oc</sub></i> value that is provided in this table. | 1.72E+00     |
| <b>Dermal Exposure Factors</b>             |   |              |
| <i>K<sub>p</sub></i> (cm/hr)               | <i>K<sub>p</sub></i> value was obtained from U.S. EPA (1992b).  | 4.50E-03     |
| <i>t</i>                                   | <i>t</i> value was obtained from U.S. EPA (1992b).  | 3.60E-01     |
| <i>t*</i>                                  | <i>t*</i> value was obtained from U.S. EPA (1992b).   | 8.60E-01     |
| <i>B</i>                                   | <i>B</i> value was obtained from U.S. EPA (1992b).  | 2.40E-03     |
| <b>Biotransfer Factors for Animals</b>     |   |              |
| <i>BCF<sub>fish</sub></i> (L/kg FW tissue) | <i>BCFs</i> were used for compounds with a log <i>K<sub>ow</sub></i> value below 4.0, as cited in U.S. EPA (1995g). <i>BCF<sub>fish</sub></i> value calculated using the correlation equation with <i>K<sub>ow</sub></i> obtained from Veith, Macek, Petrocelli, and Carroll (1980)   | 3.2E+00      |
| <i>BAF<sub>fish</sub></i> (L/kg FW)        | --  | NA           |

**TABLE A-1b-39**

**CHEMICAL-SPECIFIC INPUTS FOR  
METHYL METHACRYLATE (80-62-6)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997b)  | 8.00E-02     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 2.80E-01     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-40

**CHEMICAL-SPECIFIC INPUTS FOR  
2-METHYLNAPHTHALENE (91-57-6)**

| Parameter                              | Reference and Explanation   | Value    |
|--|---|----------|
| <b>Chemical/Physical Properties</b>    |   |          |
| $MW$ (g/mole)                          | CRC Handbook (1995)   | 142.20   |
| $T_m$ (K)                              | CRC Handbook (1995)   | 307.55   |
| $V_p$ (atm)                            | --  | ND       |
| $S$ (mg/L)                             | --  | ND       |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).   | 5.05E-04 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 4.80E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 7.84E-06 |
| $K_{ow}$ (unitless)                    | --  | ND       |
| $K_{oc}$ (mL/g)                        | --  | ND       |
| $Kd_s$ (cm <sup>3</sup> /g)            | --  | ND       |
| $Kd_{sw}$ (L/Kg)                       | --  | ND       |
| <b>Dermal Exposure Factors</b>         |   |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).                    | 1.42E-01 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).                     | 6.44E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).                   | 4.87E+00 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).                     | 7.24E-01 |
| <b>Biotransfer Factors for Animals</b> |   |          |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --  | ND       |
| $BAF_{fish}$ (L/kg FW)                 | --  | ND       |



**TABLE A-1b-40**

**CHEMICAL-SPECIFIC INPUTS FOR  
2-METHYLNAPHTHALENE (91-57-6)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>             | <b>Value</b> |
|---|--|--------------|
| <b>Health Benchmarks</b>                        |  |              |
| <i>RfD</i> (mg/kg/day)                          | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --   | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                              | --   | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-41

**CHEMICAL-SPECIFIC INPUTS FOR  
MOLYBDENUM (7439-98-7)**

| Parameter                              | Reference and Explanation                       | Value    |
|--|---|----------|
| <b>Chemical/Physical Properties</b>    |   |          |
| $MW$ (g/mole)                          | CRC Handbook (1995)                             | 95.94    |
| $T_m$ (K)                              | --  | N/A      |
| $Vp$ (atm)                             | --  | N/A      |
| $S$ (mg/L)                             | --  | N/A      |
| $H$ (atm·m <sup>3</sup> /mol)          | --  | N/A      |
| $D_a$ (cm <sup>2</sup> /s)             | --  | N/A      |
| $D_w$ (cm <sup>2</sup> /s)             | --  | N/A      |
| $K_{ow}$ (unitless)                    | --  | N/A      |
| $K_{oc}$ (mL/g)                        | --  | N/A      |
| $Kd_s$ (cm <sup>3</sup> /g)            | U.S. EPA (1995g)                                | 2.00E+01 |
| $Kd_{sw}$ (L/Kg)                       | --  | 2.00E+01 |
| <b>Dermal Exposure Factors</b>         |   |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).  | 1.00E-03 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).   | ND       |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b). | ND       |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).   | ND       |
| <b>Biotransfer Factors for Animals</b> |   |          |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --  | ND       |
| $BAF_{fish}$ (L/kg FW)                 | --  | ND       |

**TABLE A-1b-41**

**CHEMICAL-SPECIFIC INPUTS FOR  
MOLYBDENUM (7439-98-7)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>             | <b>Value</b> |
|---|--|--------------|
| <b>Health Benchmarks</b>                        |  |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997c)                             | 5.00E-03     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --   | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                              | U.S. EPA (1996c)                             | 2.40E+00     |

Note: NA= Not applicable, ND= No data available

TABLE A-1b-42

**CHEMICAL-SPECIFIC INPUTS FOR  
1,4-NAPHTHAQUINONE (130-15-4)**

| Parameter                              | Reference and Explanation   | Value    |
|--|---|----------|
| <b>Chemical/Physical Properties</b>    |   |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)                     | 158.15   |
| $T_m$ (K)                              | Budavari, O'Neil, Smith, and Heckelman (1989)                     | 399.15   |
| $V_p$ (atm)                            | --  | NA       |
| $S$ (mg/L)                             | --  | NA       |
| $H$ (atm·m <sup>3</sup> /mol)          | --  | NA       |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 3.60E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 8.98E-06 |
| $K_{ow}$ (unitless)                    | --  | NA       |
| $K_{oc}$ (mL/g)                        | --  | NA       |
| $Kd_s$ (cm <sup>3</sup> /g)            | --  | NA       |
| $Kd_{sw}$ (L/Kg)                       | --  | NA       |
| <b>Dermal Exposure Factors</b>         |   |          |
| $K_p$ (cm/hr)                          | $K_p$ value was obtained from U.S. EPA (1992b).                   | 3.39E-03 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).                     | 8.05E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).                   | 1.93E+00 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).                     | 5.13E-03 |
| <b>Biotransfer Factors for Animals</b> |   |          |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --  | NA       |
| $BAF_{fish}$ (L/kg FW)                 | --  | NA       |

**TABLE A-1b-42**

**CHEMICAL-SPECIFIC INPUTS FOR  
1,4-NAPHTHAQUINONE (130-15-4)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>             | <b>Value</b> |
|---|--|--------------|
| <b>Health Benchmarks</b>                        |  |              |
| <i>RfD</i> (mg/kg/day)                          | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --   | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                              | --   | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-43

**CHEMICAL-SPECIFIC INPUTS FOR  
2-NAPHTHYLAMINE (91-59-8)**

| Parameter                              | Reference and Explanation   | Value    |
|--|---|----------|
| <b>Chemical/Physical Properties</b>    |   |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)   | 143.18   |
| $T_m$ (K)                              | Budavari, O'Neil, Smith, and Heckelman (1989)   | 323.15   |
| $V_p$ (atm)                            | --  | ND       |
| $S$ (mg/L)                             | --  | ND       |
| $H$ (atm·m <sup>3</sup> /mol)          | U.S. EPA (1995g)  | 6.03E-07 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).   | 4.51E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).   | 8.39E-06 |
| $K_{ow}$ (unitless)                    | --  | ND       |
| $K_{oc}$ (mL/g)                        | --  | ND       |
| $Kd_s$ (cm <sup>3</sup> /g)            | --  | ND       |
| $Kd_{sw}$ (L/Kg)                       | --  | ND       |
| <b>Dermal Exposure Factors</b>         |   |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).  | 1.10E-02 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).   | 6.50E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).   | 1.60E+00 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).   | 1.90E-02 |
| <b>Biotransfer Factors for Animals</b> |   |          |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980) | 1.70E+01 |
| $BAF_{fish}$ (L/kg FW)                 | --  | NA       |

**TABLE A-1b-43**

**CHEMICAL-SPECIFIC INPUTS FOR  
2-NAPHTHYLAMINE (91-59-8)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>             | <b>Value</b> |
|---|--|--------------|
| <b>Health Benchmarks</b>                        |  |              |
| <i>RfD</i> (mg/kg/day)                          | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --   | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                              | --   | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-44

**CHEMICAL-SPECIFIC INPUTS FOR  
5-NITRO-O-TOLUIDINE (2-METHYL-5-NITROANILINE) (99-55-8)**

| Parameter                              | Reference and Explanation   | Value    |
|--|---|----------|
| <b>Chemical/Physical Properties</b>    |   |          |
| $MW$ (g/mole)                          | CRC Handbook (1995)   | 152.15   |
| $T_m$ (K)                              | CRC Handbook (1995)   | 378.65   |
| $Vp$ (atm)                             | --  | NA       |
| $S$ (mg/L)                             | --  | NA       |
| $H$ (atm·m <sup>3</sup> /mol)          | --  | NA       |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 8.00E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 8.00E-06 |
| $K_{ow}$ (unitless)                    | --  | NA       |
| $K_{oc}$ (mL/g)                        | --  | NA       |
| $Kd_s$ (cm <sup>3</sup> /g)            | --  | NA       |
| $Kd_{sw}$ (L/Kg)                       | --  | NA       |
| <b>Dermal Exposure Factors</b>         |   |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).                    | 4.78E-03 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).                     | 7.41E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).                   | 1.78E+00 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).                     | 7.41E-03 |
| <b>Biotransfer Factors for Animals</b> |   |          |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --  | NA       |
| $BAF_{fish}$ (L/kg FW)                 | --  | NA       |



**TABLE A-1b-44**

**CHEMICAL-SPECIFIC INPUTS FOR  
5-NITRO-O-TOLUIDINE (2-METHYL-5-NITROANILINE) (99-55-8)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>             | <b>Value</b> |
|---|--|--------------|
| <b>Health Benchmarks</b>                        |  |              |
| <i>RfD</i> (mg/kg/day)                          | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --   | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                              | --   | ND           |

Note: NA= Not applicable, ND= No data available

TABLE A-1b-45

**CHEMICAL-SPECIFIC INPUTS FOR  
2-NITROPROPANE (79-46-9)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 89.09    |
| $T_m$ (K)                              | U.S. EPA (1995g)   | NA       |
| $Vp$ (atm)                             | $Vp$ value cited in U.S. EPA (1995g).  | 2.40E-02 |
| $S$ (mg/L)                             | $S$ value cited in U.S. EPA (1995g).   | 1.70E+05 |
| $H$ (atm·m <sup>3</sup> /mol)          | U.S. EPA (1995g)   | 1.23E-05 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 9.23E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 1.01E-05 |
| $K_{ow}$ (unitless)                    | $K_{ow}$ value cited in U.S. EPA (1995g)   | 7.40E+00 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value was calculated by using the correlation equation with $K_{ow}$ for phthalates and PAHs, / all nonionizing organics except phthalates, PAHs, dioxins, and furans, cited in U.S. EPA (1994c). $K_{oc}$ value was calculated by using the recommended $K_{ow}$ value that is provided in this table.   | 7.20E+00 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 7.20E-02 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 5.40E-01 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 2.30E-03 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 3.10E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 7.30E-01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 7.40E-04 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 7.20E-01 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-45**

**CHEMICAL-SPECIFIC INPUTS FOR  
2-NITROPROPANE (79-46-9)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | Calculated from <i>RfC</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 5.70E-03     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | U.S. EPA (1997b)  | 2.00E-02     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | U.S. EPA (1997c)  | 9.40E+00     |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA= Not applicable, ND= No data available

TABLE A-1b-46

CHEMICAL-SPECIFIC INPUTS FOR  
4-NITROQUINOLINE-1-OXIDE (56-57-5)

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | CRC Handbook (1995)  | 190.16   |
| $T_m$ (K)                              | CRC Handbook (1995)  | 427.15   |
| $Vp$ (atm)                             | --   | NA       |
| $S$ (mg/L)                             | --   | NA       |
| $H$ (atm·m <sup>3</sup> /mol)          | --   | NA       |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was calculated using the equation cited in U.S. EPA (1996a). | 8.00E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was calculated using the equation cited in U.S. EPA (1996a). | 8.00E-06 |
| $K_{ow}$ (unitless)                    | --   | NA       |
| $K_{oc}$ (mL/g)                        | --   | NA       |
| $Kd_s$ (cm <sup>3</sup> /g)            | --   | NA       |
| $Kd_{sw}$ (L/Kg)                       | --   | NA       |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).                           | ND       |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).                            | 4.40E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).                          | ND       |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).                            | ND       |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$<br>(L/kg, FW tissue)      | --   | NA       |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-46**

**CHEMICAL-SPECIFIC INPUTS FOR  
4-NITROQUINOLINE-1-OXIDE (56-57-5)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>             | <b>Value</b> |
|---|--|--------------|
| <b>Health Benchmarks</b>                        |  |              |
| <i>RfD</i> (mg/kg/day)                          | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --   | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                              | --   | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-47

CHEMICAL-SPECIFIC INPUTS FOR  
N-NITROSODIETHYLAMINE (55-18-5)

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 102.14   |
| $T_m$ (K)                              | --   |          |
| $Vp$ (atm)                             | U.S. EPA (1995g)   | 2.60E-03 |
| $S$ (mg/L)                             | U.S. EPA (1995g)   | 2.00E+05 |
| $H$ (atm·m <sup>3</sup> /mol)          | U.S. EPA (1995g)   | 3.63E-06 |
| $D_a$ (cm <sup>2</sup> /s)             | U.S. EPA (1995g)   | 8.00E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | U.S. EPA (1995g)   | 8.00E-06 |
| $K_{ow}$ (unitless)                    | U.S. EPA (1995g)   | 3.00E+00 |
| $K_{oc}$ (mL/g)                        | U.S. EPA (1995g)   | 3.00E+00 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 3.00E-02 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 2.10E-01 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 9.90E-04 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 3.70E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 8.80E-01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 3.00E-04 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$<br>(L/kg FW tissue)       | U.S. EPA (1995g)   | 6.30E-01 |
| $BAF_{fish}$ (L/kg FW)                 | --   | ND       |

**TABLE A-1b-47**

**CHEMICAL-SPECIFIC INPUTS FOR  
N-NITROSODIETHYLAMINE (55-18-5)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>                                      | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | --  | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | U.S. EPA (1997b)  | 1.52E+02     |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --  | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | Value based on <i>Oral CSF</i> assuming route-to-route extrapolation. | 1.52E+02     |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.                          | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-48

**CHEMICAL-SPECIFIC INPUTS FOR  
N-NITROSODIMETHYLAMINE (62-75-9)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neill, Smith, and Heckelman (1989)   | 74.08    |
| $T_m$ (K)                              | Budavari, O'Neill, Smith, and Heckelman (1989)   | ND       |
| $V_p$ (atm)                            | Geometric mean value cited in U.S. EPA (1994c).  | 7.10E-03 |
| $S$ (mg/L)                             | Geometric mean value cited in U.S. EPA (1994c).  | 1.00E+06 |
| $H$ (atm·m <sup>3</sup> /mol)          | U.S. EPA (1995g)   | 5.30-E07 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-06 |
| $K_{ow}$ (unitless)                    | Geometric mean value cited in U.S. EPA (1994c).  | 2.70E-01 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value was calculated by using the correlation equation with $K_{ow}$ for all nonionizing organics except phthalates, PAHs, dioxins, and furans as cited in U.S. EPA (1994c). $K_{oc}$ value was calculated by using the recommended $K_{ow}$ value that is provided in this table.  | 2.80E-01 |
| $Kd_s$ (mL/g)                          | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed fraction organic carbon of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 2.80E-03 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 2.10E-02 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 2.70E-04 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 2.50E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 5.90E-01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 2.70E-05 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 7.40E-01 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |



**TABLE A-1b-48**

**CHEMICAL-SPECIFIC INPUTS FOR  
N-NITROSODIMETHYLAMINE (62-75-9)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>             | <b>Value</b> |
|---|--|--------------|
| <b>Health Benchmarks</b>                        |  |              |
| <i>RfD</i> (mg/kg/day)                          | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | U.S. EPA (1997b)                             | 5.10E+01     |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --   | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | U.S. EAP (1997c)                             | 4.90E+01     |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                              | --   | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-49

**CHEMICAL-SPECIFIC INPUTS FOR  
N-NITROSOMETHYLETHYLAMINE (10595-95-6)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neill, Smith, and Heckelman (1989)   | 88.13    |
| $T_m$ (K)                              | Budavari, O'Neill, Smith, and Heckelman (1989)   |          |
| $V_p$ (atm)                            | Geometric mean value cited in U.S. EPA (1994c).  | 3.00E-03 |
| $S$ (mg/L)                             | Geometric mean value cited in U.S. EPA (1994c).  | 3.00E+05 |
| $H$ (atm·m <sup>3</sup> /mol)          | U.S. EPA (1995g)   | 8.90E-07 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-06 |
| $K_{ow}$ (unitless)                    | Geometric mean value cited in U.S. EPA (1994c).  | 7.60E-01 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value was calculated by using the correlation equation with $K_{ow}$ for all nonionizing organics except phthalates, PAHs, dioxins, and furans as cited in U.S. EPA (1994c). $K_{oc}$ value was calculated by using the recommended $K_{ow}$ value that is provided in this table.  | 7.60E-01 |
| $Kd_s$ (mL/g)                          | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed fraction organic carbon of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 7.60E-03 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 5.70E-02 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 4.50E-04 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 3.00E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 7.20E-01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 7.60E-05 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 3.10E+00 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-49**

**CHEMICAL-SPECIFIC INPUTS FOR  
N-NITROSOMETHYLETHYLAMINE (10595-95-6)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>                                      | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | --  | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | U.S. EPA (1997b)  | 2.20E+01     |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --  | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | Value based on <i>Oral CSF</i> assuming route-to-route extrapolation. | 2.20E+01     |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.                          | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA= Not applicable, ND= No data available

**TABLE A-1b-50**

**CHEMICAL-SPECIFIC INPUTS FOR  
N-NITROSOMORPHOLINE (59-89-2)**

| <b>Parameter</b>                       | <b>Reference and Explanation</b>                                  | <b>Value</b> |
|--|---|--------------|
| <b>Chemical/Physical Properties</b>    |   |              |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)                     | 116.11       |
| $T_m$ (K)                              | Budavari, O'Neil, Smith, and Heckelman (1989)                     | 302.15       |
| $V_p$ (atm)                            | --  | NA           |
| $S$ (mg/L)                             | --  | NA           |
| $H$ (atm·m <sup>3</sup> /mol)          | --  | NA           |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 1.00E-02     |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 1.00E-05     |
| $K_{ow}$ (unitless)                    | --  | NA           |
| $K_{oc}$ (mL/g)                        | --  | NA           |
| $Kd_s$ (cm <sup>3</sup> /g)            | --  | NA           |
| $Kd_{sw}$ (L/Kg)                       | --  | NA           |
| <b>Dermal Exposure Factors</b>         |   |              |
| $K_p$ (cm/hr)                          | $K_p$ value was obtained from U.S. EPA (1992b).                   | ND           |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).                     | 4.50E-01     |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).                   | 1.10E+00     |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).                     | 3.60E-05     |
| <b>Biotransfer Factors for Animals</b> |   |              |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --  | NA           |
| $BAF_{fish}$ (L/kg FW)                 | --  | NA           |

**TABLE A-1b-50**

**CHEMICAL-SPECIFIC INPUTS FOR  
N-NITROSOMORPHOLINE (59-89-2)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>             | <b>Value</b> |
|---|--|--------------|
| <b>Health Benchmarks</b>                        |  |              |
| <i>RfD</i> (mg/kg/day)                          | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --   | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                              | --   | ND           |

Note: NA= Not applicable, ND= No data available

TABLE A-1b-51

**CHEMICAL-SPECIFIC INPUTS FOR  
N-NITROSOPIPERIDINE (100-75-4)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 114.5    |
| $T_m$ (K)                              | --   | ND       |
| $V_p$ (atm)                            | Geometric mean value cited in U.S. EPA (1994c).  | 1.90E-04 |
| $S$ (mg/L)                             | Geometric mean value cited in U.S. EPA (1994c).  | 1.50E+05 |
| $H$ (atm·m <sup>3</sup> /mol)          | U.S. EPA (1995g)   | 1.40E-07 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-06 |
| $K_{ow}$ (unitless)                    | Geometric mean value cited in U.S. EPA (1994c).  | 4.30E+00 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value was calculated by using the correlation equation with $K_{ow}$ for phthalates and PAHs, / all nonionizing organics except phthalates, PAHs, dioxins, and furans, cited in U.S. EPA (1994c). $K_{oc}$ value was calculated by using the recommended $K_{ow}$ value that is provided in this table.   | 4.20E+00 |
| $Kd_s$ (mL/g)                          | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 4.20E-02 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 3.15E-01 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 1.10E-03 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 4.40E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 1.00E+00 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 4.30E-04 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 8.20E-01 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-51**

**CHEMICAL-SPECIFIC INPUTS FOR  
N-NITROSOPIPERIDINE (100-75-4)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>                                      | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | --  | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | U.S. EPA (1997b)  | 3.80E+01     |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --  | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | Value based on <i>Oral CSF</i> assuming route-to-route extrapolation. | 3.80E+01     |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.                          | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-52

**CHEMICAL-SPECIFIC INPUTS FOR  
N-NITROSOPYRROLIDINE (930-55-2)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 100.11   |
| $T_m$ (K)                              | --   |          |
| $Vp$ (atm)                             | $Vp$ value cited in Montgomery and Weldom (1991).  | 2.30E-04 |
| $S$ (mg/L)                             | $S$ value cited in U.S. EPA (1995g).   | 7.80E+05 |
| $H$ (atm·m <sup>3</sup> /mol)          | U.S. EPA (1995g)   | 2.90E-08 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 7.36E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 1.04E-05 |
| $K_{ow}$ (unitless)                    | $K_{ow}$ value cited in U.S. EPA (1995g).  | 6.50E-01 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value cited in U.S. EPA (1995g).  | 6.50E-01 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 6.50E-03 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 4.90E-02 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 3.40E-04 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 3.60E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 8.60E-01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 6.50E-05 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCF_s$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Caroll (1980)  | 1.90E-01 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |



**TABLE A-1b-52**

**CHEMICAL-SPECIFIC INPUTS FOR  
N-NITROSPYRROLIDINE (930-55-2)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>                                      | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | --  | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | U.S.EPA (1997b)   | 2.10E+00     |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --  | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | Value based on <i>Oral CSF</i> assuming route-to-route extrapolation. | 2.10E+00     |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.                          | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-53

**CHEMICAL-SPECIFIC INPUTS FOR  
OCTAMETHYLPYROPHOSPHORAMIDE (152-16-9)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 286.26   |
| $T_m$ (K)                              | --   |          |
| $Vp$ (atm)                             | $Vp$ value cited in U.S. EPA (1995g).  | 1.30E-06 |
| $S$ (mg/L)                             | $S$ value cited in U.S. EPA (1995g).   | 1.00E+06 |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value cited in U.S. EPA (1995g).   | 3.80E-10 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 1.80E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 5.62E-06 |
| $K_{ow}$ (unitless)                    | $K_{ow}$ value cited in U.S. EPA (1995g).  | 3.00E-01 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value cited in U.S. EPA (1995g).  | 3.10E-01 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 3.10E-03 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 2.30E-02 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 1.50E-05 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 4.90E+00 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 1.20E+01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 3.00E-05 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 1.00E-01 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-53**

**CHEMICAL-SPECIFIC INPUTS FOR  
OCTAMETHYLPYROPHOSPHORAMIDE (152-16-9)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S.EPA (1997b)   | 2.00E-03     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 7.00E-03     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-54

**CHEMICAL-SPECIFIC INPUTS FOR  
PARATHION (ETHYL) (56-38-2)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 291.27   |
| $T_m$ (K)                              | --   |          |
| $V_p$ (atm)                            | $V_p$ value cited in U.S. EPA (1995g).   | 1.30E-08 |
| $S$ (mg/L)                             | $S$ value cited in U.S. EPA (1995g).   | 6.50E+00 |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value cited in U.S. EPA (1995g).   | 5.70E-07 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 1.70E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 5.79E-06 |
| $K_{ow}$ (unitless)                    | $K_{ow}$ value cited in U.S. EPA (1995g).  | 6.80E+03 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value cited in U.S. EPA (1995g).  | 5.80E+03 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 5.80E+01 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 4.35E+02 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 1.70E-02 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 5.20E+00 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 3.90E+01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 6.80E-01 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | BCFs were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 2.70E+02 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-54**

**CHEMICAL-SPECIFIC INPUTS FOR  
PARATHION (ETHYL) (56-38-2)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997b)  | 6.00E-03     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 2.10E-02     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | Ambient Water Quality Criteria, U.S. EPA, Office of Water.  | 1.3E-02      |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-55

**CHEMICAL-SPECIFIC INPUTS FOR  
PENTACHLOROETHANE (76-01-7)**

| Parameter                              | Reference and Explanation   | Value     |
|--|---|-----------|
| <b>Chemical/Physical Properties</b>    |   |           |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)                     | 202.31    |
| $T_m$ (K)                              | Budavari, O'Neil, Smith, and Heckelman (1989)                     | 244.15    |
| $V_p$ (atm)                            | --  | ND        |
| $S$ (mg/L)                             | --  | ND        |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).   | 1.815E-03 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 6.60E-02  |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 7.30E-06  |
| $K_{ow}$ (unitless)                    | --  | ND        |
| $K_{oc}$ (mL/g)                        | --  | ND        |
| $Kd_s$ (cm <sup>3</sup> /g)            | --  | ND        |
| $Kd_{sw}$ (L/Kg)                       | --  | ND        |
| <b>Dermal Exposure Factors</b>         |   |           |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).                    | 1.63E-02  |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).                     | 1.50E+00  |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).                   | 4.05E+00  |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).                     | 1.12E-01  |
| <b>Biotransfer Factors for Animals</b> |   |           |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --  | ND        |
| $BAF_{fish}$ (L/kg FW)                 | --  | ND        |

**TABLE A-1b-55**

**CHEMICAL-SPECIFIC INPUTS FOR  
PENTACHLOROETHANE (76-01-7)**

| <b>Parameter</b>                                   | <b>Reference and Explanation</b>             | <b>Value</b> |
|--|--|--------------|
| <b>Health Benchmarks</b>                           |  |              |
| <i>RfD</i> (mg/kg/day)                             | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>          | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                    | --   | ND           |
| <i>Inhalation CSF</i><br>(mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>   | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                                 |  | ND           |

Note: NA = Not applicable, ND = No data available

**TABLE A-1b-56**

**CHEMICAL-SPECIFIC INPUTS FOR  
PHENACETIN (62-44-2)**

| <b>Parameter</b>                           | <b>Reference and Explanation</b>  | <b>Value</b>     |
|--|---|------------------|
| <b>Chemical/Physical Properties</b>        |   |                  |
| <i>MW</i> (g/mole)                         | Budavari, O'Neill, Smith, and Heckelman (1989)  | 179.21           |
| <i>T<sub>m</sub></i> (K)                   | Budavari, O'Neill, Smith, and Heckelman (1989)  | 407.15 to 408.15 |
| <i>V<sub>p</sub></i> (atm)                 | --  | ND               |
| <i>S</i> (g/1310ml of H <sub>2</sub> O)    | <i>S</i> value cited in U.S. EPA (1995b).   | 1.0              |
| <i>H</i> (atm·m <sup>3</sup> /mol)         | <i>H</i> value was calculated by using the theoretical equation from Lyman, Reehl, and Rosenblatt (1982), which defines the constant. Recommended value was calculated by using the <i>MW</i> , <i>S</i> , and <i>V<sub>p</sub></i> values that are provided in this table. | 1.41E-06         |
| <i>D<sub>a</sub></i> (cm <sup>2</sup> /s)  | <i>D<sub>a</sub></i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 5.70E-02         |
| <i>D<sub>w</sub></i> (cm <sup>2</sup> /s)  | <i>D<sub>w</sub></i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 6.82E-06         |
| <i>K<sub>ow</sub></i> (unitless)           | --  | ND               |
| <i>K<sub>oc</sub></i> (mL/g)               | --  | ND               |
| <i>Kd<sub>s</sub></i> (cm <sup>3</sup> /g) | --  | ND               |
| <i>Kd<sub>sw</sub></i> (L/Kg)              | --  | ND               |
| <b>Dermal Exposure Factors</b>             |   |                  |
| <i>K<sub>p</sub></i> (cm/hr)               | <i>K<sub>p</sub></i> value was obtained from U.S. EPA (1992b).  | 2.03E-03         |
| <i>t</i>                                   | <i>t</i> value was obtained from U.S. EPA (1992b).  | 1.08E+00         |
| <i>t*</i>                                  | <i>t*</i> value was obtained from U.S. EPA (1992b).   | 2.60E+00         |
| <i>B</i>                                   | <i>B</i> value was obtained from U.S. EPA (1992b).  | 3.80E-03         |
| <b>Biotransfer Factors for Animals</b>     |   |                  |
| <i>BCF<sub>fish</sub></i> (L/kg FW tissue) | --  | ND               |
| <i>BAF<sub>fish</sub></i> (L/kg FW)        | --  | ND               |



**TABLE A-1b-56**

**CHEMICAL-SPECIFIC INPUTS FOR  
PHENACETIN (62-44-2)**

| <b>Parameter</b>                                   | <b>Reference and Explanation</b>             | <b>Value</b> |
|--|--|--------------|
| <b>Health Benchmarks</b>                           |  |              |
| <i>RfD</i> (mg/kg/day)                             | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>          | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                    | --   | ND           |
| <i>Inhalation CSF</i><br>(mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>   | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                                 | --   | ND           |

Note: NA= Not applicable, ND= No data available

TABLE A-1b-57

**CHEMICAL-SPECIFIC INPUTS FOR  
PHENYL MERCURIC ACETATE (62-38-4)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 336.75   |
| $T_m$ (K)                              | --   |          |
| $V_p$ (atm)                            | $V_p$ value cited in U.S. EPA (1995g).   | 4.00E-09 |
| $S$ (mg/L)                             | $S$ value cited in U.S. EPA (1995g).   | 4.40E+03 |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value cited in U.S. EPA (1995g).   | 2.04E-11 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 1.58E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 4.62E-06 |
| $K_{ow}$ (unitless)                    | $K_{ow}$ value cited in U.S. EPA (1995g).  | 1.70E+02 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value cited in U.S. EPA (1995g).  | 1.60E+02 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 1.60E+00 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 1.20E+01 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 6.40E-04 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 9.90E+00 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 2.40E+01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 1.70E-02 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$<br>(L/kg FW tissue)       | U.S. EPA (1995g)   | 1.50E+01 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-57**

**CHEMICAL-SPECIFIC INPUTS FOR  
PHENYL MERCURIC ACETATE (62-38-4)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S.EPA (1997b)   | 8.00E-05     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 2.80E-04     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-58

**CHEMICAL-SPECIFIC INPUTS FOR  
1,3-PHENYLENEDIAMINE (108-45-2)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 108.6    |
| $T_m$ (K)                              | --   |          |
| $Vp$ (atm)                             | $Vp$ value cited in U.S. EPA (1995g).  | 3.00E-05 |
| $S$ (mg/L)                             | $S$ value cited in U.S. EPA (1995g).   | 3.50E+05 |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value cited in U.S. EPA (1995g).   | 9.20E-09 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 6.63E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 9.88E-06 |
| $K_{ow}$ (unitless)                    | $K_{ow}$ value cited in U.S. EPA (1995g).  | 1.10E+00 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value cited in U.S. EPA (1995g).  | 1.10E+00 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 1.10E-02 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 8.25E-02 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 4.50E-04 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 4.00E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 9.60E-01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 1.10E-04 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg, FW tissue)         | U.S. EPA (1995g)   | 2.90E-01 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-58**

**CHEMICAL-SPECIFIC INPUTS FOR  
1,3-PHENYLENEDIAMINE (108-45-2)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997b)  | 6.00E-03     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 2.10E-02     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-59

**CHEMICAL-SPECIFIC INPUTS FOR  
A-PICOLINE (109-06-8)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 93.12    |
| $T_m$ (°K)                             | --   | 203.15   |
| $Vp$ (mm@°C)                           | $Vp$ value cited in U.S. EPA (1995g).  | 8.00E+00 |
| $S$ (mg/L)                             | $S$ value cited in U.S. EPA (1995g).   |          |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value cited in U.S. EPA (1995g).   | 4.10E-04 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 7.50E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 9.60E-06 |
| $K_{ow}$ (unitless)                    | $K_{ow}$ value cited in U.S. EPA (1995g).  | 2.00E+06 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value cited in U.S. EPA (1995g).  | 1.50E+06 |
| $Kd_s$ (mL/g)                          | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 1.50E+04 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 1.12E+05 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 3.16E-03 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 3.24E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 7.76E-01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 1.29E-03 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | --   | ND       |
| $BAF_{fish}$ (L/kg FW)                 | --   | ND       |

**TABLE A-1b-59**

**CHEMICAL-SPECIFIC INPUTS FOR  
A-PICOLINE (109-06-8)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>             | <b>Value</b> |
|---|--|--------------|
| <b>Health Benchmarks</b>                        |  |              |
| <i>RfD</i> (mg/kg/day)                          | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --   | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                              | --   | ND           |

Note: NA= Not applicable, ND= No data available

**TABLE A-1b-60**

**CHEMICAL-SPECIFIC INPUTS FOR  
POLYCHLORINATED BIPHENYLS (AROCLORS) (1336-36-3)**

| <b>Parameter</b>                           | <b>Reference and Explanation</b>  | <b>Value</b> |
|--|---|--------------|
| <b>Chemical/Physical Properties</b>        |   |              |
| <i>MW</i> (g/mole)                         | Budavari, O'Neil, Smith, and Heckelman (1989)   | 328          |
| <i>T<sub>m</sub></i> (°K)                  | --  |              |
| <i>V<sub>p</sub></i> (atm)                 | <i>V<sub>p</sub></i> value cited in U.S. EPA (1995g).   | 1.00E-07     |
| <i>S</i> (mg/L)                            | <i>S</i> value cited in U.S. EPA (1995g).   | 8.00E-02     |
| <i>H</i> (atm·m <sup>3</sup> /mol)         | <i>H</i> value cited in U.S. EPA (1995g).   | 2.60E-03     |
| <i>D<sub>a</sub></i> (cm <sup>2</sup> /s)  | <i>D<sub>a</sub></i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-02     |
| <i>D<sub>w</sub></i> (cm <sup>2</sup> /s)  | <i>D<sub>w</sub></i> value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 1.00E-05     |
| <i>K<sub>ow</sub></i> (unitless)           | <i>K<sub>ow</sub></i> value cited in U.S. EPA (1995g).  | 2.00E+06     |
| <i>K<sub>oc</sub></i> (mL/g)               | <i>K<sub>oc</sub></i> value cited in U.S. EPA (1998c).  | 9.83E+04     |
| <i>Kd<sub>s</sub></i> (mL/g)               | <i>Kd<sub>s</sub></i> value was calculated by using the correlation equation with <i>K<sub>oc</sub></i> that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate <i>Kd<sub>s</sub></i> , because the value varies, depending on the fraction of organic carbon in soil. Recommended <i>Kd<sub>s</sub></i> value was calculated by using the <i>K<sub>oc</sub></i> value that is provided in this table.   | 9.83E+02     |
| <i>Kd<sub>sw</sub></i> (L/Kg)              | <i>Kd<sub>sw</sub></i> value was calculated by using the correlation equation with <i>K<sub>oc</sub></i> that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate <i>Kd<sub>sw</sub></i> , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended <i>Kd<sub>sw</sub></i> value was calculated by using the <i>K<sub>oc</sub></i> value that is provided in this table. | 7.37E+03     |
| <b>Dermal Exposure Factors</b>             |   |              |
| <i>K<sub>p</sub></i> (cm/hr)               | <i>K<sub>p</sub></i> value was obtained from U.S. EPA (1992b).  | 1.30E+00     |
| <i>t</i>                                   | <i>t</i> value was obtained from U.S. EPA (1992b).  | 5.30E+00     |
| <i>t*</i>                                  | <i>t*</i> value was obtained from U.S. EPA (1992b).   | 2.50E+01     |
| <i>B</i>                                   | <i>B</i> value was obtained from U.S. EPA (1992b).  | 3.20E+02     |
| <b>Biotransfer Factors for Animals</b>     |   |              |
| <i>BCF<sub>fish</sub></i> (L/kg FW tissue) | --  | NA           |
| <i>BAF<sub>fish</sub></i> (L/kg FW)        |   | 6.70E+05     |



**TABLE A-1b-60**

**CHEMICAL-SPECIFIC INPUTS FOR  
POLYCHLORINATED BIPHENYLS (AROCLORS) (1336-36-3)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>                                      | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | --  | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | U.S. EPA (1997b)  | 2.00E+00     |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --  | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | Value based on <i>Oral CSF</i> assuming route-to-route extrapolation. | 2.00E+00     |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.                          | 5E-04        |
| Aquatic TRV (µg/l)                              | Ambient Water Quality Criteria, U.S. EPA, Office of Water.            | 1.4E-02      |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-61

**CHEMICAL-SPECIFIC INPUTS FOR  
TETRAETHYL DITHIOPYROPHOSPHATE (SULFOTEP) (3689-24-5)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 322.31   |
| $T_m$ (K)                              | --   |          |
| $Vp$ (atm)                             | $Vp$ value cited in U.S. EPA (1995g).  | 3.30E-07 |
| $S$ (mg/L)                             | $S$ value cited in U.S. EPA (1995g).   | 2.50E+01 |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value cited in U.S. EPA (1995g).   | 4.20E-06 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-06 |
| $K_{ow}$ (unitless)                    | $K_{ow}$ value cited in U.S. EPA (1995g).  | 6.80E+03 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value cited in U.S. EPA (1995g).  | 5.80E+03 |
| $Kd_s$ (mL/g)                          | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 5.80E+01 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 4.35E+02 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 1.10E-02 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 8.10E+00 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 6.00E+01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 6.80E-01 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | U.S. EPA (1995g)   | 2.80E+02 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-61**

**CHEMICAL-SPECIFIC INPUTS FOR  
TETRAETHYL DITHIOPYROPHOSPHATE (SULFOTEP) (3689-24-5)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997b)  | 5.00E-04     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 1.75E-03     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

**TABLE A-1b-62**

**CHEMICAL-SPECIFIC INPUTS FOR  
THIONAZIN (297-97-2)**

| <b>Parameter</b>                       | <b>Reference and Explanation</b>                                  | <b>Value</b> |
|--|---|--------------|
| <b>Chemical/Physical Properties</b>    |   |              |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)                     | 248.26       |
| $T_m$ (K)                              | Budavari, O'Neil, Smith, and Heckelman (1989)                     | 271.45       |
| $Vp$ (mmHg)                            | U.S. EPA (1995g)  | 3.00E-03     |
| $S$ (mg/L)                             | --  | ND           |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).   | 8.60E-07     |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 8.00E-02     |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 8.00E-06     |
| $K_{ow}$ (unitless)                    | --  | ND           |
| $K_{oc}$ (mL/g)                        | --  | ND           |
| $Kd_s$ (cm <sup>3</sup> /g)            | --  | ND           |
| $Kd_{sw}$ (L/Kg)                       | --  | ND           |
| <b>Dermal Exposure Factors</b>         |   |              |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).                    | ND           |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).                     | 2.86E+00     |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).                   | ND           |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).                     | ND           |
| <b>Biotransfer Factors for Animals</b> |   |              |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --  | ND           |
| $BAF_{fish}$ (L/kg FW)                 | --  | ND           |

**TABLE A-1b-62**

**CHEMICAL-SPECIFIC INPUTS FOR  
THIONAZIN (297-97-2)**

| <b>Parameter</b>                                   | <b>Reference and Explanation</b>             | <b>Value</b> |
|--|--|--------------|
| <b>Health Benchmarks</b>                           |  |              |
| <i>RfD</i> (mg/kg/day)                             | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>          | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                    | --   | ND           |
| <i>Inhalation CSF</i><br>(mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>   | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                                 | --   | ND           |

Note: NA = Not applicable, ND = No data available

**TABLE A-1b-63**

**CHEMICAL-SPECIFIC INPUTS FOR  
TIN (7440-31-5)**

| <b>Parameter</b>                       | <b>Reference and Explanation</b>                                  | <b>Value</b> |
|--|---|--------------|
| <b>Chemical/Physical Properties</b>    |   |              |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)                     | 118.69       |
| $T_m$ (K)                              | Budavari, O'Neil, Smith, and Heckelman (1989)                     | 505.05       |
| $Vp$ (atm)                             | --  | ND           |
| $S$ (mg/L)                             | --  | ND           |
| $H$ (atm·m <sup>3</sup> /mol)          | --  | ND           |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 8.00E-02     |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d). | 8.00E-06     |
| $K_{ow}$ (unitless)                    | --  | ND           |
| $K_{oc}$ (mL/g)                        | --  | ND           |
| $Kd_s$ (cm <sup>3</sup> /g)            | --  | ND           |
| $Kd_{sw}$ (L/Kg)                       | --  | ND           |
| <b>Dermal Exposure Factors</b>         |   |              |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).                    | 1.00E-03     |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).                     | ND           |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).                   | ND           |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).                     | ND           |
| <b>Biotransfer Factors for Animals</b> |   |              |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --  | ND           |
| $BAF_{fish}$ (L/kg FW)                 | --  | ND           |

**TABLE A-1b-63**

**CHEMICAL-SPECIFIC INPUTS FOR  
TIN (7440-31-5)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997b)  | 6.00E-01     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 2.10E+00     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-64

**CHEMICAL-SPECIFIC INPUTS FOR  
2,4-TOLUENEDIAMINE (95-80-7)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 122.17   |
| $T_m$ (K)                              | --   |          |
| $Vp$ (atm)                             | $Vp$ value cited in U.S. EPA (1995g).  | 1.10E-07 |
| $S$ (mg/L)                             | $S$ value cited in U.S. EPA (1995g).   | 7.50E+03 |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value cited in U.S. EPA (1995g).   | 7.92E-10 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 5.69E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 9.05E-06 |
| $K_{ow}$ (unitless)                    | $K_{ow}$ value cited in U.S. EPA (1995g).  | 2.50E+00 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value cited in U.S. EPA (1995g).  | 2.50E+00 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 2.50E-02 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 1.88E-01 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 6.60E-04 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 4.90E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 1.20E+00 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 2.50E-04 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$<br>(unitless FW tissue)   | U.S. EPA (1995g)   | 4.60E+01 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |



**TABLE A-1b-64**

**CHEMICAL-SPECIFIC INPUTS FOR  
2,4-TOLUENEDIAMINE (95-80-7)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>                                      | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | --  | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | U.S. EPA (1997b)  | 3.20E+00     |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --  | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | Value based on <i>Oral CSF</i> assuming route-to-route extrapolation. | 3.20E+00     |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.                          | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note:NA = Not applicable, ND = No data available

TABLE A-1b-65

CHEMICAL-SPECIFIC INPUTS FOR  
P-TOLUIDINE (106-49-0)

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 107.15   |
| $T_m$ (°K)                             | Budavari, O'Neil, Smith, and Heckelman (1989)  | 317.65   |
| $Vp$ (atm)                             | $Vp$ value cited in U.S. EPA (1995g).  | 4.30E-04 |
| $S$ (mg/L)                             | $S$ value cited in U.S. EPA (1995g).   | 7.60E+03 |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value cited in U.S. EPA (1995g).   | 6.10E-06 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 6.97E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 9.43E-06 |
| $K_{ow}$ (unitless)                    | $K_{ow}$ value cited in U.S. EPA (1995g).  | 2.50E+01 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value cited in U.S. EPA (1995g).  | 2.40E+01 |
| $Kd_s$ (mL/g)                          | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 2.40E-01 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 1.80E+00 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 4.20E-03 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 3.90E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 9.50E-01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 2.50E-03 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | U.S. EPA (1995g)   | 3.50E+00 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-65**

**CHEMICAL-SPECIFIC INPUTS FOR  
P-TOLUIDINE (106-49-0)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>                                      | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | --  | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | U.S. EPA (1997b)  | 1.90E-01     |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --  | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | Value based on <i>Oral CSF</i> assuming route-to-route extrapolation. | 1.90E-01     |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.                          | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-66

**CHEMICAL-SPECIFIC INPUTS FOR  
TOXAPHENE (CHLORINATED CAMPHENES) (8001-35-2)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 414      |
| $T_m$ (K)                              | Budavari, O'Neil, Smith, and Heckelman (1989)  | 350.65   |
| $Vp$ (atm)                             | $Vp$ value cited in U.S. EPA (1995g).  | 4.30E-04 |
| $S$ (mg/L)                             | $S$ value cited in U.S. EPA (1995g).   | 6.79E-01 |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value cited in U.S. EPA (1995g).   | 3.40E-06 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 1.16E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 4.34E-06 |
| $K_{ow}$ (unitless)                    | $K_{ow}$ value cited in U.S. EPA (1995g).  | 3.20E+05 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value cited in U.S. EPA (1995g).  | 2.60E+05 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 2.60E+03 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 1.95E+04 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 4.60E-02 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 2.90E+01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 1.40E+02 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 3.20E+01 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | --   | NA       |
| $BAF_{fish}$ (L/kg FW)                 | (U.S. EPA 1995g)   | 2.10E+06 |

**TABLE A-1b-66**

**CHEMICAL-SPECIFIC INPUTS FOR  
TOXAPHENE (CHLORINATED CAMPHENES) (8001-35-2)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>                                      | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | --  | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | U.S. EPA (1997b)  | 1.10E+00     |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --  | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | Value based on <i>Oral CSF</i> assuming route-to-route extrapolation. | 1.10E+00     |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.                          | 3E-03        |
| Aquatic TRV (µg/l)                              | Ambient Water Quality Criteria, U.S. EPA, Office of Water.            | 2E-04        |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-67

**CHEMICAL-SPECIFIC INPUTS FOR  
1,1,2-TRICHLORO-1,2,2-TRIFLUOROETHANE (76-13-1)**

| Parameter                              | Reference and Explanation  | Value     |
|--|--|-----------|
| <b>Chemical/Physical Properties</b>    |  |           |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 187.38    |
| $T_m$ (°K)                             | --   |           |
| $Vp$ (atm)                             | $Vp$ value cited in U.S. EPA (1995g).  | 4.80E-01  |
| $S$ (mg/L)                             | $S$ value cited in U.S. EPA (1995g).   | 1.70E+02  |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value cited in U.S. EPA (1995g).   | 4.815E-01 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 7.80E-02  |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.20E-06  |
| $K_{ow}$ (unitless)                    | $K_{ow}$ value cited in U.S. EPA (1995g).  | 1.40E+03  |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value cited in U.S. EPA (1995g).  | 1.30E+03  |
| $Kd_s$ (mL/g)                          | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 1.30E+01  |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 9.81E+01  |
| <b>Dermal Exposure Factors</b>         |  |           |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 2.40E-02  |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 1.20E+00  |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 4.10E+00  |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 1.40E-01  |
| <b>Biotransfer Factors for Animals</b> |  |           |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 8.20E+01  |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA        |

**TABLE A-1b-67**

**CHEMICAL-SPECIFIC INPUTS FOR  
1,1,2-TRICHLORO-1,2,2-TRIFLUOROETHANE (76-13-1)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>             | <b>Value</b> |
|---|--|--------------|
| <b>Health Benchmarks</b>                        |  |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997b)                             | 3.00E+01     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | U.S. EPA (1997c)                             | 8.57E+00     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                              | --   | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-68

**CHEMICAL-SPECIFIC INPUTS FOR  
TRICHLOROPHENOXY PROPIONIC ACID (SILVEX) (93-72-1)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 269.51   |
| $T_m$ (K)                              | --   |          |
| $Vp$ (atm)                             | $Vp$ value cited in U.S. EPA (1995g).  | 6.80E-09 |
| $S$ (mg/L)                             | $S$ value cited in U.S. EPA (1995g).   | 1.40E+02 |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value cited in U.S. EPA (1995g).   | 1.30E-08 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-06 |
| $K_{ow}$ (unitless)                    | $K_{ow}$ value cited in U.S. EPA (1995g).  | 2.60E+03 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value cited in U.S. EPA (1995g).  | 2.30E+03 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 2.30E+01 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 1.71E+02 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 1.10E-02 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 3.90E+00 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 1.90E+01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 2.60E-01 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Caroll (1980)   | 1.30E+02 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |



**TABLE A-1b-68**

**CHEMICAL-SPECIFIC INPUTS FOR  
TRICHLOROPHENOXY PROPIONIC ACID (SILVEX) (93-72-1)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997b)  | 8.00E-03     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 2.80E-02     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | 5E-02        |
| Aquatic TRV (µg/l)                              | --.   | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-69

**CHEMICAL-SPECIFIC INPUTS FOR  
2,4,5-TRICHLOROPHENOXYACETIC ACID (93-76-5)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 255.49   |
| $T_m$ (K)                              | --   |          |
| $Vp$ (atm)                             | $Vp$ value cited in U.S. EPA (1995g).  | 9.10E-10 |
| $S$ (mg/L)                             | $S$ value cited in U.S. EPA (1995g).   | 2.80E+02 |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value cited in U.S. EPA (1995g).   | 8.68E-09 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 8.00E-06 |
| $K_{ow}$ (unitless)                    | $K_{ow}$ value cited in U.S. EPA (1995g).  | 2.00E+03 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value cited in U.S. EPA (1995g).  | 1.80E+03 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 1.80E+01 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 1.35E+02 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 1.20E-02 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 3.20E+00 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 1.30E+01 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 2.00E-01 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (unitless FW tissue)      | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Caroll (1980)   | 7.80E+01 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-69**

**CHEMICAL-SPECIFIC INPUTS FOR  
2,4,5-TRICHLOROPHENOXYACETIC ACID (93-76-5)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997a)  | 1.00E-02     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 3.50E-02     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note:NA = Not applicable, ND = No data available

TABLE A-1b-70

**CHEMICAL-SPECIFIC INPUTS FOR  
O,O,O-TRIETHYLPHOSPHOROTHIA TE (126-68-1)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | CRC Handbook (1995)  | 198.22   |
| $T_m$ (°K)                             | --   | ND       |
| $Vp$ (atm)                             | --   | ND       |
| $S$ (mg/L)                             | --   | ND       |
| $H$ (atm·m <sup>3</sup> /mol)          | --   | ND       |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was calculated using the equation cited in U.S. EPA (1996a). | 3.17E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was calculated using the equation cited in U.S. EPA (1996a). | 6.23E-06 |
| $K_{ow}$ (unitless)                    | --   | ND       |
| $K_{oc}$ (mL/g)                        | --   | ND       |
| $Kd_s$ (mL/g)                          | --   | ND       |
| $Kd_{sw}$ (L/Kg)                       | --   | ND       |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).                           | ND       |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).                            | 1.42E+00 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).                          | ND       |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).                            | ND       |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --   | ND       |
| $BAF_{fish}$ (L/kg FW)                 | --   | ND       |

**TABLE A-1b-70**

**CHEMICAL-SPECIFIC INPUTS FOR  
O,O,O-TRIETHYLPHOSPHOROTHIA TE (126-68-1)**

| <b>Parameter</b>                                   | <b>Reference and Explanation</b>             | <b>Value</b> |
|--|--|--------------|
| <b>Health Benchmarks</b>                           |  |              |
| <i>RfD (water)</i><br>(mg/kg/day)                  | --   | ND           |
| <i>RfD (food)</i> (mg/kg/day)                      | --   | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>          | --   | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                    | --   | ND           |
| <i>Inhalation CSF</i><br>(mg/kg/day) <sup>-1</sup> | --   | ND           |
| <i>MCL</i>   | National Primary Drinking Water Regulations. | ND           |
| Aquatic TRV (µg/l)                                 | --   | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-71

**CHEMICAL-SPECIFIC INPUTS FOR  
TRIS(2,3-DIBROMOPROPYL) PHOSPHATE B19 (126-72-7)**

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 697.93   |
| $T_m$ (K)                              | --   | ND       |
| $Vp$ (atm)                             | $Vp$ value cited in U.S. EPA (1995g).  | 2.00E-07 |
| $S$ (mg/L)                             | $S$ value cited in U.S. EPA (1995g).   | 4.70E+00 |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value cited in U.S. EPA (1995g).   | 3.00E-05 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 5.50E-03 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 3.66E-06 |
| $K_{ow}$ (unitless)                    | $K_{ow}$ value cited in U.S. EPA (1995g).  | 3.20E+03 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value cited in U.S. EPA (1995g).  | 2.80E+03 |
| $Kd_s$ (cm <sup>3</sup> /g)            | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 2.80E+01 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 2.10E+02 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 3.30E-05 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 1.60E+03 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 8.60E+03 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 3.20E-01 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW tissue)          | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Caroll (1980)   | 1.60E+02 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-71**

**CHEMICAL-SPECIFIC INPUTS FOR  
TRIS(2,3-DIBROMOPROPYL) PHOSPHATE B19 (126-72-7)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>                                      | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | --  | ND           |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | U.S. EPA (1997b)  | 9.80E+00     |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | --  | ND           |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | Value based on <i>Oral CSF</i> assuming route-to-route extrapolation. | 9.80E+00     |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.                          | ND           |
| Aquatic TRV (µg/l)                              | --  | ND           |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-72

## CHEMICAL-SPECIFIC INPUTS FOR VANADIUM (7440-62-2)

| Parameter                              | Reference and Explanation                      | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | --   | 50.94    |
| $T_m$ (K)                              | --   |          |
| $Vp$ (atm)                             | --   | N/A      |
| $S$ (mg/L)                             | --   | N/A      |
| $H$ (atm·m <sup>3</sup> /mol)          | --   | NA       |
| $D_a$ (cm <sup>2</sup> /s)             | --   | NA       |
| $D_w$ (cm <sup>2</sup> /s)             | --   | NA       |
| $K_{ow}$ (unitless)                    | --   | N/A      |
| $K_{oc}$ (mL/g)                        | --   | N/A      |
| $Kd_s$ (cm <sup>3</sup> /g)            | U.S. EPA (1995g)                               | 5.00E+01 |
| $Kd_{sw}$ (L/Kg)                       | --   | 5.00E+01 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b). | 1.00E-03 |
| $t$                                    | --   | ND       |
| $t^*$                                  | --   | ND       |
| $B$                                    | --   | ND       |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$<br>(L/kg FW tissue)       | --   | NA       |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |



**TABLE A-1b-72**

**CHEMICAL-SPECIFIC INPUTS FOR VANADIUM (7440-62-2)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997b)  | 7.00E-03     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 2.45E-02     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | ND           |
| Aquatic TRV (µg/l)                              | U.S. EPA (1996c)  | 1.9E+01      |

Note: NA = Not applicable, ND = No data available

TABLE A-1b-73

CHEMICAL-SPECIFIC INPUTS FOR  
XYLENES (TOTAL) (1330-20-7)

| Parameter                              | Reference and Explanation  | Value    |
|--|--|----------|
| <b>Chemical/Physical Properties</b>    |  |          |
| $MW$ (g/mole)                          | Budavari, O'Neil, Smith, and Heckelman (1989)  | 106.17   |
| $T_m$ (°K)                             | --   |          |
| $V_p$ (atm)                            | $V_p$ value cited in U.S. EPA (1995g).   | 1.10E-02 |
| $S$ (mg/L)                             | $S$ value cited in U.S. EPA (1995g).   | 1.90E+02 |
| $H$ (atm·m <sup>3</sup> /mol)          | $H$ value cited in U.S. EPA (1995g).   | 6.00E-03 |
| $D_a$ (cm <sup>2</sup> /s)             | $D_a$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 7.14E-02 |
| $D_w$ (cm <sup>2</sup> /s)             | $D_w$ value was obtained from CHEMDAT8 database (U.S. EPA 1994d).  | 9.34E-06 |
| $K_{ow}$ (unitless)                    | $K_{ow}$ value cited in U.S. EPA (1995g).  | 1.50E+03 |
| $K_{oc}$ (mL/g)                        | $K_{oc}$ value cited in U.S. EPA (1995g).  | 1.30E+03 |
| $Kd_s$ (mL/g)                          | $Kd_s$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.01 in soil. Measured organic carbon in soil, specific to site conditions, should be used to calculate $Kd_s$ , because the value varies, depending on the fraction of organic carbon in soil. Recommended $Kd_s$ value was calculated by using the $K_{oc}$ value that is provided in this table.   | 1.30E+01 |
| $Kd_{sw}$ (L/Kg)                       | $Kd_{sw}$ value was calculated by using the correlation equation with $K_{oc}$ that is cited in U.S. EPA (1993) for an assumed organic carbon fraction of 0.075 in suspended sediment. Measured organic carbon in suspended sediment, specific to site conditions, should be used to calculate $Kd_{sw}$ , because the value varies, depending on the fraction of organic carbon in suspended sediment. Recommended $Kd_{sw}$ value was calculated by using the $K_{oc}$ value that is provided in this table. | 9.75E+01 |
| <b>Dermal Exposure Factors</b>         |  |          |
| $Kp$ (cm/hr)                           | $Kp$ value was obtained from U.S. EPA (1992b).   | 7.60E-02 |
| $t$                                    | $t$ value was obtained from U.S. EPA (1992b).  | 3.90E-01 |
| $t^*$                                  | $t^*$ value was obtained from U.S. EPA (1992b).  | 1.30E+00 |
| $B$                                    | $B$ value was obtained from U.S. EPA (1992b).  | 1.50E-01 |
| <b>Biotransfer Factors for Animals</b> |  |          |
| $BCF_{fish}$ (L/kg FW)                 | $BCFs$ were used for compounds with a log $K_{ow}$ value below 4.0, as cited in U.S. EPA (1995g). $BCF_{fish}$ value calculated using the correlation equation with $K_{ow}$ obtained from Veith, Macek, Petrocelli, and Carroll (1980)  | 7.50E+01 |
| $BAF_{fish}$ (L/kg FW)                 | --   | NA       |

**TABLE A-1b-73**

**CHEMICAL-SPECIFIC INPUTS FOR  
XYLENES (TOTAL) (1330-20-7)**

| <b>Parameter</b>                                | <b>Reference and Explanation</b>  | <b>Value</b> |
|---|---|--------------|
| <b>Health Benchmarks</b>                        |   |              |
| <i>RfD</i> (mg/kg/day)                          | U.S. EPA (1997b)  | 2.00E+00     |
| <i>Oral CSF</i> (mg/kg/day) <sup>-1</sup>       | --  | ND           |
| <i>RfC</i> (mg/m <sup>3</sup> )                 | Calculated from <i>RfD</i> using an inhalation rate of 20 m <sup>3</sup> /day and a human body weight of 70 kg. | 7.00E+00     |
| <i>Inhalation CSF</i> (mg/kg/day) <sup>-1</sup> | --  | ND           |
| <i>MCL</i>                                      | National Primary Drinking Water Regulations.  | 1.0E+01      |
| Aquatic TRV (µg/l)                              | U.S. EPA (1996c)  | 1.80E+00     |

Note: Not applicable, ND = No data available