



Dow AgroSciences

MRID 49940501 -

Dow AgroSciences LLC  
9330 Zionsville Road Indianapolis, IN 46268 USA

www.dowagro.com

February 29, 2016

Document Processing Desk (REGFEE)  
Office of Pesticide Programs  
U.S. EPA, Room S-4400  
One Potomac Yard (South Building)  
2777 South Crystal Drive  
Arlington, VA 22202-4501

**RE: Addendum to add additional minor crop to previously submitted application (December 8, 2015) for PRIA 3 M007: Request to Extend Exclusive Use Data Period for Spinetoram (62719-539)**

Dear Mr. Stephen Schiable:

Dow AgroSciences LLC (DAS) previously submitted an application to extend the Exclusive Use Data period for Spinetoram (62719-539) in December of 2015. During initial screening of the petition, EPA determined that one of the crops (tomatoes) in the petition was not considered a minor crop, and therefore could not be used within the petition. DAS was provided an opportunity to submit a write up on an additional minor crop, which could be added to the petition currently under review to replace the tomato write up. DAS is grateful for the opportunity to add the minor crop, and is herein submitting an addendum to the current M007 petition on file, with a write-up on avocados.

**Included in this submission:**

1. Transmittal document
2. Addendum to DAS petition to extend exclusive use data period for Spinetoram – PRIA M007 - Adding Avocado as minor crop to replace Tomato.
3. Complimentary copy of letter dated December 1, 2015.

Sincerely,

Jennifer Hughes  
U.S. Regulatory Manager  
317-337-7993  
JLHughes@dow.com

**Addendum to DAS petition to Extend Exclusive Use Data Period for Spinetoram (62719-539) –  
PRIA M007 - adding Avocado as minor crop to replace Tomatoes**

**DAS satisfies requirements II, III and IV (alternatives pose greater risk to environment/human health; managing pest resistance; and role in IPM) from FIFRA § 3(c)(1)(F)(ii) for the following Crop:**

**Avocado**

A key pest in the minor use crop, avocado (*Persea americana*), is avocado thrips, *Scirtothrips perseae* Nakahara. Avocado thrips were first discovered in June 1996 in Ventura County, California and has a highly restrictive host range, with larvae being found only on avocados (1). As an invasive pest, avocado thrips caused significant crop damage in avocado in 1997 and by 1999 the spread had stabilized in California with 95% of California acreage infested in 2002 (2). Avocado thrips management costs have reduced industry revenue by 12%, increased production costs by 4.5%, and economic models indicate long-term losses to the California avocado industry to be approximately \$4-8 million per year (5).

Avocado thrips have little effect on tree health but feed directly on immature fruit (3). Avocado thrips adults lay eggs within the underside of leaves, in young fruit, and in fruit petioles. The thrips develop through two larval and two pupal stages with six or more generations per year. Avocado thrips develop well under cooler temperatures, with populations finally being suppressed by warm, dry conditions. However, these higher temperatures occur late in the season and coincide with when fruit are larger and no longer susceptible to damage (3).

Prior to the introduction of avocado thrips, natural enemies were successful in keeping a variety of insect and mite pests below economically injurious levels (5). Thrips predator and parasitoids that may be present in avocado orchards include generalist predators such as lacewings (*Chrysoperla* spp), predatory thrips (*Frankliniella vespiformis*), and predatory mites (*Euseius hibisci*) (4). Two parasitic wasps are found in avocado, which parasitize thrips larvae, are *Thripobius semiluteus* and *Ceraninus menes*. For all of these predators and parasitoids that do occur in avocado orchards, the impact of them on avocado thrips is unknown (4). Worldwide, there are few examples of good biological control of pestiferous thrips species (10). Therefore, it is advisable to conserve beneficial arthropods by applying pesticides only when necessary (3,4).

In addition to the possibility of biological control to suppress avocado thrips, several cultural controls are recommended. It is encouraged to modify fertilization application regimes and pruning in a way that will reduce the length of time trees produce tender foliage and will thus decrease thrips abundance (3,6,7). Although the effectiveness to control thrips in commercial fields is uncertain, coarse organic mulches spread beneath trees in a thick layer may reduce survival of thrip pupae (7). Labor costs of adding mulch may not be justified for the purpose of exclusively suppressing thrips though (3).

Insecticidal controls are required to reduce crop loss from avocado thrips in avocado, when it is clear that economic levels of avocado thrips (approximately three to ten immature thrips per fruit) are present on a significant fruit set and are likely to cause economically significant damage (5,8,9). Pesticides in five

activity groups – abamectin (MOA Group 6), spinosad and spinetoram (MOA Group 5), spirotetramet (MOA Group 23), fenpropathrin (Group 3A), and sabadilla (MOA Group 1B) - are recommended for management of avocado thrips (3). University of California ranks these active ingredients according to having the greatest IPM fit, which encompasses being the most efficacious and least harmful to natural enemies, bees, and the environment (from greatest to least IPM benefit) – abamectin, spinetoram, spirotetramet, fenpropathrin, spinosad, and sabadilla (3). Insecticide Resistance Action Committee guidelines (<http://www.irac-online.org/>), which are also recommended by University of California, state that chemicals with a different mode-of-action group number should be rotated, and to not use products with the same mode-of-action group number more than twice per season to help prevent the development of resistance (3).

Spinosyns are a recommended for avocado thrips control, with spinetoram demonstrating even greater persistence and effectiveness than spinosad (3,10,13). Spinosad insecticide products were first introduced in the late 1990s and have a unique attribute of having exceptional activity against thrips (12). Spinetoram, a new and more active insecticide, was registered for use in 2008. Spinetoram represents a unique mode of action (Group 5 insecticides) with both contact and ingestion activity, and is one of the most effective insecticides to suppress thrips because it provides excellent control and it is a reduced risk insecticide that does not suppress beneficial populations at labeled rates (12). Spinetoram has a very favorable toxicity profile, with no evidence of teratogenicity, mutagenicity, carcinogenicity, or adverse reproductive effects. Compared to other registered chemicals spinetoram clearly demonstrates reduced risk to human health and non-target organisms (12). The rapid environmental degradation of spinetoram, combined with only requiring low rates for control of thrips, reduces the overall load of chemicals in the environment.

In numerous University of California studies conducted, spinetoram has demonstrated consistent performance in the field and provided excellent control of avocado thrips (11,13). Spinetoram provides reductions of thrips adults and immatures, resulting in protection from fruit scarring (damage) (11). When compared to other registered products for avocado thrips control in avocado, spinetoram offers one of the lowest PHIs (pre-harvest interval) and REIs (re-entry interval) (Table 1).

*Table 1. PHI and REI for insecticides recommended for management of avocado thrips, ranked in order from greatest to least IPM fit (3). Source of REI and PHI data is CDMS <http://www.cdms.net/Label-Database>*

Active Ingredient	PHI	REI
Abamectin	14	12
Spinetoram	1	4
Spirotetramet	1	24
fenpropathrin	1	24
Spinosad	1	4
Sabadilla	Do not apply at harvest	12

Spinetoram is a significant tool for IPM programs for control of avocado thrips in avocado, and has the following benefits:

- It is a selective insecticide with a low use rate.
- It has a favorable toxicological profile and is soft on beneficial arthropods.
- The unique mode of action of spinetoram makes it effective against avocado thrips as well as providing an alternate mode of action for managing against pesticide resistance.
- It is an ideal and widely used rotation partner, allowing up to 3 applications at the labeled rate per calendar year in avocado.

#### References:

1. Hoddle, M.S., S. Nakahara, and P.A. Phillips. 2002. Foreign exploration for *Scirtothrips perseae* Nakahara (Thysanoptera: Thripidae) and associated natural enemies on avocado (*Persea americana* Miller). *Biological Control* 24: 251-265.
2. Hoddle, M.S., J.G. Morse, P.A. Phillips, B.A. Faber, and K.M. Jetter. 2002. Avocado thrips: new challenge for growers. *California Agriculture*, May-June, pp. 104-107.
3. B.A. Faber, J.G. Morse, and M.S. Hoddle. 2015. Avocado thrips. <http://www.ipm.ucdavis.edu/PMG/r8300311.html>
4. Hoddle, M. 2013. The biology and management of the avocado thrips, *Scirtothrips perseae* Nakahara (Thysanoptera: Thripidae). <http://biocontrol.ucr.edu/hoddle/avocadonthrips.html>
5. Hoddle, M.S., K.M. Jetter, and J.G. Morse. 2003. The economic impact of *Scirtothrips perseae* Nakahara (Thysanoptera: Thripidae) on California avocado production. *Crop Production* 22: 485-493.
6. Oevering, P.E., B. Faber, and P. Phillips. 2004. Time of avocado pruning affects fruit set and avocado thrips occurrence in southern California groves. *California Avocado Society Yearbook*: 83-96. [http://www.avocadosource.com/cas\\_yearbooks/cas\\_87\\_2004-2005/cas\\_2004-05\\_v87\\_pg\\_083-096.pdf](http://www.avocadosource.com/cas_yearbooks/cas_87_2004-2005/cas_2004-05_v87_pg_083-096.pdf)
7. Hoddle, M.S., J.G. Morse, P. Oevering, P. Phillips, and B. Faber. 2002. Further progress on avocado thrips biology and management. *California Avocado Society*. [http://www.avocadosource.com/ARAC/Symposium\\_2002/a1.pdf](http://www.avocadosource.com/ARAC/Symposium_2002/a1.pdf)
8. *Scirtothrips perseae* (avocado thrips). Updated 20 January 2015. *Invasive Species Compendium*. <http://www.cabi.org/isc/datasheet/49072>
9. Hoddle, M.S. and J.G. Morse. 2003. Avocado thrips biology and control. *AvoResearch*, Spring 2003, 8 pp. [http://www.avocadosource.com/papers/research\\_articles/hoddlemark2003.pdf](http://www.avocadosource.com/papers/research_articles/hoddlemark2003.pdf)
10. Morse, J., A. Urena, and L. Robinson. 2009. Research update – avocado thrips, perseae mite, and pesticide resistance management. [http://www.avocadosource.com/arac/symposium\\_2008-2009/april\\_morse-pest.pdf](http://www.avocadosource.com/arac/symposium_2008-2009/april_morse-pest.pdf)
11. Bender, G., F. Byrne, A. Urena, and L. Robinson. 2013. 2013 Arthropod pest management update – focus on pesticide resistance management. *CAS-CAC-UCCE Seminar Series*. <http://slideplayer.com/slide/4401686/>
12. Dripps, J.E., R.E. Boucher, A. Chloridis, C.B. Cleveland, C.V. Deamicis, L.E. Gomez, D.L. Paroonagian, L.A. Pavan, T.C. Sparks, and G.B. Watson. Chapter 5, the spinosyn insecticides. *In*, "Green Trends in Insect Control" O. Lopez and J.G. Fernandez-Bolanos [eds], pp. 163-212.

13. Morse, J. 2008. Management and resistance monitoring of avocado thrips and perseia mite. California Avocado Commission 2008 Production Research Report.  
[http://www.avocadosource.com/ARAC/Symposium\\_2007-2008/MORSE-PESTS\\_EOY.pdf](http://www.avocadosource.com/ARAC/Symposium_2007-2008/MORSE-PESTS_EOY.pdf)



Dow AgroSciences

# Complimentary Copy

Dow AgroSciences LLC  
9330 Zionsville Road Indianapolis, IN 46268 USA

www.dowagro.com

December 1, 2015

Document Processing Desk (REGFEE)  
Office of Pesticide Programs  
U.S. EPA, Room S-4400  
One Potomac Yard (South Building)  
2777 South Crystal Drive  
Arlington, VA 22202-4501

**RE: PRIA 3 M007: Request to Extend Exclusive Use Data Period for Spinetoram (62719-539)**

Dear Mr. Stephen Schiabe:

Dow AgroSciences LLC (DAS) hereby requests EPA to extend the Exclusive Use Period for Spinetoram (62719-539) by three years for a total exclusive use of data period of September 28, 2007 until September 28, 2020. As set forth below, DAS respectfully submits that it has satisfied the requirements for extending the traditional ten year exclusive use of data period for new active ingredients by three years, in accordance with the minor use provisions of FIFRA § 3(c)(1)(F)(ii). Specifically, DAS has registered Spinetoram, a Presidential Green Chemistry Challenge Award winner and a Reduced Risk Pesticide, on more than nine minor use crops within seven years of its original registration and DAS has satisfied three conditions of FIFRA 3(c)(1)(F)(ii); the alternatives posed greater risks; spinetoram plays or will play a significant part in managing pest resistance; and spinetoram plays or will play a significant part in an integrated pest management program.

DAS representatives would be pleased to meet with the Agency to discuss the matter.

**Background:**

EPA first registered Spinetoram Technical (originally XDE-175; 62719-539), as well as two formulations (Delegate WG - 62719-541 and Radiant SC - 62719-545) on September 28, 2007. Spinetoram represents the next generation of spinosyn insecticides. Spinosyns are naturally derived fermentation products for arthropod pest control produced by the soil organism *Saccharopolyspora spinosa*, a novel bacterium of the order Actinomycetales. Spinetoram is a synthetically-modified spinosyn that is prepared from a mixture of two natural spinosyns produced by *S. spinosa* and possess the same insecticidal mode of action (MOA) as spinosad. Spinosad, a highly successful reduced risk insecticide, is the only other spinosyn insecticide commercially available today.

Spinetoram provides control of lepidopteran pests, leafminers and thrips in vegetables that is equal to or superior than spinosad, at use rates that are one-half (or less) of the rates used for spinosad. Spinetoram is more potent than spinosad, as it has increased activity at the (nicotinic acetylcholine receptor) target site. In addition, spinetoram is more stable in the sunlight and provides longer residual activity. Spinetoram provides control of Lepidoptera, thrips and other key pests of pome fruit, stone fruit, and tree nuts at relatively low use rates. Additionally, spinetoram controls other pests that spinosad does not, or is less active against, including codling moth, oriental fruit moth, navel orangeworm, plum curculio, and pear psylla. This increased level of control and residuality has allowed spinetoram to replace older products in markets, such as in pome and stone fruit, where spinosad use was limited.

Although spinosad and spinetoram are sometimes labeled on the same crops/uses, most growers prefer spinetoram due to the reduced rates required, and the additional control provided. Growers will either use spinosad or spinetoram, and normally do not use both on their crops. At this time, most spinosad use is in the organic markets, with conventional growers having converted to spinetoram products, with a few exceptions.

**DAS Spinetoram Minor Use Registrations:**

To date, DAS has registered more than 25 minor uses on spinetoram products. However, at this time, the focus of the following report will be on just 9 crops that currently have production data to demonstrate that the crops are below 300,000 acres, thus deemed "Minor" crops.

The current minor crops that are labeled for spinetoram and that will be discussed in the document below are:

Crop	2013 (acres harvested)	2012 (acres harvested)	Source
Blueberries	78960	77700	USDA
Raspberries	5600	5400	USDA
Strawberries	58190	55840	USDA
Cranberries	42000	40300	USDA
Tomato (fresh market) <sup>1</sup>	277,000	276,300	USDA
Tomato (for processing) <sup>1</sup>	99,600	101,000	USDA
Peppers	64,800	65,900	USDA
Green onion <sup>2</sup>	178,300	194,500	USDA
Bulb onion <sup>2</sup>	143,340	146,870	USDA
Head Lettuce	129,000	141,600	USDA

<sup>1</sup> Tomato (fresh/processing) are classified as separate minor use crops, but will be grouped together in document due to similarity in use pattern and pest pressure. We request they still be counted as 2 crops.

<sup>2</sup> Bulb onion and green onion are separate minor use crops, but will be grouped together in the document due to similarity in use pattern and pest pressure. We request they still be counted as 2 crops.

**DAS satisfies requirements II, III and IV (alternatives pose greater risk to environment/human health; managing pest resistance; and role in IPM) from FIFRA § 3(c)(1)(F)(ii) for the following 9 Crops:**

**Blueberry**

Spinetoram controls target pests through contact, ingestion and ovicidal activity in a wide variety of crops. One of the major pests is spotted wing drosophila (SWD), which is found in the minor use crop blueberry.

There are two types of blueberries grown in United States: Highbush blueberry (*Vaccinium corymbosum*) and lowbush blueberry (*V. angustifolium*, *V. myrtilloides*).

SWD (*Drosophila suzukii*) is an invasive species from Asia that has become a significant agricultural pest on blueberries in the United States. By 2014, the presence of SWD has become widespread throughout the western and eastern U.S., as well as in many European countries, Brazil, and Mexico (1). SWD are invasive vinegar flies (fruit flies) with multiple generations per year that can attack unripened fruit (2). Female SWD cut into intact fruit with their serrated ovipositor to lay eggs under the skin. This allows larvae of SWD to be present during ripening, leading to a risk of detection in ripe fruit after harvest. During egg-laying and larval feeding, sour rot and fungal diseases can also be introduced by SWD, further affecting fruit quality. There is a greater risk of fruit contamination at harvest from SWD compared with native species that lay eggs only in already-damaged and rotting fruit (3).

Several predatory insects feed on SWD adults and pupae, but not yet in sufficient quantities to provide significant control. A tiny predatory wasp that parasitizes SWD pupae is present in the Pacific Northwest and mid-Atlantic region and thus may be found in other regions as well. Thorough research is needed to fully understand whether and how this species may be utilized in long-term SWD management. As a result, blueberry growers cannot solely rely on natural enemies to manage SWD (4).

Successful spotted wing drosophila management programs should be focused on effective monitoring, cultural control and use of selective pesticides (5). An example of a cultural practice is to harvest all ripe and cull fruit from the planting and then dispose of the unwanted fruit to keep the entire planting completely clean. But this practice is very challenging and expensive, especially on larger acreages. In addition, SWD will multiply on wild fruit (raspberries or blackberries in hedgerows, mulberries, wild cherries, etc.) as well as cultivated fruit, and thus wild stands of these hosts can be reservoirs for SWD. Traps are typically used to detect adult SWD and determine whether control measures are needed, and are not intended to provide control (3). Extremely fine mesh (0.039 inches) screens can be used to protect plantings from SWD in high tunnels or greenhouses, but this practice reduces air flow and restricts access of pollinators during bloom (6). These types of cultural practices that are recommended by universities can be useful, however, most growers have found them to be insufficient to provide good control of annual and numerous populations of SWD.

Use of selective chemistry along with cultural practices for IPM program is important for managing SWD. Pesticides in three MOA groups – pyrethroids (IRAC, MOA Group 3), *spinosyns* (MOA Group 5), and organophosphates (MOA Group 1B) have demonstrated good efficacy against SWD adults. Neonicotinoids (MOA Group 4A) have very low efficacy and are not recommended by universities (7).



To prevent developing resistance in SWD, rotation of chemical families with different Modes of Action is essential for effective management of this pest (6, 8). Depending on environmental conditions, SWD populations can be very high with multiple generations per year, conditions which contribute to a high level of risk of resistance development. There is already documented resistance of SWD to pyrethrins in West Coast production (6).

Alternating insecticides with different MOAs is a very important part of IPM programs. To manage SWD during the season, a 7-14 day retreatment cycle is recommended, and the same product should not be used more than two consecutive times before rotating to a different class. Rotating among the various classes delays the onset of resistance to the pesticides that are used in the management program (22). Spinetoram is an ideal and widely used rotation partner, allowing up to 6 applications at the labeled rate for SWD per season.

In numerous university studies across the US, spinetoram demonstrated consistent performance in the field and provided excellent control of SWD (3, 7, 9, 10, 11, 12, 13, 14, 15, 16, 27). Spinetoram provides quick knockdown and residual control via contact and ingestion activity. It has a very favorable toxicity profile, with no evidence of teratogenicity, mutagenicity, carcinogenicity, or adverse reproductive effects. Compared to other registered chemicals such as acetamiprid, azinphos-methyl, lambda cyhalothrin, methomyl and phosmet; spinetoram clearly demonstrates reduced risk to human health and non-target organisms (17). The rapid environmental degradation of spinetoram, combined with low rates needed for control (Appendix 1) reduces the overall load of chemicals in the environment. Spinetoram, as a selective (18) reduced risk insecticide, plays a key role in adoption of IPM strategies as it has unique MOA and minimal impact on beneficial arthropod species such as and including big-eyed bugs (*Geocoris* sp.), damsel bugs (*Nabis* sp.), ladybird beetles (Coccinellidae) and lacewings (*Chrysopa* sp.) (Appendix 1).

Furthermore, there are products that are labeled in blueberries for SWD control such as Diazinon, Danitol, Mustang, Asana, Brigade and Lannate that are "restricted use pesticides (RUP)" due to acute toxicity to humans with possible oncogenicity and/or toxicity to fish, birds and aquatic organisms (19). Organophosphate insecticides within this list as well as Imidan, Diazinon and Malathion, have higher application rates and greater impacts on beneficial arthropods as compared to spinetoram. While pyrethroid insecticides are used at lower doses, they are broad spectrum and thus active on both beneficial and pest insects. Predator insects may be susceptible to lower dose than the pest, thus disrupting the predator-prey relationship, which leads to flaring up of mites and knock down of beneficial insects (18, 20, 21).

When compared to other registered products for SWD control in blueberries, spinetoram offers the shortest re-entry interval (REI), which is very important for U-pick farm operations, and also has a lower PHI at 7 days, compared to Asana (Appendix 1). Further work is being done by IR-4 to reduce the PHI on blueberries for SWD as it is a desired use pattern for the growers.

Spinetoram is a significant tool for IPM programs for control of SWD in blueberries, and has the following benefits:

- It is a selective insecticide with a low use rate
- It has a favorable toxicological profile and is soft on beneficial arthropods
- The unique mode of action of spinetoram makes it effective against SWD that become resistant to other insecticides as well as providing an alternate mode of action for managing against pesticide resistance.

## Raspberry

Spotted wing drosophila (SWD) is also a major pest in raspberry (*Rubus idaeus*), another minor use crop. SWD (*Drosophila suzukii*) is an invasive species from Asia that has become a significant agricultural pest on raspberries in the United States. By 2014, the presence of SWD has become widespread throughout the western and eastern U.S., as well as in many European countries, Brazil, and Mexico (1). SWD are invasive vinegar flies (fruit flies) with multiple generations per year that can attack unripened fruit (23, 24). Female SWD cut into intact fruit with their serrated ovipositor to lay eggs under the skin. This allows larvae of SWD to be present during ripening, leading to a risk of detection in ripe fruit after harvest. During egg-laying and larval feeding, sour rot and fungal diseases can also be introduced by SWD, further affecting fruit quality. There is a greater risk of fruit contamination at harvest from SWD compared with native species that lay eggs only in already-damaged and rotting fruit (24).

Several predatory insects feed on SWD adults and pupae, but not yet in sufficient quantities to provide significant control. A tiny predatory wasp that parasitizes SWD pupae is present in the Pacific Northwest and mid-Atlantic region and thus may be found in other regions as well. Thorough research is needed to fully understand whether and how this species may be utilized in long-term SWD management. As a result, caneberry growers cannot solely rely on natural enemies to manage SWD (4).

Successful spotted wing drosophila management programs should be focused on effective monitoring, cultural control and use of selective pesticides (5). An example of a cultural practice is to harvest all ripe and cull fruit from the planting and then dispose of the unwanted fruit to keep the entire planting completely clean. But, this practice is very challenging and expensive, especially on larger acreages. In addition, SWD will multiply on wild fruit (raspberries or blackberries in hedgerows, mulberries, wild cherries, etc.) as well as cultivated fruit, and thus wild stands of these hosts can be reservoirs of SWD. Traps are typically used to detect adult SWD and determine whether control measures are needed, and are not intended to provide control (25). Extremely fine mesh (0.039 inches) screens can be used to protect plantings from SWD in high tunnels or greenhouses, but this practice reduces air flow and restricts access of pollinators during bloom (6). Early harvest of fruit can be important in reducing exposure of fruit to the pest. These types of cultural practices recommended by universities can be useful, however, most growers have found them to be insufficient to provide good control of SWD populations.

Use of selective chemistry along with cultural practices for IPM program is important for managing SWD. Pesticides in three MOA groups – pyrethroids (IRAC, MOA Group 3), *spinosyns* (MOA Group 5), and organophosphates (MOA Group 1B) have demonstrated good efficacy against SWD adults. Neonicotinoids (MOA Group 4A) have very low efficacy and are not recommended by universities (7).

To prevent developing resistance in SWD, rotation of chemical families with different Mode of Action is essential for effective management of this notorious pest (6, 8, 26). Depending on environmental conditions, SWD populations can be very high with multiple generations per year, which most likely will lead to resistance development. There is already documented resistance of SWD to pyrethrins in the West Coast (6).

Alternating insecticides with different MOAs is a very important part of IPM programs. To manage SWD during the season, a 7-14 day retreatment cycle is recommend, and the same product should not be used more than two consecutive times before rotating to a different class. Rotating among the various classes delays the onset of resistance to the pesticides that are used in the management program (22). Spinetoram is an ideal and widely used rotation partner, allowing up to 6 applications at the labeled rate for SWD per season.

In numerous university studies across the US, spinetoram demonstrated consistent performance in the field and provided excellent control of SWD (3, 7, 9, 10,11,12,13,14, 15, 16). Spinetoram provides quick knockdown and residual control via contact and ingestion activity. It has a very favorable toxicity profile, with no evidence of teratogenicity, mutagenicity, carcinogenicity, or adverse reproductive effects. Compared to other registered chemicals such as acetamiprid, azinphos-methyl, lambda cyhalothrin, methomyl and phosmet, spinetoram clearly demonstrates reduced risk to human health and non-target organisms (17). The rapid environmental degradation of spinetoram, combined with only needing low rates for control (Appendix 1) will reduce the overall load of chemicals in the environment. Spinetoram, as a selective (18) reduced risk insecticide, plays a key role in adoption of IPM strategies. It has a unique MOA and induces little or no disruption of beneficial arthropod species such as and including big-eyed bugs (*Geocoris* sp.), damsel bugs (*Nabis* sp.), ladybird beetles (*Coccinellidae*) and lacewings (*Chrysopa* sp.) (Appendix 1).

Furthermore, there are products that are labeled in raspberries for SWD control such as Diazinon, Danitol, Mustang, Asana, Brigade and Lannate that are "restricted use pesticides (RUP)" due to acute toxicity to humans with possible oncogenicity and/or toxic to fish, birds and aquatic organisms (19). In addition, the pyrethroid class of chemistry is toxic to all insects, both beneficial and pests. Predator insects may be susceptible to a lower dose than the pest, disrupting the predator-prey relationship, which leads to flaring up of the mites and overall reduction in beneficial arthropod densities (18, 20, 21). Other registered products such as Imidan, Diazinon and Malathion are organophosphates, which compared to spinetoram have much higher application rates, and are extremely toxic to humans, beneficials and the environment (25).

Additionally, compared to other registered products for SWD control in raspberries, spinetoram offers the lowest REI, which is very important for U-pick farm operations, and low PHI compared to Asana that offers extended fruit protection in the storage. Growers need to spray up to harvest to help protect the fruit from SWD infestations.

Spinetoram is a significant tool for IPM programs for control of SWD in blueberries, and has the following benefits:

- It is a selective insecticide with a low use rate
- It has a favorable toxicological profile and is soft on beneficial arthropods
- The unique mode of action of spinetoram makes it effective against SWD that become resistant to other insecticides as well as providing an alternate mode of action for managing against pesticide resistance.

## Strawberry

A major pest in the minor use crop, strawberry (*Fragaria x ananassa*), is western flower thrips *Frankliniella occidentalis* (Perganda) (WFT). WFT are widely distributed throughout the United States, feed on a wide variety of plants and have multiple, overlapping generations per year on a broad range of plant species. WFT are slender, small insects that rasp the surface of the fruit and feed on strawberry blossoms, which causes the stigmas and anthers to turn brown and decline prematurely (30, 31). Typically, WFT adults insert bean-shaped eggs into leaf, flower, or fruit tissues and can survive for 4-5 weeks and oviposit about 50 eggs per one generation.

Pre-bloom feeding deforms blossoms and leaves, reducing fruit set or weakening the fruit so it is more susceptible to frost and temperature stresses. In addition, as fruit develops, thrips feeding may cause a russetting (30). Significant crop damage leading to severe yield losses can be caused by thrips that act as a vector of *Tospoviruses* (32). Thrips-transmitted tospoviruses (genus *Tospovirus*, family *Bunyaviridae*) are a major group of plant viruses affecting at least 1,090 host-plant species in 15 monocotyledonous and 69 dicotyledonous families worldwide (33, 34, 35, 36). Strawberry plantations often have a mixed population of thrips that includes a low percentage of onion thrips,

(*Thrips tabaci*), among other species that do not cause economic damage. By properly identifying WFT, unnecessary insecticide applications can be avoided. (38).

Natural predators such as adults and nymphs of the minute pirate bug, green lacewings and certain parasitic wasps are known to prey on plant feeding thrips. The naturally occurring minute pirate bugs, *Orius insidiosus*, are effective predators of thrips in fruiting vegetable crops and strawberries. Their effectiveness is predictable based on the number of the predator relative to the number of thrips prey (118). However, releasing beneficial insects and predatory mites against thrips in most cases will not provide sufficient control of this pest on their own (37). Spinosad and spinetoram are the most effective insecticides to suppress WFT, and they are reduced risk selective insecticides that do not suppress populations of *O. insidiosus* at labeled rates (40, 41, 42, 118). The use of selective insecticides that have minimal effects on natural enemy populations, is vital in the control of WFT and preservation of predators.

Successful WFT management programs should be focused on cultural controls, effective monitoring and identification of the thrips species and use of selective pesticides (37, 38). IPM encompasses the simultaneous management of multiple pests, regular monitoring of pests, their natural enemies and antagonists, use of economic or treatment thresholds when applying pesticides, and integrated use of multiple suppressive tactics. Integrated Pest Management practices that combine use of cultural and biological control as well the use of the most selective or least-toxic insecticides is essential for effective management of thrips. Choosing the right site and planting date for cultivars susceptible to WFT may help to decrease the crops susceptibility to thrips damage. Removal and disposal of spent flowers that can harbor thrips may also be helpful, although the general benefit of this practice in landscapes is unknown; and old blossoms also commonly shelter beneficial predators of thrips. Timely pruning of injured and infested terminals may help to promote the increase of predaceous mite populations. Another option is to increase the use of mulches that reflect light, which can interfere with certain flying insects' ability to locate plants and can delay or reduce the extent to which young plants become infested by thrips (44). These types of cultural practices that are recommended by universities can be useful, however, most growers have found them to be insufficient to provide good control of the numerous yearly populations of WFT on their own and that a selective insecticide is required.

Integrated resistance management (IRM) includes the rotation of insecticides from different chemical classes, the use of recommended rates, the limitation of maximum number of applications and product per acre per year or season, and the avoidance of sequential treatments within a single planting and across sequential crops/plantings.

Pesticides in four activity groups – pyrethrins (IRAC, MOA Group 3), spinosyns (MOA Group 5), neonicotinoids (MOA Group 4A), sulfoximines (Group4C), and organophosphates (MOA Group 1B) have varied activity against WFT ranging from suppression to control (43). Insecticides continue to have an important role to play in WFT management, although the use of insecticides must be judicious. Decisions regarding which insecticides are to be used and the timing should be made in the context of both short term and long-term management goals. Minimizing resistance development and avoiding the flaring of WFT populations by their release from natural enemies are critical factors in insecticide use decisions (44). Behavior and ecology of WFT can minimize exposure to insecticides, due to the species being well suited to evolve resistance to multiple classes of insecticides. There have been numerous incidences of resistance reported to all major classes of insecticides from all regions of the world (102, 103, 104, 105, 106, 107, 108). In addition to pest management failures and resistance development, there is a limited pool of efficacious insecticides for use against WFT (47), which further increases the selection pressure on the remaining chemistries.

The key to managing resistance is to reduce selection pressure by rotating between insecticides with different modes of action and reducing the number of insecticide applications. It may be necessary to use nonchemical control methods and rotate to insecticides that may not provide the highest level of control.

The negative impacts of broad-spectrum insecticides, such as pyrethroids, on natural enemies and competitor species, in addition to continuous, overlapping generations typical for WFT populations in crops makes it especially challenging to implement IRM strategy (45).

The most efficacious insecticides for WFT, at present, are in the spinosyn class. No other insecticide class provides a similar level of effectiveness against WFT (44, 46). Spinosad insecticide products were first introduced in the late 1990s and have been highly effective in controlling WFT and widely used for this purpose (39). Spinetoram, a new and more active insecticide, was registered for use in 2008. Spinetoram represents a unique mode of action (Group 5 insecticides) and is one of the most effective insecticides to suppress WFT because it provides excellent control and it is a reduced risk insecticide that does not suppress beneficial populations, including minute pirate bug (*O. insidiosus*), at labeled rates (40, 41, 42). Rotating highly active insecticides such as spinetoram with moderately active insecticides could achieve good levels of control of the WFT while mitigating the development of resistance to all insecticides in the rotation. Other insecticides registered for suppression of WFT may not be as efficacious as spinosyns, however, they should be included as part of IPM program (44).

As mentioned, alternating insecticides with different MOAs is a very important part of IPM programs (44, 50). To manage WFT during the season, a 3-7 day retreatment cycle is recommended and the same product should not be used more than two consecutive times before rotating to a different class. Rotating among the various classes delays the onset of resistance to the pesticides that are used in the management program (44).

In numerous university studies across the US, spinetoram demonstrated consistent performance in the field and provided excellent control of WFT (40, 41, 42, 44, 46, 50). Spinetoram provides quick knockdown and residual control via contact and ingestion activity. It has a very favorable toxicity profile, with no evidence of teratogenicity, mutagenicity, carcinogenicity, or adverse reproductive effects. Compared to other registered chemicals such as acetamiprid, naled, malathion, pyrethrins + piperonyl butoxide and clothianidin, spinetoram clearly demonstrates reduced risk to human health and non-target organisms (17). The rapid environmental degradation of spinetoram, combined with only requiring low rates for control of WFT, reduces the overall load of chemicals in the environment.

For the Southeastern United States, there are only two products registered for WFT, spinosad and spinetoram (48). For California and Florida, there are a number of registered insecticides available with variable efficacy. Dibrom and Malathion are organophosphates that have much higher application rates than Spinetoram, and are more toxic to humans, beneficial arthropods and the environment (25, 31). In addition, Dibrom and Malathion provide only 50% control or less of WFT and can induce spider mite outbreaks (37). If registered in the state, neonicotinoids such as Belay and Assail can be a good rotation partner for spinetoram as a different class of chemistry and MOA. However, neonicotinoid insecticides have reported effects on predaceous arthropods and can have variable activity on WFT (37, 46, 49). Horticultural oil, natural pyrethrins (plus piperonyl butoxide), or insecticidal soap have very low efficacy and require multiple applications.

When compared to other registered products for WFT control in strawberries, spinetoram offers one of the lowest REI, which is very important for U-pick farm operations.

Spinetoram is a significant tool for IPM programs for control of WFT in strawberries, and has the following benefits:

- It is a selective insecticide with a low use rate.
- It has a favorable toxicological profile and is soft on beneficial arthropods.
- The unique mode of action of spinetoram makes it effective against WFT as well as providing an alternate mode of action for managing against pesticide resistance.
- It is an ideal and widely used rotation partner, allowing up to 3 applications at the labeled rate per calendar year in strawberries.

### Cranberry

One of the major pests in the minor use crop, cranberry (*Oxycoccus microcarpus*), is the blackheaded fireworm (BHF<sub>W</sub>) (*Phopobota naevana*). Proper identification plays a major role in achieving control of this pest. Larvae are approximately 7-9 mm in length and have a distinct shiny black head. The body is a greenish, greenish-yellowish, or grayish color. BHF<sub>W</sub> is one of the most significant cranberry pests because of the damage it is able to inflict on cranberry (88). BHF<sub>W</sub> are widely distributed throughout the United States and have 2-3 generations per year. Females of both generations lay about 70-80 eggs (89). BHF<sub>W</sub> overwinters in the egg stage and the hatching period in the spring can last up to 6 weeks. Newly hatched larvae burrow into cranberry leaves or mine into unopened terminal buds as they swell. On new growth, the larva uses silk to gather 2 or more leaves in the tip area and then feeds within. Older caterpillars use webbing to construct up to 5-6 tents during the course of development (90). The primary damage caused by BHF<sub>W</sub> on cranberries is larval feeding on foliage and fruit. The foliage turns brown over time as a result of larval feeding. BHF<sub>W</sub> larva feed on the surface of the fruit as well. Second generation of BHF<sub>W</sub> usually reduce the crop for the following year because the feeding on the tips of the upright results in a failure to form normal fruit buds.

Minimizing damage to the plants and crop by insect pests is one of the most important challenges in cranberry production. Failure to manage pest insects properly can result in severe crop loss, vine damage, or in extreme cases, the death of large areas of the bog.

Predators and parasitoids which coexist in the bog environment play an important role in regulating cranberry pest populations. Natural predators such as adults and nymphs of the ladybird beetles, green lacewings, spined soldier bug, praying mantis and certain parasitic insects are known to prey on plant feeding BHF<sub>W</sub> (93). However, maintaining naturally occurring populations of these beneficials may be difficult. If there is low presence of insects for ladybeetles to feed on, they will disperse to more productive hunting grounds. Although the green lacewing adults also feed on insects, they require other food sources such as many soft insects and mites, and if any necessary type of food is not available, they will fly elsewhere. Spined soldier bug also feed on a variety of slow-moving, soft-bodied insects that live on plants. However, no research has been done to fully evaluate the full benefit of this predator in cranberries (92). As of today, limited research-based information is available on effectiveness of beneficials. Releasing beneficial insects to manage BHF<sub>W</sub> in most cases will not provide sufficient control of this notorious pest and can be very costly (92). Furthermore, bio-control is often too late, as the crop damage occurs before the predators and parasites can build up to adequate numbers.

Successful BHF<sub>W</sub> management programs should be focused on effective monitoring, cultural control and use of selective pesticides (91, 93). IPM encompasses the simultaneous management of multiple pests, regular monitoring of pest and their natural enemies and antagonists, use of economic or treatment thresholds when applying pesticides, and integrated use of multiple suppressive tactics. Integrated Pest Management practices that combine use of cultural and biological control as well the use of the most selective or least-toxic insecticides is essential for effective management of BHF<sub>W</sub>. Use of IPM traps for monitoring the flight pattern and peak catch is an important part of monitoring for BHF<sub>W</sub>. Scouting is difficult, particularly for small larvae. However, this stage is very critical for successful management of this pest due to increased sensitivity to chemical treatments (88). Ecological management such as sanding and flooding may help to reduce the number of larvae, but will not eliminate all larvae.

Providing appropriate cultural care to keep plants vigorous may help to increase their tolerance to BHF<sub>W</sub> damage. Use of more tolerant cranberry varieties may be helpful as well. Although -these types of cultural practices recommended by universities can be useful, most growers have found them to be insufficient to provide good control of BHF<sub>W</sub> (94). Spraying chemical pesticides is the most common practice to manage herbivorous insects in cranberries.

Insecticide resistance management (IRM) includes the rotation of insecticides from different chemical classes, the use of recommended rates, the limitation of maximum number of applications of products per acre per year or season, and the avoidance of sequential treatments within a single planting and across sequential crops/plantings. The use of selective insecticides that have minimal effects on natural enemy populations is important in the control of BHF<sub>W</sub>. Pesticides in three major activity groups - *spinosyns* (MOA Group 5), neonicotinoids (MOA Group 4A), sulfoximines (Group 4C) and organophosphates (MOA Group 1B) have varied activity against BHF<sub>W</sub> ranging from suppression to control (93, 95, 97). However, the use of insecticides must be judicious. Decisions regarding the proper insecticide to use and timing need to be made in the context of both short term and long-term management goals. Critical factors in insecticide use decisions for BHF<sub>W</sub> include minimizing resistance development and minimizing effect on natural enemies (94). Current control methods for the management of BHF<sub>W</sub> include the repeated use of insecticides which can lead to resistance (96).

Alternating insecticides with different MOAs is a very important part of IPM programs (44, 50). To manage BHF<sub>W</sub> during the season, it is recommended to use a 7-14 day retreatment cycle and the same product should not be used more than two consecutive times before rotating to a different class. Rotating among the various classes delays the onset of resistance to the pesticides that are used in the management program (44).

The most efficacious insecticides for BHF<sub>W</sub>, at present, are in the spinosyn class. Spinetoram is one of the most efficacious synthetic insecticides against BHF<sub>W</sub>. Spinosad insecticide products were first introduced in the late 1990s and have been highly effective in controlling BHF<sub>W</sub> and widely used for this purpose (39). Spinetoram, a new and more active insecticide, was registered for use in 2008. Spinosyn insecticides represent a unique mode of action (Group 5 insecticides). Spinosad and spinetoram are the most effective insecticides to suppress *Phopobota naevana*, and they are reduced risk insecticides that do not suppress populations of predatory insects at labeled rates (40, 41, 42).

Rotating highly active insecticides such as spinetoram with moderately active insecticides could achieve good levels of control of the BHF<sub>W</sub> while mitigating the development of resistance to all insecticides in the rotation. Other insecticides registered for suppression of BHF<sub>W</sub> may not be as efficacious as spinosyns, however, they should be included as part of IPM program (97).

In numerous university studies conducted across the US, spinetoram demonstrated consistent performance in the field and provided excellent control of BHF<sub>W</sub> (98, 99, 100, 101). Spinetoram provides quick knockdown and residual control via contact and ingestion activity. It has a very favorable toxicity profile, with no evidence of teratogenicity, mutagenicity, carcinogenicity, or adverse reproductive effects. Compared to other registered chemicals such as acetamiprid, diazinon, clothianidin, carbaryl and phosmet, spinetoram clearly demonstrates reduced risk to human health and non-target organisms (17, 97). The rapid environmental degradation of spinetoram, combined with low application rates, reduces the overall load of chemicals in the environment when managing BHF<sub>W</sub>. Spinetoram, as a selective (18) reduced risk insecticide, plays a key role in adoption of IPM strategies as it has unique MOA and little or no disruption of beneficial arthropod species such as and including big-eyed bugs (*Geocoris* sp.), damsel bugs (*Nabis* sp.), ladybird beetles (*Coccinellidae*) and lacewings (*Chrysopa* sp.).

There is list of registered insecticides available to manage BHF<sub>W</sub> on cranberries with variable efficacy. Diazinon, phosmet and acephate are organophosphate insecticides that have much higher application rates when compared to spinetoram, and are more toxic to humans, beneficial arthropods and the environment (25, 31). Also, organophosphate and carbamate insecticides are considered "high risk" by FQPA and may be unavailable for use in the near future (66).

Furthermore, Diazinon and Actara that are labeled in cranberries for BHF<sub>W</sub> control are "restricted use pesticides (RUP)" due to acute toxicity to humans with possible oncogenicity and/or toxic to fish, birds and aquatic organisms (19). Neonicotinoids such as Belay, Assail, Scorpion and *Bt* products can be a good rotation partner for spinetoram due to its different class of chemistry. However, these classes of chemistry may not effectively reduce overall infestations of BHF<sub>W</sub>.

When compared to other registered products for BHF<sub>W</sub> control in cranberries, spinetoram offers the lowest REI (4 hours) compared to Assail, Avaunt, Belay, Diazinon, Imidan, Orthene and Sevin and the lowest PHI (1day) compared to Avaunt and Orthene (97).

Spinetoram is a significant tool for IPM programs for control of BHF<sub>W</sub> in cranberries, and has the following benefits:

- It is a selective insecticide with a low use rate.
- It has a favorable toxicological profile and is soft on beneficial arthropods.
- The unique mode of action of spinetoram makes it effective against BHF<sub>W</sub> as well as providing an alternate mode of action for managing against pesticide resistance.
- It is an ideal and widely used rotation partner, allowing up to 6 applications at the labeled rate per calendar year in cranberries.

#### **Tomatoes (fresh market and processing)**

One of the major pests in the minor use crop tomatoes (*Solanum lycopersicum* L.), is western flower thrips, *Frankliniella occidentalis* (Perganda), (WFT). WFT is one of the most significant agricultural pests globally because of the damage it is able to inflict on a wide range of crops (47). WFT are widely distributed throughout the United States and have multiple, overlapping generations per year on a broad range of plant species. Adults and larvae feed by piercing plant tissues with their needle-shaped mandible and draining the contents of punctured cells (47, 54). Feeding by adults and larvae produces scarring on foliage, flowers and fruits, which results in aesthetic crop damage and disrupts plant growth and physiology. Also, oviposition can produce a wound response in fruiting structures, which reduces the marketability of certain horticultural produce (47, 55). Typically, WFT adults insert



bean-shaped eggs into leaf, flower, or fruit tissues and can survive for 4-5 weeks and oviposit about 50 eggs per one generation.

The primary damage caused by WFT to tomatoes is the vectoring of tomato spotted wilt virus, genus *Tospovirus*. The virus can only be acquired by the immature stage of thrips, whereas plant-to-plant transmission primarily occurs by adults. The adult thrips can transmit the virus for the remainder of their lives, which can last 30 to 45 days.

Tomato plantations often have a mixed population of thrips that includes a low percentage of the onion thrips, *Thrips tabaci*. Thrips feeding on plants can damage fruit, leaves, and shoots and very noticeably affect plants' cosmetic appearance. In addition, as fruit develops, thrips feeding may cause a russetting (30). Significant crop damage leading to severe yield losses can be caused by thrips that serve as vectors of *Tospoviruses* (32). Thrips-transmitted tospoviruses (genus *Tospovirus*, family *Bunyaviridae*) are a major group of plant viruses affecting at least 1,090 host-plant species in 15 monocotyledonous and 69 dicotyledonous families worldwide (33, 34, 35, 36). The actual amounts of economic losses attributable to tomato spotted wilt virus alone caused over US\$1 billion in losses annually on a global basis. This estimate did not include the direct damage also caused by WFT (56, 57).

Natural predators such as adults and nymphs of the minute pirate bug, green lacewings and certain parasitic wasps are known to prey on plant feeding thrips. The naturally occurring minute pirate bugs, *Orius insidiosus* (Say), are effective predators of thrips in fruiting vegetable crops. Their effectiveness is predictable based on the number of the predator relative to the number of thrips prey (118). Spinosad and spinetoram are the most effective insecticides to suppress WFT, and they are reduced risk selective insecticides that do not suppress populations of *O. insidiosus* at labeled rates, making the spinosyns extremely useful for thrips control (40, 41, 42, 118). Unfortunately, minute pirate bug species, which preferentially preys on WFT, do not have an affinity for tomato (44, 60, 61). Releasing beneficial insects and predatory mites against thrips in most cases will not provide sufficient control (37). Furthermore, biocontrol is often too late, as the crop damage is already present before the predators and parasites can build up to adequate numbers (59).

Successful WFT management programs should be focused on effective monitoring, cultural control and use of selective pesticides (37). IPM encompasses the simultaneous management of multiple pests, regular monitoring of pests and their natural enemies and antagonists, use of economic or treatment thresholds when applying pesticides, and integrated use of multiple suppressive tactics. Integrated Pest Management practices that combine use of cultural and biological control as well the use of the most selective or least-toxic insecticides is essential for effective management of thrips. Choosing the right site for planting WFT susceptible plants and providing appropriate cultural care to keep plants vigorous may help to increase their tolerance to thrips damage. Removal and disposal of spent flowers that can harbor thrips may also be helpful. However, the general benefit of this practice in landscapes is unknown; and old blossoms also commonly shelter beneficial predators of thrips. Timely pruning of injured and infested terminals may help to promote increases in predaceous mite populations (37). Additional cultural practices include increasing the use of mulches that reflect light, which can interfere with certain flying insects' ability to locate plants and can delay or reduce the extent to which young plants become infested by thrips (44). These types of cultural practices that are recommended by universities can be useful; however, most growers have found them to be insufficient to provide good control of the numerous yearly populations of WFT by themselves.

Integrated resistance management (IRM) includes the rotation of insecticides from different chemical classes, the use of recommended rates, the limitation of maximum number of applications and product per acre per year or season, and the avoidance of sequential treatments within a single planting and across sequential crops/plantings.

Pesticides in four activity groups – pyrethrins (IRAC, MOA Group 3), *spinosyns* (MOA Group 5), neonicotinoids (MOA Group 4A), sulfoximines (Group 4C), and organophosphates (MOA Group 1B) have varied activity against WFT ranging from suppression to control (43). Insecticides continue to have an important role to play in WFT management, although the use of insecticides must be judicious. Decisions regarding which insecticides are to be used and the timing should be made in the context of both short term and long-term management goals. Minimizing resistance development and avoiding the flaring of WFT populations by their release from natural enemies is a critical factor in insecticide use decisions (44). Behavior and ecology of WFT can minimize exposure to insecticides, due to the species being well suited to evolve resistance to multiple classes of insecticides. There have been numerous incidences of resistance reported to all major classes of insecticides from all regions of the world (47, 62, 63, 64, 65). In addition to pest management failures and resistance development, there is a limited pool of efficacious insecticides for use against WFT (47), which further increases the selection pressure on the remaining chemistries.

The key to managing resistance is to reduce selection pressure by rotating between insecticides with different modes of action and reducing the number of insecticide applications. It may be necessary to use nonchemical control methods and rotate to insecticides that may not provide the highest level of control.

The negative impacts of broad-spectrum insecticides, such as pyrethroids, on natural enemies and competitor species, in addition to continuous, overlapping generations typical for WFT populations in crops makes it especially challenging to implement IRM strategies (45).

As mentioned, alternating insecticides with different MOAs is a very important part of IPM programs (44, 50). To manage WFT during the season, a 3-7 day retreatment cycle is recommended, and the same product should not be used more than two consecutive times before rotating to a different class. Rotating among the various classes delays the onset of resistance to the pesticides that are used in the management program (44).

The most efficacious insecticides for WFT, at present, are in the spinosyn class. No other insecticide class provides a similar level of effectiveness against WFT (44, 46). Spinosad insecticide products were first introduced in the late 1990s and have been highly effective in controlling WFT and widely used for this purpose (39). Spinetoram, a new and more active insecticide, was registered for use in 2008. Spinetoram represents a unique mode of action (Group 5 insecticides) and is the most effective insecticide against WFT, as it provides excellent control and is a reduced risk insecticide that does not suppress populations of natural predators, including minute pirate bugs, at labeled rates (40, 41, 42).

Rotating highly active insecticides such as spinetoram with moderately active insecticides could achieve good levels of control of the WFT while mitigating the development of resistance to all insecticides in the rotation. Other insecticides registered for suppression of WFT may not be as efficacious as spinosyns, however, they should be included as part of IPM program (44).

In numerous university studies across the US, spinetoram demonstrated consistent performance in the field and provided excellent control of WFT (40, 41, 42, 44, 46, 50). Spinetoram provides quick knockdown and residual control via contact and ingestion activity. It has a very favorable toxicity profile, with no evidence of teratogenicity, mutagenicity, carcinogenicity, or adverse reproductive effects.

Compared to other registered chemicals such as acetamiprid, bifenthrin, beta-cyfluthrin, clothianidin, fenpropathrin, lambda cyhalothrin, zeta-cypermethrin, dinotefuran, methomyl and dimethoate, spinetoram clearly demonstrates reduced risk to human health and non-target organisms (17). The rapid environmental degradation of spinetoram, combined with only needing low rates for control of WFT will reduce the overall load of chemicals in the environment.

There is list of registered insecticides available to manage WFT on tomatoes with efficacy that varies widely. Dimethoate is an organophosphate, which compared to spinetoram has much higher application rates, and higher levels of toxicity to humans, beneficials and the environment (25, 31). Also, organophosphate and carbamate insecticides are considered "high risk" by FQPA and may be unavailable for use in the near future (66). In addition, dimethoate has low efficacy on WFT and can cause spider mite outbreaks (67). Furthermore, there are large number of pyrethroids that are labeled in tomato for WFT control such as Danitol, Mustang, Hero, Asana, Brigade, Karate, Leverage, Proaxis, Warrior II, Athena, Baythroid and Lannate (carbamate) that are "restricted use pesticides (RUP)" due to acute toxicity to humans with possible oncogenicity and/or toxic to fish, birds and aquatic organisms (19). In addition, the pyrethroid class of chemistry is not highly selective and is toxic to both beneficials and pests. Predator insects may be susceptible to a lower dose than the pest, disrupting the predator-prey relationship, which leads to flaring up of the mites and reduction in beneficial insect populations (18, 20, 21).

If registered in the state, neonicotinoids such as Belay and Assail can be a good rotation partners for spinetoram because they are from a different class of chemistry with a different MOA. However, neonicotinoid insecticides have reported effects on predaceous arthropods and can have variable activity on WFT (37, 46, 49). Horticultural oil, natural pyrethrins, or insecticidal soap have very low efficacy and require multiple applications.

Additionally, compared to other registered products for WFT control in tomatoes, spinetoram offers one of the lowest REI, which is very important for fresh market farm operations and the lowest PHI (1 day) compared to Asana, Assail, Platinum, Athena, Gladiator, Karate, Proaxis, Belay, Danitol, Sevin, Warrior II and Durivo (68).

Spinetoram is a significant tool for IPM programs for control of WFT in tomatoes, and has the following benefits:

- It is a selective insecticide with a low use rate.
- It has a favorable toxicological profile and is soft on beneficial arthropods.
- The unique mode of action of spinetoram makes it effective against WFT as well as providing an alternate mode of action for managing against pesticide resistance.
- It is an ideal and widely used rotation partner, allowing up to 6 applications at the labeled rate per calendar year in tomatoes.

## Peppers

One of the major pests in the minor use crop peppers (*Capsicum annuum L*) is western flower thrips, *Frankliniella occidentalis* (Perganda), (WFT). WFT are widely distributed throughout the United States, feed on a wide variety of plants and have multiple, overlapping generations per year on a broad range of plant species. Thrips feeding on plants can damage fruit, leaves, and shoots and very noticeably affect plants' cosmetic appearance. Adults and larvae feed by piercing plant tissues with their needle-shaped mandible and draining the contents of punctured cells (47, 54). Feeding by adults and larvae produces "flecking" on fruits, which results in aesthetic crop damage and disrupts plant growth and physiology. Peppers display a range of symptoms, and the ring spots are not always present on the leaves. Typically, WFT adults insert bean-shaped eggs into leaf, flower, or fruit tissues and can survive for 4-5 weeks and oviposit about 50 eggs per one generation.

The primary damage caused by WFT to peppers is the vectoring of tomato spotted wilt virus, groundnut ringspot virus and tomato chlorotic spot virus (71). The virus can only be acquired by the immature stage of thrips, whereas plant-to-plant transmission primarily occurs by adults. The adult thrips can transmit the virus for the remainder of their lives, which can last 30 to 45 days. Significant crop damage leading to severe yield losses can be caused by

thrips that act as a vector of *Tospoviruses* (32). Thrips-transmitted tospoviruses (genus *Tospovirus*, family *Bunyaviridae*) are a major group of plant viruses affecting at least 1,090 host-plant species in 15 monocotyledonous and 69 dicotyledonous families worldwide (33, 34, 35, 36).

Across the US, pepper plantations often have a mixed population of thrips that can include a percentage of one or more of the following: *Thrips tabaci*, *F. Tritici*, *F. bispinosa* and *F. fusca*. Large populations of *F. occidentalis*, *F. tritici*, and *F. bispinosa* migrate into the spring crop of flowering peppers, causing reduced-marketability from thrips feeding damage and from their vectoring of tomato spotted wilt virus (69, 70).

Natural predators such as adults and nymphs of the minute pirate bug, green lacewings and certain parasitic wasps are known to prey on plant feeding thrips. The naturally occurring minute pirate bugs, *Orius insidiosus*, are very effective predators of thrips in peppers, as pepper is a better reproductive host for minute pirate bugs than tomato. Adults of minute pirate bugs are highly mobile, and can rapidly invade pepper and eggplant fields to control the WFT adults and larvae (42, 45, 118). Their effectiveness is predictable based on the number of the predator relative to the number of thrips prey (118). Spinosad and spinetoram are the most effective insecticides to control WFT, and they are also reduced risk selective insecticides that do not suppress populations of minute pirate bugs at labeled rates (40, 41, 42, 45, 118). Conversely, pyrethroids are known to suppress populations of minute pirate bugs, while increasing populations of the WFT in pepper (45). The use of selective insecticides that have minimal effects on natural enemy populations is vital in the control of WFT and preservation of predators.

Releasing beneficial insects and predatory mites against thrips in most cases will not provide sufficient control of this key pest (37). Furthermore, bio-control is often too late, as the crop damage occurs before the predators and parasites can build up to adequate numbers (59). In northern Florida, minute pirate bugs are not active during the winter and early spring. There is usually a lag time in spring pepper and eggplant during which populations of thrips increase before natural populations of the minute pirate bugs invade in sufficient numbers to suppress the pests (40, 41).

Successful WFT management programs should be focused on effective monitoring, cultural control and use of selective pesticides (37). IPM encompasses the simultaneous management of multiple pests, regular monitoring of pests and their natural enemies and antagonists, use of economic or treatment thresholds when applying pesticides, and integrated use of multiple suppressive tactics. Integrated Pest Management practices that combine use of cultural and biological control as well the use of the most selective or least-toxic insecticides is essential for effective management of thrips. Choosing the right site for planting peppers and providing appropriate cultural care to keep plants vigorous may help to increase their tolerance to thrips damage. Removal and disposal of spent flowers that can harbor thrips may also be helpful. However, the general benefit of this practice in fields is unknown; and old blossoms also commonly shelter beneficial predators of thrips. Timely pruning of injured and infested terminals may help to promote increases in predaceous mite populations (37). An additional cultural practice includes increasing the use of mulches that reflect light, which can interfere with certain flying insects' ability to locate plants and can delay or reduce the extent to which young plants become infested by thrips (44). These types of cultural practices that are recommended by universities can be useful, however most growers have found them to be insufficient to provide good control of the numerous yearly populations of WFT on their own.

Integrated resistance management (IRM) includes the rotation of insecticides from different chemical classes, the use of recommended rates, the limitation of maximum number of applications and product per acre per year or season, and the avoidance of sequential treatments within a single planting and across sequential crops/plantings. Pesticides in four activity groups – pyrethrins (IRAC, MOA Group 3), *spinosyns* (MOA Group 5), neonicotinoids (MOA Group 4A), sulfoximines (Group 4C) and organophosphates (MOA Group 1B) have varied activity against WFT ranging from suppression to control (43). Insecticides continue to have an important role to play in WFT

management, although the use of insecticides must be judicious. Decisions regarding which insecticides are to be used and the timing should be made in the context of both short term and long-term management goals. Minimizing resistance development and avoiding the flaring of WFT populations by their release from natural enemies are critical factors in insecticide use decisions (44). Behavior and ecology of WFT can minimize exposure to insecticides and increase risk of resistance due to the species being well suited to evolve resistance to multiple classes of insecticides. There have been numerous incidences of resistance reported in WFT to all major classes of insecticides from all regions of the world (47, 62, 63, 64, 65). In addition to pest management failures and resistance development, there is a limited pool of efficacious insecticides for use against WFT (47), which further increases the selection pressure on the remaining chemistries.

The key to managing resistance is to reduce selection pressure by rotating between insecticides with different modes of action and reducing the number of insecticide applications. It may be necessary to use nonchemical control methods and rotate to insecticides that may not provide the highest level of control.

The negative impacts of broad-spectrum insecticides, such as pyrethroids, on natural enemies and competitor species, in addition to continuous overlapping generations typical for WFT populations in crops makes it especially challenging to implement IRM strategies (45).

As mentioned, alternating insecticides with different MOAs is a very important part of IPM programs (44, 50). To manage WFT during the season, a 3-7 day retreatment cycle is recommended, and the same product should not be used more than two consecutive times before rotating to a different class. Rotating among the various classes delays the onset of resistance to the pesticides that are used in the management program (44).

The most efficacious insecticides for WFT, at present, are in the spinosyn class. No other insecticide class provides a similar level of effectiveness against WFT (44, 46). Spinosad insecticide products were first introduced in the late 1990s and have been highly effective in controlling WFT and widely used for this purpose (39). Spinetoram, a new and more active insecticide, was registered for use in 2008. Spinetoram represents a unique mode of action (Group 5 insecticides), and is one of the most effective insecticides to control WFT, as it provides excellent control and is a reduced risk insecticide that does not suppress beneficial populations, including minute pirate bug, at labeled rates (40, 41, 42).

Rotating highly active insecticides such as spinetoram with moderately active insecticides could achieve good levels of control of the WFT while mitigating the development of resistance to all insecticides in the rotation. Other insecticides registered for suppression of WFT may not be as efficacious as spinosyns, however, they should be included as part of IPM program (44).

In numerous university studies conducted across the US, spinetoram demonstrated consistent performance in the field and provided excellent control of WFT (40, 41, 42, 44, 46, 50). Spinetoram provides quick knockdown and residual control via contact and ingestion activity. It has a very favorable toxicity profile, with no evidence of teratogenicity, mutagenicity, carcinogenicity, or adverse reproductive effects.

Compared to other registered chemicals such as dimethoate, acetamiprid, azinphos-methyl, lambda cyhalothrin, methomyl and phosmet, spinetoram clearly demonstrates reduced risk to human health and non-target organisms (17). The rapid environmental degradation of spinetoram, combined with only needing low rates for control of WFT will reduce the overall load of chemicals in the environment.

There is list of registered insecticides available to manage WFT on peppers with efficacy that varies widely. Dimethoate is an organophosphate that has much higher application rates, and higher levels of toxicity to humans, beneficials and the environment when compared to Spinetoram (25, 31). Additionally, dimethoate has low efficacy on WFT and can cause spider mite outbreaks (67). In general, organophosphate and carbamate insecticides are considered "high risk" by FQPA and may be unavailable for use in the near future (66).

Furthermore, there are a large number of pyrethroids labeled in peppers for WFT control such as Danitol, Endigo, Hero, Asana, Brigade, Proclaim, Warrior II, Athena, Baythroid and Lannate (carbamate) that are "restricted use pesticides (RUP)" due to acute toxicity to humans with possible oncogenicity and/or toxicity to fish, birds and aquatic organisms (19). In addition, the pyrethroid class of chemistry is toxic to all insects, both beneficial and pests. Predator insects may be susceptible to a lower dose than the pest, disrupting the predator-prey relationship, which leads to flaring up of mites and knocking down of beneficial insects (18, 20, 21).

If registered in the state, neonicotinoids such as Belay and Assail can be a good rotation partner for spinetoram because they are in a different class of chemistry with a different MOA. However, neonicotinoid insecticides have reported effects on predaceous arthropods and can have variable activity on WFT (37, 46, 49). Horticultural oil, natural pyrethrins, or insecticidal soap have very low efficacy and require multiple applications.

When compared to other registered products for WFT control in peppers, spinetoram offers one of the lowest REI, which is very important for fresh market farm operations and the lowest PHI (1 day) compared to Asana, Assail, Plutonium, Athena, Gladiator, Belay, Danitol, Sevin, Warrior II and Durivo (68).

Spinetoram is a significant tool for IPM programs for control of WFT in peppers, and has the following benefits:

- It is a selective insecticide with a low use rate.
- It has a favorable toxicological profile and is soft on beneficial arthropods.
- The unique mode of action of spinetoram makes it effective against WFT as well as providing an alternate mode of action for managing against pesticide resistance.
- It is an ideal and widely used rotation partner, allowing up to 6 applications at the labeled rate per calendar year in peppers.

#### **Onion (green and bulb)**

A major pest in the minor use crop onion, *Allium cepa* L, is onion thrips, *Thrips tabaci* Lindeman (Thysanoptera: Thripidae) (OT). Onion thrips is the key insect pest in most onion production regions of the world and can reduce yield by 30-50%. Losses can be even more severe if thrips infest the crop with Iris yellow spot virus (72, 73). OT causes both direct and indirect damage to onion by feeding and ovipositing on leaves that may cause green onions (scallions) to be unmarketable and dry bulb onion size to be reduced. To prevent many unnecessary insecticide applications to control nonthreatening thrips, proper identification of thrips species is an essential part of IPM. OT may not be the only thrips species present in the field (80).

OT are widely distributed throughout the United States and have multiple, overlapping generations per year on a broad range of plant species, and have an ability to disperse rapidly with short generation time (71, 74, 75). OT can reproduce asexually (parthenogenesis) and sexually, where females are produced from fertilized eggs and males are produced from unfertilized eggs. Adults and nymphs feed by piercing plant tissues with their needle-shaped mandible and draining the contents of punctured cells (74, 75). Feeding by adults and larvae produces silvery patches or streaks on the leaves, which also create entry points for pathogens. Typically, OT feed mainly within the sheaths of newly emerging onion leaves. This feeding habit makes the insects difficult to detect unless the inner leaves are examined. The damage may cause leaves to dry up, wither, turn brown, and eventually lodge (bend or break the stalk). Adults insert bean-shaped eggs into leaf tissues and can oviposit about 50 eggs per one generation, with 5-8 generations per year (75).

The primary damage caused by OT to onions is the vectoring of iris yellow spot virus (IYSV), genus *Tospovirus*, that causes significant economic losses for both onion seed and bulb crops produced in the United States (74, 75, 76). IYSV can be propagated in the vector's body and transmitted in a persistent and circulative manner and adults are capable of transmitting IYSV for life (77, 78). Large populations of OT adults move from overwintering sites to onion fields over several months beginning in early May, causing reduced-marketability from thrips feeding damage and from their vectoring of IYSV.

Thrips-transmitted tospoviruses (genus *Tospovirus*, family *Bunyaviridae*) are a major group of plant viruses affecting at least 1,090 host-plant species in 15 monocotyledonous and 69 dicotyledonous families worldwide (33, 34, 35, 36).

Natural predators such as adults and nymphs of the minute pirate bug, green lacewings and certain parasitic wasps are known to prey on plant feeding thrips. Predators of onion thrips can be numerous, but are not usually in abundance until late summer, after the majority of thrips feeding injury has occurred (74, 79). Releasing beneficial insects and predatory mites against thrips in most cases will not provide sufficient control of this pest (37). In addition, biological controls seldom reduce thrips populations below the economic injury level (74).

Successful onion thrips management programs should be focused on effective monitoring, cultural control and use of selective pesticides (73, 80). IPM encompasses the simultaneous management of multiple pests, regular monitoring of pests and their natural enemies and antagonists, use of economic or treatment thresholds when applying pesticides, and integrated use of multiple suppressive tactics. Integrated Pest Management practices that combine use of cultural and biological control, as well the use of the most selective or least-toxic insecticides, is essential for effective management of thrips.

Choosing the appropriate field location that will not be in close proximity to alfalfa, small grain fields and other onions will decrease the likelihood of OT migration into the onion fields (75). Inspecting onion transplants for thrips infestation and discarding infested plants may be helpful. Use of straw or mulch may help to reduce thrips populations and reducing nitrogen applications and using overhead irrigation may also reduce thrips densities (73). Removing or destroying debris and volunteer onion plants after harvesting may reduce overwintering sites of OT. Although these types of cultural practices recommended by universities can be useful, most growers have found them to be insufficient to provide good control of the numerous yearly populations of OT on their own.

Integrated resistance management (IRM) includes the rotation of insecticides from different chemical classes, the use of recommended rates, the limitation of maximum number of applications and product per acre per year or season, and the avoidance of sequential treatments within a single planting and across sequential crops/plantings. The use of selective insecticides that have minimal effects on natural enemy populations is vital in the control of OT. Pesticides in six activity groups – pyrethrins (IRAC, MOA Group 3), *spinosyns* (MOA Group 5), spirotetramat (MOA Group 23), neonicotinoids (MOA Group 4A), sulfoximines (Group 4C), diamides (Group 28), and organophosphates (MOA Group 1B) have varied activity against different life stages of OT ranging from suppression to control (43, 72, 75, 81, 82). However, the use of insecticides must be judicious. Decisions regarding which proper insecticides to use and timing need to be made in the context of both short term and long-term management goals. Minimizing resistance development and avoiding the flaring of onion thrip populations by their release from natural enemies need to be critical factors in insecticide use decisions. Behavior and ecology of thrips can minimize exposure to insecticides and increase risk of resistance due to species being well suited to evolve resistance to multiple classes of insecticides and because OT feed in protected areas of the plant. There have been numerous incidences of resistance reported to all major classes of insecticides from all regions of the world (47, 62, 63, 64, 65, 75). In addition to pest management failures and resistance development, there is a limited pool of

efficacious insecticides for use against OT, which further increases the selection pressure against the remaining chemistries (81, 82).

The key to managing resistance is to reduce selection pressure by rotating between insecticides with different modes of action and reducing the number of insecticide applications. Insecticides are the most common tactic for onion thrips management and their effectiveness against the pest vary based on inherent toxicity to the pesticide, as well as evolving resistance due to overdependence on a single insecticide or class of insecticides (73, 75).

As mentioned, alternating insecticides with different MOAs is a very important part of IPM programs (44, 50, 75). To manage OT during the season, a 7-10 day retreatment cycle is recommended (based on weather and pest pressure) and the same product should not be used more than two consecutive times before rotating to a different class. Rotating among the various classes delays the onset of resistance to the pesticides that are used in the management program (44). Spinetoram is an ideal and widely used rotation partner, allowing up to 5 applications at the labeled rate for OT per season.

The most efficacious insecticides for onion thrips at the present time are in the spinosyn class. No other insecticide class provides a similar level of effectiveness against onion thrips (75, 80, 81, 82). Spinosad insecticide products were first introduced in the late 1990s and have been highly effective in controlling onion thrips and widely used for this purpose (39). Spinetoram, a new and more active insecticide, was registered for use in 2008. Spinetoram represents a unique mode of action (Group 5 insecticides) and is one of the most effective insecticides to suppress OT due to the excellent control, and the fact it is a reduced risk insecticide that does not suppress beneficial populations, including minute pirate bug, *O. insidiosus*, at labeled rates (40, 41, 42).

Rotating highly active insecticides such as spinetoram with moderately active insecticides could achieve good levels of control of the onion thrips while mitigating the development of resistance to all insecticides in the rotation. Other insecticides registered for suppression of OT may not be as efficacious as spinosyns, however, they should be included as part of IPM program (44, 75).

In numerous university studies conducted across the US, spinetoram demonstrated consistent performance in the field and provided excellent control of OT (72, 73, 74, 75, 80, 81, 82). Spinetoram provides quick knockdown and residual control via contact and ingestion activity. It has a very favorable toxicity profile, with no evidence of teratogenicity, mutagenicity, carcinogenicity, or adverse reproductive effects. Compared to other registered chemicals such as oxydemeton methyl, diazinon, methyl parathion, acetamiprid, lambda cyhalothrin, methomyl and permethrin, spinetoram clearly demonstrates reduced risk to human health and non-target organisms (17). The rapid environmental degradation of spinetoram, combined with only requiring low rates for control of OT reduces the overall load of chemicals in the environment.

There is a list of registered insecticides available to manage OT on onions with wide ranging efficacy. Oxydemeton methyl (MSR), diazinon, and methyl parathion are organophosphates, which have much higher application rates when compared to spinetoram, and are more toxic to humans, beneficials and the environment (25, 31). Also, organophosphate and carbamate insecticides are considered "high risk" by FQPA and may also be unavailable for use in the near future (66).

Pyrethroids labeled in onions for OT control include Ambush, Ammo, Proaxis, Mustang Max, Pounce, Warrior II and Lannate (carbamate). These are "restricted use pesticides (RUP)" due to acute toxicity to humans with possible oncogenicity and/or toxicity to fish, birds and aquatic organisms (19). In addition, the pyrethroid class of chemistry is broad spectrum and controls both beneficial and pest insects. Predator insects may be susceptible to a lower dose than the pest, disrupting the predator-prey relationship, which leads to flaring up of mites and knocking down of beneficial insects (18, 20, 21). Furthermore, there is documented resistance of OT to organophosphate (e.g.,



azinphosmethyl, diazinon, and methyl parathion) and synthetic pyrethroid (e.g., cypermethrin, and permethrin) insecticides across the United States (75, 83, 84, 85, 86). Managing resistance to insecticides is critical and can be mitigated by limiting the frequency of insecticide applications, rotating insecticides used in a sequence, and maintaining thorough coverage to prolong the effectiveness of insecticides (75, 87).

Neonicotinoids such as Assail can be a good rotation partner for spinetoram due to its different class of chemistry. Agri-Mek (MOA Group 6) and Exirel (MOA Group 28) provide variable control of onion thrip larvae and adults, and have short residual activity (5–7 days). Movento (MOA Group 23) is highly efficacious against larval OT infestations, but it does not provide satisfactory control against adults or late in the season when plants are maturing (75). Horticultural oil, natural pyrethrins (plus piperonyl butoxide), or insecticidal soap have very low efficacy (75).

When compared to other registered products for OT control in onions, spinetoram offers the lowest REI and the lowest PHI (1 day) compared to PennCap-M, Diazinon, Assail, Warrior II and Lannate (68, 80).

Spinetoram is a significant tool for IPM programs for control of OT in onions, and has the following benefits:

- It is a selective insecticide with a low use rate.
- It has a favorable toxicological profile and is soft on beneficial arthropods.
- The unique mode of action of spinetoram makes it effective against OT as well as providing an alternate mode of action for managing against pesticide resistance.
- It is an ideal and widely used rotation partner, allowing up to 5 applications at the labeled rate per calendar year in peppers.

## Lettuce

One of major pests in the minor use crop lettuce (*Lactuca sativa*), is western flower thrips, *Frankliniella occidentalis* (Perganda), (WFT). WFT are widely distributed throughout the United States, feed on a wide variety of plants and have multiple, overlapping generations per year on a broad range of plant species (71). Adults and larvae feed by piercing plant tissues with their needle-shaped mandible and then draining the contents of the punctured cells (47, 54, 109). Feeding by adults and larvae produces silvery appearance that eventually turns to brown scarring and can be confused with windburn or blown sand damage. Typically, WFT adults insert bean-shaped eggs into leaf tissue and can survive for 4-5 weeks, and ovipositing approximately 50 eggs per generation. Thrips also cause damage on outside leaves of head lettuce and can contaminate the inside of heads at harvest.

The primary damage caused by WFT to lettuce is the vectoring of tomato spotted wilt virus and impatiens necrotic spot virus, genus *Tospovirus* (109). The virus can only be acquired by the immature larva stage of thrips, whereas plant-to-plant transmission primarily occurs by adults. The adult thrips can transmit the virus for the remainder of their lives, which can last 30 to 45 days. Large populations of WFT migrate into the spring crop of lettuce from weeds and other vegetation, causing reduced-marketability from thrips feeding damage and from their vectoring of tomato spotted wilt virus (69, 70). On lettuce plants, adults reproduce and rapidly colonize into large populations (109).

Thrips feeding on plants can damage leaves and noticeably affect plants' cosmetic appearance. Significant crop damage leading to severe yield losses can be caused by thrips that act as a vector of Tospoviruses (32). Thrips-transmitted tospoviruses (genus *Tospovirus*, family Bunyaviridae) are a major group of plant viruses affecting at least 1,090 host-plant species in 15 monocotyledonous and 69 dicotyledonous families worldwide (33, 34, 35, 36). The actual amount of economic losses attributable to tomato spotted wilt virus alone was over US\$1 billion in losses annually on a global basis. This estimate did not include the direct damage caused by WFT (56, 57).

Insect control in high value fresh market lettuce grown in the desert southwest have relied almost exclusively on insecticides to control a complex of mobile pests such as WFT. Because lettuce is a short season annual crop with no tolerance for insect damage or contamination, biological control is generally considered unacceptable. Western lettuce growers cannot rely on beneficial insects and predatory mites against thrips in most cases as it will not provide sufficient control of this pest to meet marketing demands for fresh produce. Additionally, fresh market grading standards do not allow for contaminants such as pest or predatory species in fresh produce (120). Thus any insect species including natural predators found on marketable lettuce crops at harvest is not tolerated (120).

Successful WFT management programs should be focused on cultural controls, effective monitoring and identification of the thrips species, and use of selective pesticides (37, 38, 121). Crop placement can help if growers avoid planting lettuce crops near grain fields, weedy drains or fields or grassy areas. Overhead sprinkler irrigation has been shown to suppress WFT numbers in romaine and spinach by as much as 50%, but insecticide treatments are generally still necessary, particularly in late spring when WFT adult dispersal is high (121). Cultural methods do not provide effective control of thrip populations during the critical spring months (113).

Integrated resistance management (IRM) includes the rotation of insecticides from different chemical classes, the use of recommended rates, the limitation of maximum number of applications and product per acre per year or season, and the avoidance of sequential treatments within a single planting and across sequential crops/plantings.

Pesticides in four activity groups – pyrethrins (IRAC, MOA Group 3), *spinosyns* (MOA Group 5), neonicotinoids (MOA Group 4A), sulfoximines (MOA Group 4C) and organophosphates (MOA Group 1B) have varied activity against WFT ranging from suppression to control (43, 114, 115, 116). Insecticides continue to have an important role to play in WFT management, although the use of insecticides must be judicious. Decisions regarding which insecticides are to be used and the timing should be made in the context of both short term and long-term management goals. Minimizing resistance development and avoiding the flaring of WFT populations by their release from natural enemies are critical factors in insecticide use decisions (44). Behavior and ecology of WFT can minimize exposure to insecticides and increase the risk of resistance due to the species being well suited to evolve resistance to multiple classes of insecticides. There have been numerous incidences of resistance reported to all major classes of insecticides from all regions of the world (102, 103, 104, 105, 106, 107, 108). In addition to pest management failures and resistance development, there is a limited pool of efficacious insecticides for use against WFT (47), which further increases the selection pressure on the remaining chemistries.

The key to managing resistance is to reduce selection pressure by rotating between insecticides with different modes of action and reducing the number of insecticide applications. It may be necessary to use nonchemical control methods and rotate to insecticides that may not provide the highest level of control.

The negative impacts of broad-spectrum insecticides, such as pyrethroids, on natural enemies and competitor species, in addition to continuous, overlapping generations typical for WFT populations in crops makes it especially challenging to implement IRM strategy (45).

The most efficacious insecticides for WFT, at present, are in the spinosyn class. No other insecticide class provides a similar level of effectiveness against WFT (44, 46). Spinosad insecticide products were first introduced in the late 1990s and have been highly effective in controlling WFT and widely used for this purpose (39). Spinetoram, a new and more active insecticide, was registered for use in 2008. Spinetoram represents a unique mode of action (Group 5 insecticides) and is one of the most effective insecticides to suppress WFT due to excellent control and softness of beneficial at labeled rates. Rotating highly active insecticides such as spinetoram with moderately active insecticides could achieve good levels of control of the WFT while mitigating the development of resistance to all insecticides in

the rotation. Other insecticides registered for suppression of WFT may not be as efficacious as spinosyns, however, they should be included as part of IPM program (44).

As mentioned, alternating insecticides with different MOAs is a very important part of IPM programs (44, 50). To manage WFT during the season, a 3-7 day retreatment cycle is recommended, and the same product should not be used more than two consecutive times before rotating to a different class. Rotating among the various classes delays the onset of resistance to the pesticides that are used in the management program (44).

In numerous university studies across the US, spinetoram demonstrated consistent performance in the field and provided excellent control of WFT (40, 41, 42, 44, 46, 50). Spinetoram provides quick knockdown and residual control via contact and ingestion activity. It has a very favorable toxicity profile, with no evidence of teratogenicity, mutagenicity, carcinogenicity, or adverse reproductive effects. Compared to other registered chemicals such as dimethoate, acetamiprid, beta-cyfluthrin, zeta-cypermethrin and methomyl, spinetoram clearly demonstrates reduced risk to human health and non-target organisms (120). The rapid environmental degradation of spinetoram, combined with only needing low rates for control of WFT reduces the overall load of chemicals in the environment. There is a list of registered insecticides available to manage WFT on lettuce with widely ranging efficacy. Dimethoate is an organophosphate, which has much higher application rates, and is more toxic to humans, beneficials and the environment when compared to Spinetoram (25, 31). Also, organophosphate and carbamate insecticides are considered "high risk" by FQPA and may be unavailable for use in the near future (66). In addition, dimethoate and methomyl have low efficacy on WFT and can cause spider mite outbreaks (67).

Furthermore, there are large number of pyrethroids that are labeled in lettuce for WFT control such as Ambush, Asana, Baythroid, Brigade, Mustang Max, Warrior II and Lannate (carbamate) that are "restricted use pesticides (RUP)" due to acute toxicity to humans with possible oncogenicity and/or toxicity to fish, birds and aquatic organisms (19, 117). Neonicotinoids, such as Actara, Belay, Assail, and Scorpion, can be a good rotation partner for spinetoram because it is a different class of chemistry with a different MOA. Horticultural oil, natural pyrethrins (plus piperonyl butoxide), or insecticidal soap have very low efficacy and requires multiple applications (109, 116, 119).

When compared to other registered products for WFT control in lettuce, spinetoram offers the lowest REI and the lowest PHI (1 day) compared to Asana, Assail, Mustang Max and Belay (109).


Spinetoram is a significant tool for IPM programs for control of WFT in lettuce, and has the following benefits:

- It is a selective insecticide with a low use rate.
- It has a favorable toxicological profile.
- The unique mode of action of spinetoram makes it effective against WFT as well as providing an alternate mode of action for managing against pesticide resistance.
- It is an ideal and widely used rotation partner, allowing up to 6 applications at the labeled rate per calendar year in lettuce.

**Conclusion:**

For the reasons set forth above, DAS requests the Agency to grant an exclusive use period extension on Spinetoram of 3 years, making the end date September 28, 2020. Should you have any questions or wish to discuss this matter further, please do not hesitate to contact me. Thank you for your assistance.

Sincerely,

A handwritten signature in black ink that reads "J Hughes". The signature is written in a cursive style with a long horizontal stroke at the end.

Jennifer Hughes  
U.S. Regulatory Manager  
317-337-7993  
JLHughes@dow.com

References:

1. <http://insect.pnwhandbooks.org/pnw-insect-management-handbook/emerging-pest-spotted-wing-drosophila>
2. [www.ipm.msu.edu/SWD.htm](http://www.ipm.msu.edu/SWD.htm)
3. New England Small Fruit Management Guide, 2013-2014.
4. <http://spottedwing.org/>
5. [http://entmemdept.ufl.edu/liburd/fruitnvegipm/presentations/SWD\\_recommendations.pdf](http://entmemdept.ufl.edu/liburd/fruitnvegipm/presentations/SWD_recommendations.pdf)
6. <http://extension.psu.edu/plants/tree-fruit/news/2012/spotted-wing-drosophila-management>.
7. Commercial Blueberry Pest Control Recommendations for New Jersey, 2015, (<https://njaes.rutgers.edu/pubs/publication.asp?pid=e265>)
8. <http://insect.pnwhandbooks.org/pnw-insect-management-handbook/emerging-pest-spotted-wing-drosophila>
9. SWD Management Recommendations for Michigan Blueberry., [http://www.ipm.msu.edu/invasive\\_species/spotted\\_wing\\_drosophila/crop\\_recommendations](http://www.ipm.msu.edu/invasive_species/spotted_wing_drosophila/crop_recommendations)
10. <http://utahpests.usu.edu/ipm/html/fruits/fruit-insect-disease/spotted-wing-drosophila>
11. [http://pubs.ext.vt.edu/456/456-017/Section-2\\_Commercial\\_Small\\_Fruit-1.pdf](http://pubs.ext.vt.edu/456/456-017/Section-2_Commercial_Small_Fruit-1.pdf)
12. [http://labs.russell.wisc.edu/swd/management-2/#Chemical\\_Control](http://labs.russell.wisc.edu/swd/management-2/#Chemical_Control)
13. <http://www.fruit.cornell.edu/spottedwing/pdfs/BerrySWDinsecticidemanagement.pdf>
14. <http://ufdc.ufl.edu/UFE0046031/00001>
15. [http://spottedwing.org/system/files/Spotted\\_Wing\\_booklet-11-2.pdf](http://spottedwing.org/system/files/Spotted_Wing_booklet-11-2.pdf)
16. <http://www.isaacslab.ent.msu.edu/Images/2014%20Wise%20et%20al%20-%20Curative%20activity%20SWD.pdf>
17. XDE-175 Reduced risk Rationale, EPA, MRID 46695901
18. <http://www.growingproduce.com/vegetables/vegetable-insect-control-options-in-the-southeast/>
19. <http://www.pesticides.montana.edu/reference/labels.htm>
20. <https://www.beyondpesticides.org/info/services/pesticidefactsheets/toxic/pvretroid.php>
21. <http://thealmonddoctor.com/2010/07/26/pyrethroid-vs-reduced-risk-product-usage-for-hullsplitmay-spray/>
22. <http://edis.ifas.ufl.edu/in998>
23. <http://www.ipm.ucdavis.edu/PMG/r71300711.html>
24. <http://www.extension.umn.edu/garden/insects/find/spotted-wing-drosophila-in-home-gardens/>
25. <http://www.ipm.ucdavis.edu/PMG/PESTNOTES/pn74158.html>
26. <http://www fruitedge.umn.edu/spotted-wing-drosophila-management-recommendations-for-minnesota-berry-growers/>
27. <http://www.fruit.cornell.edu/spottedwing/pdfs/BerrySWDinsecticidemanagement.pdf>
28. <http://www.ipm.msu.edu/uploads/files/SWD/ManagementRecommendations-RaspberryBlackberrySep2012.pdf>
29. [http://spottedwing.org/system/files/Caneberry%20SWD%20Pesticides%20for%20OR%20and%20WA%20February%202014\\_0.pdf](http://spottedwing.org/system/files/Caneberry%20SWD%20Pesticides%20for%20OR%20and%20WA%20February%202014_0.pdf)
30. <http://ienny.tfrec.wsu.edu/opm/displaySpecies.php?pn=240>
31. <http://www.ipm.ucdavis.edu/PMG/r734301211.html>
32. [http://www.caes.uga.edu/topics/diseases/tswv/documents/thrips\\_vectors\\_of\\_tospoviruses.pdf](http://www.caes.uga.edu/topics/diseases/tswv/documents/thrips_vectors_of_tospoviruses.pdf)
33. Parrella, G., P. Gognalons, K. Gebre-Selassie, C. Vovlas, and G. Marchoux., 2003. An update of the host range of Tomato spotted wilt virus. *Journal of Plant Pathology* 85: 227–264.
34. Ullman, D. E., J. L. Sherwood, and T. L. German. 1997. Thrips as vectors of plant pathogens, pp. 539–565. In T. Lewis (ed), *Thrips as Crop Pests*. CAB International, New York.
35. Jones, D. R. 2005. Plant viruses transmitted by thrips. *European Journal of Plant Pathology* 113: 119–157.
36. Pappu, H. R., R.A.C. Jones, and R. K. Jain. 2009. Global status of *Tospovirus* epidemics in diverse cropping systems: successes achieved and challenges ahead. *Virus Research* 141: 219–236.
37. <http://www.ipm.ucdavis.edu/PMG/PESTNOTES/pn7429.html>
38. <http://journals.fcla.edu/flaent/article/view/75899>

39. Eger, JR., J. E., Stavisky, J., and Funderburk, J. E. 1998. Comparative toxicity of spinosad to *Frankliniella* spp. (Thysanoptera: Thripidae), with notes on a bioassay technique. *Florida Entomol* 81: 542-551.
40. Funderburk, J., Stavisky, J., and Olson, S. 2000. Predation of *Frankliniella occidentalis* (Thysanoptera: Thripidae) in field peppers by *Orius insidiosus* (Hemiptera: Anthocoridae). *Environ. Entomol.* 29: 376-382.
41. Reitz, S. R., Yearby, E. L., Funderburk, J. E., Stavisky, J., Momol, M. T., and Olson, S. M. 2003. Integrated management tactics for *Frankliniella* thrips (Thysanoptera: Thripidae) in field-grown pepper. *J. Econ. Entomol.* 96: 1201-1214.
42. Srivistava, M., Bosco, L., Funderburk, J., Olson, S., and Weiss, A. 2008. Spinetoram is compatible with the key natural enemy of *Frankliniella* species thrips in pepper. *Plant Health Progress* doi:10.1094/PHP-2008-0118-02-RS.
43. [http://programs.ifas.ufl.edu/u-scout/Tutor\\_files/2014\\_VPH.pdf](http://programs.ifas.ufl.edu/u-scout/Tutor_files/2014_VPH.pdf)
44. [http://www.ars.usda.gov/sp2UserFiles/person/11884/2012\\_Reitz\\_Funderburk\\_01.pdf](http://www.ars.usda.gov/sp2UserFiles/person/11884/2012_Reitz_Funderburk_01.pdf)
45. <http://journals.fcla.edu/flaent/article/view/83159>
46. Dripps, J. E., Gomez, L. E., Weiss, A. W., Funderburk, J., Castro, B. A., and Paroonagian, D. L. 2010. Insecticide rotation as a component of thrips resistance management programs. *Resist. Pest Mgt. Newsl.* 19: 32-35.
47. <http://cdn.intechopen.com/pdfs-wm/28269.pdf>
48. <http://www.smallfruits.org/smallfruitsregguide/guides/2014/2014sestrawberrvipmguid.pdf>
49. Jacobson, A.L., & Kennedy, G.G. (2011) The effect of three rates of cyantraniliprole on the transmission of tomato spotted wilt virus by *Frankliniella occidentalis* and *Frankliniella fusca* (Thysanoptera: Thripidae) to *Capsicum annuum*. *Crop Protection* 30: 512-515.
50. [http://whalonlab.msu.edu/wp-content/uploads/2012/06/vol.19\\_no.2.pdf](http://whalonlab.msu.edu/wp-content/uploads/2012/06/vol.19_no.2.pdf)
51. Hansen, E.A., Funderburk, J.E., Reitz, S.R., Ramachandran, S., Eger, J.E., & McAuslane, H. (2003) Within-plant distribution of *Frankliniella* species (Thysanoptera: Thripidae) and *Orius insidiosus* (Heteroptera: Anthocoridae) in field pepper. *Environmental Entomology* 32: 1035-1044.
52. <http://ir.library.oregonstate.edu/xmlui/bitstream/handle/1957/53201/ReitzStuartCropSoilScienceImpactsNaturalEnemies.pdf?sequence=1>
53. Frantz, G., & Mellinger, H.C. (2009) Shifts in WFT, *Frankliniella occidentalis* (Thysanoptera: Thripidae), population abundance and crop damage. *Florida Entomologist* 92: 29-34.
54. Kirk, W.D.J. (1997b) Feeding, In: T. Lewis (Ed.), *Thrips as crop pests*, CAB International, New York. pp. 119-174.
55. Childers, C.C. (1997) Feeding and oviposition injuries to plants, In: T. Lewis (Ed.), *Thrips as crop pests*, CAB International, New York. pp. 505-537.
56. Kirk, W.D.J., & Terry, L.I. (2003) The spread of the WFT *Frankliniella* occidentalis (Pergande). *Agricultural and Forest Entomology* 5: 301-310.
57. Reitz, S.R., Gao, Y.L., & Lei, Z.R. (2011) Thrips: Pests of concern to China and the United States. *Agricultural Sciences in China* 10: 867-892.
58. <http://westernfarmpress.com/vegetables/thrips-control-hinges-multiple-practices>
59. Baez, L., Reitz, S.R., Funderburk, J.E., & Olson, S.M. (2011) Variation within and between *Frankliniella* thrips species in host plant utilization. *Journal of Insect Science* 11: 41.
60. Pfannenstiel, R.S., & Yeargan, K.V. (1998) Association of predaceous Hemiptera with selected crops. *Environmental Entomology* 27: 232-239.
61. Bielza, P., Quinto, V., Contreras, J., Torne, M., Martin, A., & Espinosa, P.J. (2007b) Resistance to spinosad in the WFT, *Frankliniella occidentalis* (Pergande), in greenhouses of south-eastern Spain. *Pest Management Science* 63: 682-687.

63. Dağlı, F., & Tunç, I. (2007) Insecticide resistance in *Frankliniella occidentalis* (Pergande) (Thysanoptera: Thripidae) collected from horticulture and cotton in Turkey. *Australian Journal of Entomology* 46: 320-324.
64. Kay, I.R., & Herron, G.A. (2010) Evaluation of existing and new insecticides including spirotetramat and pyridalyl to control *Frankliniella occidentalis* (Pergande) (Thysanoptera: Thripidae) on peppers in Queensland. *Australian Journal of Entomology* 49: 175-181.
65. Weiss, A., Dripps, J.E., & Funderburk, J. (2009) Assessment of implementation and sustainability of integrated pest management programs. *Florida Entomologist* 92: 24- 28.
66. <http://fshs.org/proceedings-o/2003-vol-116/161-164.pdf>
67. <http://www.ipm.ucdavis.edu/PMG/r783302011.html>
68. <http://edis.ifas.ufl.edu/cv137>
69. Funderburk, J. E., Stavisky, J., and Olson, S. 2000. Predation of *Frankliniella occidentalis* (Thysanoptera: Thripidae) in field peppers by *Orius insidiosus* (Hemiptera: Anthocoridae). *Environ. Entomol.* 29:376-382.
70. Reitz, S. R., Yearby, E. L., Funderburk, J. E., Stavisky, J., Olson, S. M., and Momol, M. T. 2003. Integrated management tactics for *Frankliniella* thrips (Thysanoptera: Thripidae) in field-grown pepper. *J. Econ. Entomol.* 96:1201-1214
71. <https://edis.ifas.ufl.edu/in401>
72. [http://rvpadmin.cce.cornell.edu/pdf/veg\\_edge/pdf14\\_pdf.pdf](http://rvpadmin.cce.cornell.edu/pdf/veg_edge/pdf14_pdf.pdf)
73. <http://extension.usu.edu/files/publications/factsheet/ent-117-08pr.pdf>
74. <http://cru.cahe.wsu.edu/CEPublications/FS126E/FS126E.pdf>
75. <http://ijpm.oxfordjournals.org/content/6/1/6>
76. Kritzman, A., M. Lampel, B. Raccach, and A. Gera. 2001. Distribution and transmission of Iris yellow spot virus. *Plant Dis.* 85: 838-842.
77. Ullman, D. E., J. J. Cho, R.F.L. Mau, D. M. Westcott, and D. M. Custer. 1992. A midgut barrier to Tomato spotted wilt virus acquisition by adult WFT. *Phytopathology* 82: 1333-1342.
78. Wijkamp, L., J. van Lent, R. Kormelink, R. Goldbach, and D. Peters. 1993. Multiplication of Tomato spotted wilt virus in its insect vector, *Frankliniella occidentalis*. *J. Gen. Virol.* 74: 341-349.
79. Fok, E. J., J. D. Petersen, and B. A. Nault. 2014. Relationships between insect predator populations and their prey, *Thrips tabaci*, in onion fields grown in large-scale and small-scale cropping systems. *BioControl* 59: 739-748.
80. <http://www.ipm.ucdavis.edu/PMG/r584300111.html>
81. [http://rvpadmin.cce.cornell.edu/uploads/doc\\_320.pdf](http://rvpadmin.cce.cornell.edu/uploads/doc_320.pdf)
82. <http://insect.pnwhandbooks.org/sites/insect.pnwhandbooks.org/files/pdfsection/vegetable.pdf>
83. Cranshaw, W. S. 1989. Control of organophosphate resistant onion thrips, 1986. *Insect. Acaricide Tests* 14: 128.
84. Davis, M., Y. Anandwala, M. Bommarito, and E. Grafus. 1995. Onion thrips control, 1994. *Arthropod Manag. Tests* 20: 99.
85. Shelton, A. M., B. A. Nault, J. Plate, and J. -Z. Zhao. 2003. Regional and temporal variation in susceptibility to k-cyhalothrin in onion thrips, *Thrips tabaci* (Thysanoptera: Thripidae), in onion fields in New York. *J. Econ. Entomol.* 96: 1843-1848.
86. Shelton, A. M., J. -Z. Zhao, B. A. Nault, J. Plate, F. R. Musser, and E. Larentzaki. 2006. Patterns of insecticide resistance in onion thrips (Thysanoptera: Thripidae) in onion fields in New York. *J. Econ. Entomol.* 99: 1798-1804.
87. Gill, H. K., and H. Garg. 2014. Pesticide: Environmental impacts and management strategies, pp. 187-230. In S. Solenski, and M. L. Larramenday (eds.), *Pesticides- Toxic Effects*. Intech. Rijeka, Croatia. (<http://www.intechopen.com/books/pesticides-toxic-aspects/pesticides-environmental-impacts-andmanagement-strategies>) (last accessed August 2015).
88. <http://www.umass.edu/cranberry/downloads/Black-Headed%20Fireworm.pdf>

89. Fitzpatrick, S.M. and J.T. Troubridge. 1993. Fecundity, number of diapause eggs, and egg size of successive generations of the blackheaded fireworm (Lepidoptera: Tortricidae) on cranberries. *Environ. Entomol.* 22: 818-823.
90. Maurice, C., C. Bedard, S.M. Fitzpatrick, J. Troubridge, and D. Henderson. 2000. Integrated pest management for cranberries in western- Canada- a guide for identification, monitoring and control of key pests and diseases in cultivated cranberry fields in western Canada. Technical report 163 pacific agri-food research centre-agassiz.
91. [http://www.umass.edu/cranberry/pubs/bmp\\_insect.html](http://www.umass.edu/cranberry/pubs/bmp_insect.html)
92. <http://fruit.wisc.edu/wp-content/uploads/2011/05/Understanding-Biological-Control-and-its-Potential-for-Managing-Insect-Pests-on-Cranberry-in-Wisconsin.pdf>
93. [http://www.bccranberries.com/pdfs/ipm\\_guide.pdf](http://www.bccranberries.com/pdfs/ipm_guide.pdf)
94. <http://www.umass.edu/cranberry/downloads/bmp/IPM%20BMP.pdf>
95. <http://fruit.wisc.edu/wp-content/uploads/2011/06/Cranberry-Pest-Management-in-Wisconsin.pdf>
96. <http://www.reeis.usda.gov/web/crisprojectpages/1000372-host-plant-resistance-in-cranberry-varietal-susceptibility-of-cranberry-to-three-major-lepidopteran-pests.html>
97. <http://scholarworks.umass.edu/cgi/viewcontent.cgi?article=1194&context=cranchart>
98. [http://msdssearch.dow.com/PublishedLiteratureDAS/dh\\_08aa/0901b803808aa289.pdf?filepath=/010-21842.pdf&fromPage=GetDoc](http://msdssearch.dow.com/PublishedLiteratureDAS/dh_08aa/0901b803808aa289.pdf?filepath=/010-21842.pdf&fromPage=GetDoc)
99. <http://njaes.rutgers.edu/pubs/plantandpestadvisory/2010/cb042910.pdf>
100. <http://plant-pest-advisory.rutgers.edu/post-bloom-insect-pest-control-recommendations-2/>
101. <http://longbeach.wsu.edu/cranberries/documents/pattenbcwintercongress2013presentation.pdf>
102. Bielza, P., Quinto, V., Contreras, J., Torne, M., Martin, A., & Espinosa, P.J. (2007b) Resistance to spinosad in the WFT, *Frankliniella occidentalis* (Pergande), in greenhouses of south-eastern Spain. *Pest Management Science* 63: 682-687.
103. Dağlı, F., & Tunç, I. (2007) Insecticide resistance in *Frankliniella occidentalis* (Pergande) (Thysanoptera: Thripidae) collected from horticulture and cotton in Turkey. *Australian Journal of Entomology* 46: 320-324.
104. Jensen, S.E. (2000a) Insecticide resistance in the WFT, *Frankliniella occidentalis*. *Integrated Pest Management Reviews* 5: 131-146.
105. Kay, L.R., & Herron, G.A. (2010) Evaluation of existing and new insecticides including spirotetramat and pyridalyl to control *Frankliniella occidentalis* (Pergande) (Thysanoptera: Thripidae) on peppers in Queensland. *Australian Journal of Entomology* 49: 175-181.
106. Morishita, M. (2001) Toxicity of some insecticides to larvae of *Frankliniella occidentalis* (Pergande) (Thysanoptera: Thripidae) evaluated by the petri dish-spraying tower method. *Applied Entomology and Zoology* 36: 137-141.
107. Weiss, A., Dripps, J.E., & Funderburk, J. (2009) Assessment of implementation and sustainability of integrated pest management programs. *Florida Entomologist* 92: 24-28.
108. Zhao, G., Liu, W., Brown, J.M., & Knowles, C.O. (1995) Insecticide resistance in field and laboratory strains of WFT (Thysanoptera: Thripidae). *Journal of Economic Entomology* 88: 1164-1170.
109. <http://www.ipm.ucdavis.edu/PMG/r441302111.html>
110. <http://extension.arizona.edu/sites/extension.arizona.edu/files/pubs/az1177-1g.pdf>
111. <http://extension.arizona.edu/sites/extension.arizona.edu/files/pubs/az1177-1f.pdf>
112. [http://calgreens.org/control/uploads/Thrips\\_Management\\_in\\_Lettuce\\_11.pdf](http://calgreens.org/control/uploads/Thrips_Management_in_Lettuce_11.pdf)
113. <http://www.ipmcenters.org/cropprofiles/docs/azlettuce.pdf>
114. <http://www.ipmcenters.org/cropprofiles/docs/azlettuce.pdf>
115. <http://cals.arizona.edu/crops/vegetables/insects/veginspub.html#thrips>
116. <http://edis.ifas.ufl.edu/features/handbooks/vegetableguide.html>



117. [https://extension.arizona.edu/sites/extension.arizona.edu/files/resources/040115%20LIL\\_%2010%20yr%20summary.pdf](https://extension.arizona.edu/sites/extension.arizona.edu/files/resources/040115%20LIL_%2010%20yr%20summary.pdf)
118. <http://www.bioone.org/doi/pdf/10.1653/024.092.0101>
119. <https://edis.ifas.ufl.edu/in1078>
120. IPM for fresh-market lettuce production in the desert southwest: the produce paradox. John C Palumbo Department of Entomology, University of Arizona, Yuma, AZ 85364, USA. Pest Manag Sci 65:1311-20. 2009
121. [http://cals.arizona.edu/crops/pdfs/020613%20Insect%20Management%20on%20Desert%20Produce\\_Thrips\\_2013.pdf](http://cals.arizona.edu/crops/pdfs/020613%20Insect%20Management%20on%20Desert%20Produce_Thrips_2013.pdf)

n	Pyrethroid	3	Excellent	16 oz	3	24	2	Poor
trin	Pyrethroid	3	Excellent	4 oz	1	12	6	Poor
e	Pyrethroid	3	Excellent	9.6 oz	14	12	7	Poor
	Pyrethroid	3	Excellent	16 oz	1	12	6	Poor
	Carbamate	1A	Excellent	1lb	3	48	4	Poor
ole	Diamide	28	Excellent	13.5 oz	3	12	4	Good
l	Neonicotinoid	4A	Poor	5.3 oz	1	12	5	Poor
	Pyrethrin	3	Poor	64 oz	0.5	12	-	Very good
a	Spinosyn	5	Excellent	3-6oz	3	4	6	Good

d by IRAC = Insecticides Resistance Action Committee