

Microcystin-LR – Drinking Water Guidance in Minnesota



Inland HAB Discussion Group
January 24, 2013




Source: MPCA

Julia Dady
Minnesota Department of Health
Health Risk Assessment Unit

Minnesota Concerns

- ❑ “Land of 10,000 Lakes” – many lakeshore residential properties, agricultural lands
- ❑ Ambient/Recreational Water Quality – Swimming, boating, fishing, livestock, pets
- ❑ Drinking Water Quality
 - Surface source-waters
 - Groundwater under the influence of surface waters



Source: MPCA

MC = microcystin

MC-LR in Minnesota – Occurrence

- ❑ Interagency Work Group formed, 2004
 - Several dog deaths – algal blooms suspected; MPCA met with MDH, DNR, and the MN Vet. Med. Assoc.

- ❑ MPCA/USGS Surveys, 2006
 - Up to 12 MN eutrophic lakes

- ❑ EPA National Lake Assessment Project, 2007
 - MPCA, DNR, MDA
 - 50 MN lakes, mid- to late summer, multiple “ecoregions” – not just eutrophic lakes

MC-LR = Microcystin-LR

MPCA = MN Pollution Control Agency; DNR = MN Department of Natural Resources; MDA = MN Dept. of Agriculture

MC-LR in Minnesota

– Public Consultations



Source: MPCA

- ❑ MDH Site Consultation, July 2011 – Little Rock Lake
 - MCs detected @ 38,000 $\mu\text{g/L}$ and $> 80,000 \mu\text{g/L}$ in lake in 2007
 - Near-shore shallow residential wells sampled in Aug-Sept 2011
 - all non-detect during mild algal bloom, but wells considered vulnerable

- ❑ MDH Consultations – Budd Lake, Fairmont MN
 - Budd Lake = drinking water source for city
 - Citizen concerns raised, summer 2012

MDH Guidance Development

– Selection of Microcystin-LR

- Drinking Water Contaminants of Emerging Concern (CEC) program
 - MN Clean Water, Land and Legacy constitutional amendment, Nov. 2008
 - Nominations (Public, State Agencies, MDH Staff, etc.)
 - MDH screening and ranking for priority

- MC-LR Nominated to MDH's CEC program by MPCA in April 2011
 - MDH-ranked as high priority based on toxicity and exposure factors

MDH Guidance – MC-LR

- Health-Based Value (HBV) = $0.04 \mu\text{g/L}$
 - for short-term, subchronic and chronic durations
 - insufficient data for acute guidance
- Guidance posted on MDH website
 - Sept. 2012

MDH Guidance Development – MC-LR

❑ Critical Studies Selected

- Short-term – Heinze, 1999; 28-day drinking water study in rats, serum liver enzymes, ↑ rel liver wt, liver lesions
- Subchronic – Fawell et al., 1999; 13-wk gavage study in mice, serum liver enzymes, ↑ rel liver wt
- Chronic – Fawell et al., 1999; 13-wk gavage study in mice, serum liver enzymes, ↑ rel liver wt, degenerative liver lesions, Kupffer cell activation in liver

❑ POD (Dose causing no harm to animals)

- Short-term – 6.4 $\mu\text{g}/\text{kg}\text{-d}$
- Subchronic & Chronic - 58 $\mu\text{g}/\text{kg}\text{-d}$

Drinking Water Guidance Values: MDH & WHO Comparison

Short-term and Subchronic:

MDH (0.04 ug/L)

WHO (n/a)

Chronic:

MDH (0.04 ug/L) **25 x lower!**




WHO (1 ug/L)

Microcystin-LR - Chronic Duration

Parameter	WHO	MDH
Study Basis	Fawell et al. 1999, 13-week gavage study in mice	Fawell et al. 1999, 13-week gavage study in mice
Health Endpoints	Liver	Liver
Animal Dose Level causing no harm (POD)	0.04 mg/kg-d (NOAEL)	0.058 mg/kg-d (EPA BMDL _{1SD})
Human Equivalent Dose (HED)	n/a	0.0081 mg/kg-d (per EPA 2011 guidance)
Uncertainty Factors	1000 (10 interspecies, 10 intraspecies, 10 database uncertainty)	1000 (3 interspecies, 10 intraspecies, 10 database uncertainty, 3 subchronic to chronic)
Human Dose Level expected to cause no harm (RfD)	0.00004 mg/kg-d	0.0000081 mg/kg-d ✓ 5 x lower than WHO
Drinking water allocation factor (RSC)	0.8	0.8
Drinking water intake rates	0.0333	0.043 ✓
Drinking water guidance value (HBV)	1 ug/L	0.2 ug/L (calculated) 0.04 ug/L (set to short-term) ✓

$$HBV = [RfD \times RSC \times 1000] / \text{Intake Rate}$$

Microcystin-LR Short-term Duration (1 to 30 days)

Parameter	WHO	MDH
Study Basis	n/a	Heinze 1999, 28-day drinking water study in rats 
Health Endpoints	n/a	Liver
Animal Dose Level causing no harm (POD)	n/a	0.0064 mg/kg-d (EPA BMDL ₁₀)
Human Equivalent Dose (HED)	n/a	0.0015 mg/kg-d
Uncertainty Factors	n/a	100 (3 interspecies, 10 intraspecies, 3 database uncertainty)
Human Dose Level expected to cause no harm (RfD)	n/a	0.000015 mg/kg-d
Drinking water allocation factor (RSC)	n/a	0.8
Drinking water intake rates (infant intake rate)	n/a	0.289 L/kg bw-day 
Drinking water guidance value (HBV)	n/a	0.04 ug/L 


6-7 x higher than chronic intake rate



$$HBV = [RfD \times RSC \times 1000] / \text{Intake Rate}$$

Age-Adjusted Drinking Water Intake Rates

- 2004 EPA Estimated Water Ingestion in the U.S.
- 2006 Draft/2008 Final EPA Child-Specific Exposure Factors Handbook



Microcystin-LR - Cancer

- ❑ IARC Group 2B carcinogen (possibly carcinogenic to humans) - Liver, colon
- Tumor promotor -
 - “Strong” evidence for tumor promotion
 - Threshold dose is likely
- Epidemiology drinking water studies –
 - Reports of associations w/ cancer at microcystin levels ranging from 0.1 to 2 ug/L
 - Reference populations (control groups) reported to have exposures up to 0.04 ug/L [note that this is also the same as MDH HBV]
- ❑ The MDH non-cancer HBV is considered protective for potential carcinogenicity.

Toxicology issues: Further study needed?

- ❑ Male Reproductive Effects?? (Chen et al. 2011)
 - Sperm, hormones, testes – may be more sensitive than liver?
 - Mouse study, drinking water, “potential” HBV 4x lower
 - Addressed in database uncertainty factor
 - MDH RfDs considered protective (i.e., 6-11 times lower than the LOAEL_{HED} from repro study)

- ❑ Other microcystin congeners, MC-LF and MC-LW, may have greater toxicity than MC-LR



Monitoring Challenges

- HBV of $0.04 \mu\text{g/L}$ is below LOD ($0.15 \mu\text{g/L}$)
 - ELISA is limited in accuracy, sensitivity and specificity
 - Congener-specific methods exist– LC/MS/MS

- Sampling – seasonal, diurnal, hourly variations

- Assumes MC-LR is most toxic and abundant variant – but may not be in all cases?

Questions?



Source: MPCA



Julia Dady

Phone: 651-201-4956

Julia.dady@state.mn.us

Useful links -

MDH – Microcystin-LR in Drinking Water

<http://www.health.state.mn.us/divs/eh/risk/guidance/gw/mclrinfo.pdf>

MDH – Little Rock Lake

<http://www.health.state.mn.us/divs/eh/hazardous/sites/benton/littlerocklake/index.html>

MPCA – Blue-green Algae and Harmful Algal Blooms

<http://www.pca.state.mn.us/index.php/water/water-types-and-programs/surface-water/lakes/lake-water-quality/blue-green-algae-and-harmful-algal-blooms.html?menuid=&redirect=1&expandable=1>

MPCA – National Lakes Assessment Project report:

<http://www.pca.state.mn.us/index.php/view-document.html?gid=6231>

Extra details, if questions...

Male repro

Chen et al. 2011

- Limitations and Uncertainties:
 - Mechanistic uncertainties – toxicokinetics, blood-testes barrier
 - Methodology uncertainties – no historical control data, low sperm motility in controls, sample handling and measurement

- MDH Conclusions on Repro Tox:
 - Uncertainties prevented use of Chen et al. study as a critical study
 - Repro uncertainty addressed in database uncertainty factor
 - MDH RfDs are 6-11 times lower than the LOAEL_{HED} from Chen study (i.e., considered protective)
 - Further research is needed to replicate and support findings

Extra details, if questions....

Other microcystin congeners

- ❑ Human Hepatocytes and HEK293 cells (Fischer et al. 2010):
 - MC-LW & LF - 7 to 70x greater cytotoxicity than MC-LR
 - Due to greater OATP receptor uptake into cells
- ❑ CHO cells (Huang et al. 2009) – cytotox LF > LW > LR
- ❑ HeLa cells (Monks et al. 2007) – cytotox: LR > LF > LW; but growth inhibition (IC₅₀ OATPs): LW > LF > LR
- ❑ Fischer et al. 2010 - Concluded that risk of human microcystin toxicity may be underestimated in algal blooms where MC-LW and MC-LF are predominant.