

SAT Initiative: Vonore Middle and Elementary Schools (Vonore, TN)

This document describes the analysis of air monitoring and other data collected under EPA's initiative to assess potentially elevated air toxics levels at some of our nation's schools. The document has been prepared for technical audiences (e.g., risk assessors, meteorologists) and their management. It is intended to describe the technical analysis of data collected for this school in clear, but generally technical, terms. A summary of this analysis is presented on the page focused on this school on EPA's website (www.epa.gov/schoolair).

I. Executive Summary

- Air monitoring has been conducted at Vonore Middle School, Vonore, Tennessee as part of the EPA initiative to monitor specific air toxics in the outdoor air around priority schools in 22 states and 2 tribal areas. For the purposes of this study, results from the air monitoring at Vonore Middle School are considered to be indicative of conditions at the nearby Vonore Elementary School (located within a half mile south of Vonore Middle School).
- These schools were selected for monitoring based on information indicating the potential for elevated ambient concentrations of diisocyanates in air outside the schools. The schools were ranked 26th on the USA Today list due to 2005 Toxics Release Inventory (TRI) estimates of diisocyanates emissions from a nearby boat manufacturer.
- Air monitoring was performed from September 4, 2009 to December 9, 2009 for hexamethylene diisocyanate (1,6-HDI), 4,4-methylene diphenyl diisocyanate (MDI), and 2,4-toluene diisocyanate (2,4-TDI).
- There was no detection of 1,6-HDI, MDI, or 2,4-TDI in any of the samples collected. However, information supplied by the Tennessee Department of Environment and Conservation (TDEC) indicates that, during the monitoring period, the nearby source was operating at a significantly lower level than normal. TDEC has determined that this facility only uses MDI and polymeric MDI as part of their normal operations.
- Based on the analysis described here, EPA plans to continue air toxics monitoring at this school for an additional 60 days when the nearby source's production returns to a more normal level.
- The TDEC will continue to oversee industrial facilities in the area through air permits and other programs. Through oversight of the facility, TDEC will determine when operations have returned to more normal levels and additional monitoring should be initiated.

II. Background on this Initiative

As part of an EPA initiative to implement Administrator Lisa Jackson's commitment to assess potentially elevated air toxics levels at some of our nation's schools, EPA and state and local air pollution control agencies are monitoring specific (key) air toxics in the outdoor air around priority schools in 22 states and 2 tribal areas (<http://www.epa.gov/schoolair/schools.html>).

- The schools selected for monitoring include some schools that are near large industries that are sources of air toxics, and some schools that are in urban areas, where emissions of air toxics come from a mix of large and small industries, cars, trucks, buses and other sources.
- EPA selected schools based on information available to us about air pollution in the vicinity of the school, including results of the 2002 National-Scale Air Toxics Assessment (NATA), results from a 2008 USA Today analysis on air toxics at schools, and information from state and local air agencies. The analysis by USA Today involved use of EPA's Risk Screening Environmental Indicators tool and Toxics Release Inventory (TRI) for 2005.
 - Available information had raised some questions about air quality near these schools that EPA concluded merited investigation. In many cases, the information indicated that estimated long-term average concentrations of one or more air toxics were above the upper end of the range that EPA generally considers as acceptable (e.g., above 1-in-10,000 cancer risk for carcinogens).
- Monitors are placed at each school for approximately 60 days, and take air samples on at least 10 different days during that time. The samples are analyzed for specific air toxics identified for monitoring at the school (i.e., key pollutants).¹
- These monitoring results and other information collected at each school during this initiative allow us to:
 - assess specific air toxics levels occurring at these sites and associated estimates of longer-term concentrations in light of health risk-based criteria for long-term exposures,
 - better understand, in many cases, potential contributions from nearby sources to key air toxics concentrations at the schools,
 - consider what next steps might be appropriate to better understand and address air toxics at the school, and
 - improve the information and methods we will use in the future (e.g., NATA) for estimating air toxics concentrations in communities across the U.S.

Assessment of air quality under this initiative is specific to the air toxics identified for monitoring at each school. This initiative is being implemented in addition to ongoing state, local and national air quality monitoring and assessment activities, including those focused on criteria pollutants (e.g., ozone and particulate matter) or existing, more extensive, air toxics programs.

Several technical documents prepared for this project provide further details on aspects of monitoring and data interpretation and are available on the EPA website (e.g., www.epa.gov/schoolair/techinfo.html). The full titles of these documents are provided here:

- *School Air Toxics Ambient Monitoring Plan*
- *Quality Assurance Project Plan For the EPA School Air Toxics Monitoring Program*
- *Schools Air Toxics Monitoring Activity (2009), Uses of Health Effects Information in Evaluating Sample Results*

¹ In analyzing air samples for these key pollutants, samples are also being analyzed for some additional pollutants that are routinely included in the analytical methods for the key pollutants.

Information on health effects of air toxics being monitored² and educational materials describing risk concepts³ are also available from EPA's website.

III. Basis for Selecting this School and the Air Monitoring Conducted

Vonore Middle School was selected for monitoring in consultation with the Tennessee air agency, Tennessee Department of Environment and Conservation (TDEC). For the purposes of this study, results from the air monitoring at Vonore Middle School are considered to be indicative of conditions at the nearby Vonore Elementary School (located within a half mile south of Vonore Middle School) (Figure 1). We were interested in evaluating the ambient concentrations of diisocyanates in air outside the schools because both Vonore schools were ranked 26th on the USA Today list due to 2005 TRI estimates of total diisocyanates for a nearby boat manufacturer. The diisocyanates that are HAPs are 1,6-HDI, MDI, and 2,4-TDI. TDEC has determined that this facility only uses MDI and polymeric MDI as part of their normal operations.

Monitoring commenced at Vonore Middle School on September 4, 2009 and continued through December 9, 2009. During this period, ten valid samples of diisocyanates were analyzed for 1,6-HDI, MDI, and 2,4-TDI. All sampling methodologies are described in EPA's schools air toxics monitoring plan (www.epa.gov/schoolair/techinfo.html).⁴

IV. Monitoring Results and Analysis

A. Background for the SAT Analysis

The majority of schools being monitored in this initiative were selected based on modeling analyses that indicated the potential for annual average air concentrations of some specific (key) hazardous air pollutants (HAPs or air toxics)⁵ to be of particular concern based on approaches that are commonly used in the air toxics program for considering potential for long-term risk. For example, such analyses suggested annual average concentrations of some air toxics were greater than long-term risk-based concentrations associated with an additional cancer risk greater than 10-in-10,000 or a hazard index on the order of or above 10. To make projections of air concentrations, the modeling analyses combined estimates of air toxics emissions from industrial, motor vehicle and other sources, with past measurements of winds, and other meteorological factors that can influence air concentrations, from a weather station in the general area. In some cases, the weather station was very close (within a few miles), but in other cases,

² For example, <http://www.epa.gov/schoolair/pollutants.html>, http://www.epa.gov/ttn/fera/risk_atoxic.html.

³ For example, http://www.epa.gov/ttn/atw/3_90_022.html, http://www.epa.gov/ttn/atw/3_90_024.html.

⁴ TDEC staff operated the monitors and sent the filters to the analytical laboratory under contract to EPA.

⁵ The term hazardous air pollutants (commonly called HAPs or air toxics) refers to pollutants identified in section 112(b) of the Clean Air Act which are the focus of regulatory actions involving stationary sources described by CAA section 112 and are distinguished from the six pollutants for which criteria and national ambient air quality standards (NAAQS) are developed as described in section 108. One of the criteria pollutants, lead, is also represented, as lead compounds, on the HAP list.

it was much further away (e.g., up to 60 miles), which may contribute to quite different conditions being modeled than actually exist at the school. The modeling analyses are intended to be used to prioritize locations for further investigation.

The primary objective of this initiative is to investigate - through monitoring air concentrations of key air toxics at each school over a 2-3 month period - whether levels measured and associated longer-term concentration estimates are of a magnitude, in light of health risk-based criteria, for which follow-up activities may need to be considered. To evaluate the monitoring results consistent with this objective, we developed health risk-based air concentrations (the long-term comparison levels summarized in Appendix A) for the monitored air toxics using established EPA methodology and practices for health risk assessment⁶ and, in the case of cancer risk, consistent with the implied level of risk considered in identifying schools for monitoring. Consistent with the long-term or chronic focus of the modeling analyses, based on which these schools were selected for monitoring, we have analyzed the full record of concentrations of air toxics measured at this school, using routine statistical tools, to derive a 95 percent confidence interval⁷ for the estimate of the longer-term average concentration of each of these pollutants. In this project, we are reporting all actual numerical values for pollutant concentrations including any values below method detection limit (MDL).⁸ Additionally, a value of 0.0 is used when a measured pollutant has no value detected (ND). The projected range for the longer-term concentration estimate for each chemical (most particularly the upper end of the range) is compared to the long-term comparison levels. These long-term comparison levels conservatively presume continuous (all-day, all-year) exposure over a lifetime. The analysis of the air concentrations also includes a consideration of the potential for cumulative multiple pollutant impacts.⁹ In general, where the monitoring results indicate estimates of longer-term average concentrations that are above the comparison levels - i.e., above the cancer-based comparison levels or notably above the noncancer-based comparison levels - we will consider the need for follow-up actions such as:

⁶ While this EPA initiative will rely on EPA methodology, practices, assessments and risk policy considerations, we recognize that individual state methods, practices and policies may differ and subsequent analyses of the monitoring data by state agencies may draw additional or varying conclusions.

⁷ When data are available for only a portion of the period of interest (e.g., samples not collected on every day during this period), statisticians commonly calculate the 95% confidence interval around the dataset mean (or average) in order to have a conservative idea of how high or low the “true” mean may be. More specifically, this interval is the range in which the mean for the complete period of interest is expected to fall 95% of the time (95% probability is commonly used by statisticians). The interval includes an equal amount of quantities above and below the sample dataset mean. The interval that includes these quantities is calculated using a formula that takes into account the size of the dataset (i.e., the ‘n’) as well as the amount by which the individual data values vary from the dataset mean (i.e., the “standard deviation”). This calculation yields larger confidence intervals for smaller datasets as well as ones with more variable data points. For example, a dataset including {1.0, 3.0, and 5.0}, results in a mean of 3.0 and a 95% confidence interval of 3.0 +/- ~5 (or -2.0 to 8.0). For comparison purposes, a dataset including {2.5, 3 and 3.5} results in a mean of 3.0 and a 95% confidence interval of 3.0 +/- ~1.2 (or 1.8 to 4.2). The smaller variation within the data in the second set of values causes the second confidence interval to be smaller.

⁸ Method detection limit (MDL) is the minimum concentration of a substance that can be measured and reported with 99% confidence that the pollutant concentration is greater than zero and is determined from the analysis of a sample in a given matrix containing the pollutant.

⁹ As this analysis of a 2-3 month monitoring dataset is not intended to be a full risk assessment, consideration of potential multiple pollutant impacts may differ among sites. For example, in instances where no individual pollutant appears to be present above its comparison level, we will also check for the presence of multiple pollutants at levels just below their respective comparison levels (giving a higher priority to such instances).

- Additional monitoring of air concentrations and/or meteorology in the area,
- Evaluation of potentially contributing sources to help us confirm their emissions and identify what options (regulatory and otherwise) may be available to us to achieve emissions reductions, and
- Evaluation of actions being taken or planned nationally, regionally or locally that may achieve emission and/or exposure reductions. An example of this would be the actions taken to address the type of ubiquitous emissions that come from mobile sources.

We have further analyzed the dataset to describe what it indicates in light of some other criteria and information commonly used in prioritizing state, local and national air toxics program activities. State, local and national programs often develop long-term monitoring datasets in order to better characterize pollutants near particular sources. The 2-3 month dataset developed under this initiative will be helpful to those programs in setting priorities for longer-term monitoring projects. The intent of this analysis is to make this 2-3 month monitoring dataset as useful as possible to state, local and national air toxics programs in their longer-term efforts to improve air quality nationally. To that end, this analysis:

- Describes the air toxics measurements in terms of potential longer-term concentrations, and, as available, compares the measurements at this school to monitoring data from national monitoring programs.
- Describes the meteorological data by considering conditions on sampling days as compared to those over all the days within the 2-3 month monitoring period and what conditions might be expected over the longer-term (as indicated, for example, by information from a nearby weather station).
- Describes available information regarding activities and emissions at the nearby source(s) of interest, such as that obtained from public databases such as TRI and/or consultation with the local air pollution authority.

B. Chemical Concentrations

We developed two types of long-term health risk-related comparison levels (summarized in Appendix A below) to address our primary objective. The primary objective is to investigate through the monitoring data collected for key pollutants at the school, whether pollutant levels measured and associated longer-term concentration estimates are elevated enough in comparison with health risk-based criteria to indicate that follow-up activities be considered. These comparison levels conservatively presume continuous (all-day, all-year) exposure over a lifetime.

In developing or identifying these comparison levels, we have given priority to use of relevant and appropriate air standards and EPA risk assessment guidance and precedents.¹⁰ These levels are based upon health effects information, exposure concentrations and risk estimates developed and assessed by EPA, the U.S. Agency for Toxic Substances and Disease Registry, and the California EPA. These agencies recognize the need to account for potential differences in

¹⁰ This is described in detail in *Schools Air Toxics Monitoring Activity (2009), Uses of Health Effects Information in Evaluating Sample Results*

sensitivity or susceptibility of different groups (e.g., asthmatics) or lifestages/ages (e.g., young children or the elderly) to a particular pollutant's effects so that the resulting comparison levels are relevant for these potentially sensitive groups as well as the broader population.

Using the analysis approach described above, we analyzed the chemical concentration data (Table 1) with regard to areas of interest identified below.

Key findings drawn from the information on chemical concentrations and the considerations discussed below include:

- None of the three diisocyanate compounds was detected in any of the samples collected. However, as the nearby source of diisocyanates was operating at a level that was significantly below normal during the monitoring period, additional monitoring will need to occur before a longer-term determination regarding potential health concerns can be made.

1,6-HDI, MDI, and 2,4-TDI, key pollutants:

- Do the monitoring data indicate elevated levels that pose significant long-term health concerns?
 - The data collected, which did not include any detection of these pollutants in any of the samples, does not indicate health concerns. However, as the nearby source of diisocyanates was operating at lower than normal levels during the monitoring, additional monitoring will need to occur before a longer-term determination regarding potential health concerns can be made.

C. Wind and Other Meteorological Data

At each school monitored as part of this initiative, we are collecting meteorological data, minimally for wind speed and direction, during the sampling period. Additionally, we have identified the nearest National Weather Service (NWS) station at which a longer record is available.

In reviewing these data at each school in this initiative, we are considering if these data indicate that the general pattern of winds on our sampling dates are significantly different from those occurring across the full sampling period or from those expected over the longer-term. Additionally, we are noting, particularly for school sites where the measured chemical concentrations show little indication of influence from a nearby source, whether wind conditions on some portion of the sampling dates were indicative of a potential to capture contributions from the nearby "key" source in the air sample collected.

The nearby key source of emissions was operating at lower than normal levels during the monitoring period. Therefore, this review of meteorological data serves mainly to provide information on wind patterns near the school during the sampling period and expected longer-

term patterns that may be used to characterize the meteorology once normal operations at the facility have resumed and more sampling is performed.

The meteorological station at Vonore Middle School collected wind speed and wind direction measurements during the sampling period, beginning on August 19, 2009, continuing through the sampling period (September 4, 2009-December 9, 2009), and ending on January 28, 2010. As a result, on-site data for these meteorological parameters are available for all dates of sample collection, and also for a period before and after the sampling period, producing a continuous record of five months of on-site meteorological data. The meteorological data collected at the school site on sampling days are presented in Figure 2 and Table 2. Additional meteorological data will be collected when sampling for diisocyanates resumes.

At the completion of the additional monitoring, all the meteorological data collected will be analyzed and compared to the nearest NWS station, which is at McGhee Tyson Airport in Knoxville, Tennessee. This station is approximately 20 miles northeast of the school. Measurements taken at that station include wind, temperature, and precipitation. These are presented in Table 2 and Appendix C.

Key findings drawn from this information and the considerations discussed below include:

- The wind patterns at the monitoring site across sampling dates are not similar to those observed across the record of on-site meteorological data during the sampling period.
- The wind patterns at the nearest NWS station during the sampling period are somewhat similar to those recorded at the NWS station over the long-term (2002-2007 period; Appendix C), supporting the idea that regional meteorological patterns in the area during the sampling period were consistent with long-term patterns.
- We will resume wind data collection when monitoring resumes at the school and analyze all the meteorological data together at the end of monitoring.

- What is the direction of the key source of diisocyanate emissions in relation to the school location?
 - The nearby boat manufacturer described in section III above lies approximately 1.3 miles northwest of the school.
 - Using the property boundaries of the full facility (in lieu of information regarding the location of specific sources of diisocyanates emissions at the facility), we have identified an approximate range of wind directions to use in considering the potential influence of this facility on air concentrations at the school.
 - This general range of wind directions, from approximately 280-325 degrees, is referred to here as the expected zone of source influence (ZOI).
- How often did wind come from direction of the key source?
 - There were eight days out of ten sampling days in which the on-site wind data had a portion of the winds were from the expected ZOI (Table 2).

- What might be expected over the longer-term at the school location?
 - The wind patterns at the nearest NWS station during the sampling period are somewhat similar to those recorded at the NWS station over the long-term (2002-2007 period; Appendix C), supporting the idea that regional meteorological patterns in the area during the sampling period were consistent with long-term patterns.

V. Key Source Information

- Was the source operating as usual during the monitoring period?
 - The nearby source of diisocyanates (described in section III above) has an operating permit issued by TDEC that includes operating requirements.¹¹
 - Information from the nearby source indicates that this facility was operating at a rate less than 20% of capacity during the sampling period, in comparison to its usual conditions of 50 to 60% capacity.
 - The most recently available diisocyanate emission reports for this source (2008 TRI) are less than 500 pounds per year. The 2007 TRI emissions report is about 50% higher (0.4 tons per year) than that relied upon in the previous modeling analysis for this area (e.g., 2005 TRI). TDEC has determined that this facility only uses MDI and polymeric MDI as part of their normal operations.

VI. Integrated Summary and Next Steps

A. Summary of Key Findings

1. What are the key HAPs for this school?
 - 1,6-HDI, MDI, and 2,4-TDI are the key HAPs for this school, identified based on emissions information considered in identifying the school for monitoring.
2. Do the data collected at this school indicate an elevated level of concern, as implied by information that led to identifying this school for monitoring?
 - There was no detection of any diisocyanates in any of the samples collected.

B. Next Steps for Key Pollutants

1. Based on the analysis described here, EPA plans to resume air toxics monitoring at this school when the nearby source's production returns to a more normal level.
2. The TDEC will continue its oversight of nearby facilities through the operating permit program. Through oversight of the facility, TDEC will determine when operations have returned to more normal levels and additional monitoring should be initiated.

¹¹ Operating permits, which are issued to air pollution sources under the Clean Air Act, are described at: <http://www.epa.gov/air/oaqps/permits/>

VII. Figures and Tables

A. Tables

1. Vonore Middle School – Key Pollutant Analysis.
2. Vonore Middle School Key Pollutant Concentrations and Meteorological Data.

B. Figures

1. Vonore Middle School and Vonore Elementary School.
2. Vonore Middle School (Vonore, TN) 1,6-HDI, MDI, and 2,4-TDI Concentration and Wind Information.

VIII. Appendices

- A. Summary Description of Long-term Comparison Levels.
- B. Vonore Middle School Pollutant Concentrations.
- C. Windroses for McGhee Tyson Airport NWS Station.

Figure 1. Vonore Middle School and Vonore Elementary School.



Table 1. Vonore Middle School - Key Pollutant Analysis.

Parameter	Units	Mean of Measurements	95% Confidence Interval on the Mean	Long-term Comparison Level ^a	
				Cancer-Based ^b	Noncancer-Based ^c
Hexamethylene Diisocyanate (1,6-HDI)	µg/m ³	100% of results were ND ^d		NA	0.01
Methylene Diphenyl Diisocyanate, 4,4- (MDI)	µg/m ³	100% of results were ND ^d		NA	0.6
Toluene Diisocyanate, 2,4- (2,4-TDI)	µg/m ³	100% of results were ND ^d		9.1	0.07

µg/m³ micrograms per cubic meter

ND No results of this chemical were registered by the laboratory analytical equipment.

NA Not applicable

^a Details regarding these values are in the technical report, Schools Air Toxics Monitoring Activity (2009) Uses of Health Effects Information.

^b Air toxics for which the upper 95% confidence limit on the mean concentration is above this level will be fully discussed in the text and may be considered a priority for potential follow-up activities, if indicated in light of the full set of information available for the site. Findings of the upper 95% confidence limit below 1% of the comparison level (i.e., where the upper 95% confidence limit is below the corresponding 1-in-1-million cancer risk based concentration) are generally considered a low priority for follow-up activity. Situations where the summary statistics for a pollutant are below this comparison level but above 1% of this level are fully discussed in the text of the report.

^c Air toxics for which the upper 95% confidence limit on the mean concentration are near or below the noncancer-based comparison level are generally of low concern and will generally be considered a low priority for follow-up activity. Pollutants for which the 95% confidence limits extend appreciably above the noncancer-based comparison level are fully discussed in the school-specific report and may be considered a priority for follow-up activity, if indicated in light of the full set of information available for the site.

^d There were no detections of this compound for any of the 10 valid samples collected.

Table 2. Vonore Middle School Key Pollutant Concentrations and Meteorological Data.

Parameter	Units	9/4/2009	9/10/2009	9/16/2009	9/22/2009	9/28/2009	10/4/2009	10/16/2009	10/22/2009	10/28/2009	12/9/2009
Hexamethylene Diisocyanate (1,6-HDI)	µg/m ³	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
Methylene Diphenyl Diisocyanate, 4,4- (MDI)	µg/m ³	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
Toluene Diisocyanate, 2,4- (2,4-TDI)	µg/m ³	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
% Hours w/Wind Direction from Expected ZOI ^a	%	12.5	20.8	0.0	16.7	8.3	0.0	37.5	4.2	16.7	4.2
Wind Speed (avg. of hourly speeds)	mph	2.0	2.6	2.5	2.1	5.4	2.2	4.1	2.5	2.7	11.3
Wind Direction (avg. of unitized vector) ^b	deg.	242.0	216.1	43.2	32.7	233.9	103.1	339.0	169.1	236.1	226.6
% of Hours with Speed below 2 knots	%	66.7	50.0	45.8	79.2	16.7	66.7	0.0	45.8	54.2	0
Daily Average Temperature	° F	73.1	69.0	71.1	73.9	67.2	56.4	51.0	56.9	60.6	52.8
Daily Precipitation	inches	0.00	0.01	0.16	0.03	0.00	0.27	0.06	0.00	0.01	1.07

All precipitation and temperature data were from the McGhee Tyson Airport NWS Station.

Selected wind speed and wind direction observations for December 9 were extracted from the McGhee Tyson Airport NWS Station due to questionably high wind speed readings (> 50mph) from the school site.



µg/m³ micrograms per cubic meter

ND No detection of this chemical was registered by the laboratory analytical equipment. The method detection limit for MDI and 2,4-TDI is 0.190 µg/m³, while the method detection limit for 1,6-HDI is 0.380 µg/m³.


^a Based on count of hours for which vector wind direction is from expected zone of influence.

^b Wind direction for each day is represented by values derived by scalar averaging of hourly estimates that were produced (by wind instrumentation's logger) as unitized vectors (specified as degrees from due north).

Figure 2. Vonore Middle School (Vonore, TN) 1,6-HDI, MDI, and 2,4-TDI Concentration and Wind Information.



**Vonore Middle School
Composite Hourly Windrose
on Sample Days
(Sep. 4, 2009-Dec. 9, 2009)**



**Vonore Middle School
Composite Hourly Windrose
Across Sampling Period
(Sep. 4, 2009-Dec. 9, 2009)**

KEY

Pollutant: 1,6-HDI, MDI, and 2,4-TDI
Timeframe: Septemeber 4, 2009 - December 9, 2009

Note

Each circle denotes a 24-hour collection of air for chemical analysis. The size of the circle indicates the magnitude of the wind speed for that day (wind data shown in Table 2). The expected zone of source influence is a rough approximation of the range of directions from which winds carrying chemicals emitted by the key source may originate.

- Wind Speed: 0.1-2.5 mph
- Wind Speed: 2.5-5.0 mph
- Wind Speed: > 5.0 mph

Expected Zone of Source Influence

Appendix A. Summary Description of Long-term Comparison Levels

In addressing the primary objective identified above, to investigate through the monitoring data collected for key pollutants at the school whether levels are of a magnitude, in light of health risk-based criteria, to indicate that follow-up activities be considered, we developed two types of long-term health risk-related comparison levels. These two types of levels are summarized below.¹²

Cancer-based Comparison Levels

- For air toxics where applicable, we developed cancer risk-based comparison levels to help us consider whether the monitoring data collected at the school indicate the potential for concentrations to pose incremental cancer risk above the range that EPA generally considers acceptable in regulatory decision-making to someone exposed to those concentrations continuously (24 hours a day, 7 days a week) over an entire lifetime.¹³ This general range is from 1 to 100 in a million.
- Air toxics with long-term mean concentrations below one one-hundredth of this comparison level would be below a comparably developed level for 1-in-a-million risk (which is the lower bound of EPA's traditional acceptable risk range). Such pollutants, with long-term mean concentrations below the Agency's traditional acceptable risk range, are generally considered to pose negligible risk.
- Air toxics with long-term mean concentrations above the acceptable risk range would generally be a priority for follow-up activities. In this evaluation, we compare the upper 95% confidence limit on the mean concentration to the comparison level. Pollutants for which this upper limit falls above the comparison level are fully discussed in the school monitoring report and may be considered a priority for potential follow-up activities in light of the full set of information available for that site.

¹² These comparison levels are described in more detail *Schools Air Toxics Monitoring Activity (2009), Uses of Health Effects Information in Evaluating Sample Results*.

¹³ While no one would be exposed at a school for 24 hours a day, every day for an entire lifetime, we chose this worst-case exposure period as a simplification for the basis of the comparison level in recognition of other uncertainties in the analysis. Use of continuous lifetime exposure yields a lower, more conservative, comparison level than would use of a characterization more specific to the school population (e.g., 5 days a week, 8-10 hours a day for a limited number of years).

Noncancer-based Comparison Levels

- To consider concentrations of air toxics other than lead (for which we have a national ambient air quality standard) with regard to potential for health effects other than cancer, we derived noncancer-based comparison levels using EPA chronic reference concentrations (or similar values). A chronic reference concentration (RfC) is an estimate of a long-term continuous exposure concentration (24 hours a day, every day) without appreciable risk of adverse effect over a lifetime.¹⁴ This differs from the cancer risk-based comparison level in that it represents a concentration without appreciable risk vs a risk-based concentration.
- In using this comparison level in this initiative, the upper end of the 95% confidence limit on the mean is compared to the comparison level. Air toxics for which this upper confidence limit is near or below the noncancer-based comparison level (i.e., those for which longer-term average concentration estimates are below a long-term health-related reference concentration) are generally of low concern and will generally be considered a low priority for follow-up activity. Pollutants for which the 95% confidence limits extend appreciably above the noncancer-based comparison level are fully discussed below and may be considered a priority for follow-up activity if indicated in light of the full set of information available for the pollutant and the site.
- For lead, we set the noncancer-based comparison level equal to the level of the recently revised national ambient air quality standard (NAAQS). It is important to note that the NAAQS for lead is a 3-month rolling average of lead in total suspended particles. Mean levels for the monitoring data collected in this initiative that indicate the potential for a 3-month average above the level of the standard will be considered a priority for consideration of follow-up actions such as siting of a NAAQS monitor in the area.

In developing or identifying these comparison levels, we have given priority to use of relevant and appropriate air standards and EPA risk assessment guidance and precedents. These levels are based upon health effects information, exposure concentrations and risk estimates developed and assessed by EPA, the U.S. Agency for Toxic Substances and Disease Registry, and the California EPA. These agencies recognize the need to account for potential differences in sensitivity or susceptibility of different groups (e.g., asthmatics) or lifestyles/ages (e.g., young children or the elderly) to a particular pollutant's effects so that the resulting comparison levels are relevant for these potentially sensitive groups as well as the broader population.

¹⁴ EPA defines the RfC as “an estimate (with uncertainty spanning perhaps an order of magnitude) of a continuous inhalation exposure to the human population (including sensitive subgroups) that is likely to be without an appreciable risk of deleterious effects during a lifetime. It can be derived from a NOAEL, LOAEL, or benchmark concentration, with uncertainty factors generally applied to reflect limitations of the data used. Generally used in EPA's noncancer health assessments.” http://www.epa.gov/ncea/iris/help_gloss.htm#r

Appendix B. Vonore Middle School Pollutant Concentrations.

Parameter	Units	9/4/2009	9/10/2009	9/16/2009	9/22/2009	9/28/2009	10/4/2009	10/16/2009	10/22/2009	10/28/2009	12/9/2009	Sample Screening Level ^a
Hexamethylene Diisocyanate (1,6-HDI)	µg/m ³	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	0.2
Methylene Diphenyl Diisocyanate, 4,4- (MDI)	µg/m ³	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	6.0
Toluene Diisocyanate, 2,4- (2,4-TDI)	µg/m ³	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	0.7



Key Pollutant

µg/m³ micrograms per cubic meter

ND No detection of this chemical was registered by the laboratory analytical equipment. The method detection limit for MDI, 2,4-TDI, and 2,6-TDI is 0.190 µg/m³, while the method detection limit for 1,6-HDI is 0.380 µg/m³.

^a The individual sample screening levels and their use is summarized on the web site and described in detail in Schools Air Toxics Monitoring Activity (2009), "Uses of Health Effects Information in Evaluating Sample Results", see <http://www.epa.gov/schoolair/pdfs/UsesOfHealthEffectsInfoinEvalSampleResults.pdf>. These screening levels are based on consideration of exposure all day, every day over a period ranging up to at least a couple of weeks, and longer for some pollutants.

Appendix C. Windroses for McGhee Tyson Airport NWS Station.



¹McGhee Tyson Airport NWS Station (WBAN 13891) is 20.42 miles from Vonore Middle School.